

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

A-level FURTHER MATHEMATICS

Paper 2

Thursday 6 June 2019

Afternoon

Time allowed: 2 hours

Materials

- You must have the AQA formulae and statistical tables booklet for A-level Mathematics and A-level Further Mathematics.
- You should have a scientific calculator that meets the requirements of the specification. (You may use a graphical calculator.)

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do **not** write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 100.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
TOTAL	



Answer **all** questions in the spaces provided.

- 1 Given that z is a complex number, and that z^* is the complex conjugate of z , which of the following statements is **not** always true?

Circle your answer.

[1 mark]

$(z^*)^* = z$ $zz^* = |z|^2$ $(-z)^* = -(z^*)$ $z - z^* = z^* - z$


$(x+iy) - (x-iy) \stackrel{?}{=} (x-iy) - (x+iy)$
 $2iy = -2iy \Rightarrow \text{only true if } y=0$

- 2 Which of the straight lines given below is an asymptote to the curve

$$y = \frac{ax^2}{x-1}$$

where a is a non-zero constant?

Circle your answer.



[1 mark]

$y = ax + a$ $y = ax$ $y = ax - a$ $y = a$

$y = mx + c$ is an asymptote if $\lim_{x \rightarrow \infty} [f(x) - (mx + b)] = 0$
 $\lim_{x \rightarrow \infty} ax + a + \frac{a}{x-1} - (ax + a) = 0$

- 3 The set \mathcal{A} is defined by $\mathcal{A} = \{x : -\sqrt{2} < x < 0\} \cup \{x : 0 < x < \sqrt{2}\}$

Which of the inequalities given below has \mathcal{A} as its solution?

Circle your answer.

[1 mark]

$|x^2 - 1| > 1$ $|x^2 - 1| \geq 1$ $|x^2 - 1| < 1$ $|x^2 - 1| \leq 1$

$x^2 - 1 = 1$ $-x^2 + 1 = 1$
 $x^2 - 2 = 0$ $x^2 = 0$
 $x = \pm\sqrt{2}$ $x = 0$

$\Rightarrow -\sqrt{2} < x < 0 \text{ \& } 0 < x < \sqrt{2}$



- 4 The positive integer k is such that

$$\sum_{r=1}^k (3r - k) = 90$$

Find the value of k .

[3 marks]

$$\sum_{r=1}^k (3r - k) = 3 \sum_{r=1}^k r - \sum_{r=1}^k k$$

$$= 3 \left(\frac{1}{2} k(k+1) \right) - k(k) \quad \text{using standard results}$$

$$= \frac{3}{2} k(k+1) - k^2$$

$$\Rightarrow \frac{1}{2} k^2 + \frac{3}{2} k = 90 \rightarrow k^2 + 3k - 180 = 0$$

$$(k+15)(k-12) = 0$$

k must be +ve so $k=12$

Turn over for the next question

Turn over ►



5 A curve has equation $y = \cosh x$

Show that the arc length of the curve from $x = a$ to $x = b$, where $0 < a < b$, is equal to

$$\sinh b - \sinh a$$

[4 marks]

$$S = \int_a^b \left(1 + \left(\frac{dy}{dx}\right)^2\right)^{\frac{1}{2}} dx \quad \text{standard result}$$

$$\frac{dy}{dx} = \sinh x$$

identity

$$1 + \left(\frac{dy}{dx}\right)^2 = 1 + \sinh^2 x = \cosh^2 x$$

$$\therefore S = \int_a^b (\cosh^2 x)^{\frac{1}{2}} dx = \int_a^b \cosh x dx$$

$$= [\sinh x]_a^b$$

$$= \sinh b - \sinh a$$




6

A circle C in the complex plane has equation $|z - 2 - 5i| = a$

The point z_1 on C has the least argument of any point on C , and $\arg(z_1) = \frac{\pi}{4}$

Prove that $a = \frac{3\sqrt{2}}{2}$

[6 marks]



$$\tan\left(\frac{\pi}{4}\right) = 1 = \frac{y}{x}$$

$$\arg(z_1) = \frac{\pi}{4} \Rightarrow z_1 \text{ is}$$

on line $y=x$.

let $z_1 = k+ki$, as $y=x$.

gradient of tangent = 1 \therefore gradient
of radius = -1

radius \perp tangent

gradients of
perpendicular lines
multiply to give -1

\therefore line between $(2,5)$ & (k,k) has gradient -1

$$\Rightarrow \frac{5-k}{2-k} = -1 \Rightarrow 5-k = k-2$$

$$2k = 7$$

$$k = \frac{7}{2}$$

equation of circle: $(x-x_1)^2 + (y-y_1)^2 = a^2$

sub in values $(2,5)$ & $(\frac{7}{2}, \frac{7}{2})$

$$a^2 = \left(5 - \frac{7}{2}\right)^2 + \left(2 - \frac{7}{2}\right)^2 = \left(\frac{3}{2}\right)^2 + \left(-\frac{3}{2}\right)^2 = 2 \times \frac{9}{4} = \frac{9}{2} = 4.5$$

$$a^2 = 4.5$$

$$\Rightarrow a = \sqrt{\frac{9}{2}} = \frac{3\sqrt{2}}{2} \quad (a \text{ is +ve root})$$

Turn over ►



7 The points A , B and C have coordinates $A(4, 5, 2)$, $B(-3, 2, -4)$ and $C(2, 6, 1)$

7 (a) Use a vector product to show that the area of triangle ABC is $\frac{5\sqrt{11}}{2}$

[4 marks]

$$\vec{AB} = \begin{pmatrix} -3 \\ 2 \\ -4 \end{pmatrix} - \begin{pmatrix} 4 \\ 5 \\ 2 \end{pmatrix} = \begin{pmatrix} -7 \\ -3 \\ -6 \end{pmatrix} \quad \vec{AC} = \begin{pmatrix} 2 \\ 6 \\ 1 \end{pmatrix} - \begin{pmatrix} 4 \\ 5 \\ 2 \end{pmatrix} = \begin{pmatrix} -2 \\ 1 \\ -1 \end{pmatrix}$$

$$\vec{AB} \times \vec{AC} = \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ -7 & -3 & -6 \\ -2 & 1 & -1 \end{vmatrix} = \underline{i}(3+6) - \underline{j}(7-12) + \underline{k}(-7-6)$$

$$= \begin{pmatrix} 9 \\ 5 \\ -13 \end{pmatrix}$$

$$\text{area} = \frac{1}{2} |\vec{AB} \times \vec{AC}| = \frac{\sqrt{9^2 + 5^2 + 13^2}}{2} = \frac{5\sqrt{11}}{2}$$

7 (b) The points A , B and C lie in a plane.

Find a vector equation of the plane in the form $\mathbf{r} \cdot \mathbf{n} = k$

[1 mark]

we need a vector that is \perp to the plane containing the points $A, B, \& C$. the cross product is $\perp \Rightarrow \mathbf{r} \cdot \begin{pmatrix} 9 \\ 5 \\ -13 \end{pmatrix} = k$

$$\mathbf{n} = 9\mathbf{i} + 5\mathbf{j} - 13\mathbf{k} \leftarrow$$

pick point in plane: $\begin{pmatrix} 4 \\ 5 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 9 \\ 5 \\ -13 \end{pmatrix} = 36 + 25 - 26 = 35$

$$\therefore \mathbf{r} \cdot \begin{pmatrix} 9 \\ 5 \\ -13 \end{pmatrix} = 35$$



7 (c) Hence find the exact distance of the plane from the origin.

[1 mark]

$$|\underline{n}| = 5\sqrt{11} \text{ magnitude of normal}$$

$\underline{r} \cdot \underline{n}$ gives projection of \underline{r} on \underline{n} so distance
is given by $\frac{\underline{r} \cdot \underline{n}}{|\underline{n}|}$

$$\frac{35}{5\sqrt{11}} = \frac{7}{\sqrt{11}} = \frac{7\sqrt{11}}{11}$$

Turn over for the next question

Turn over ►



8 A parabola P_1 has equation $y^2 = 4ax$ where $a > 0$

P_1 is translated by the vector $\begin{bmatrix} b \\ 0 \end{bmatrix}$, where $b > 0$, to give the parabola P_2

8 (a) The line $y = mx$ is a tangent to P_2

Prove that $m = \pm\sqrt{\frac{a}{b}}$

Solutions using differentiation will be given no marks.

[4 marks]

translation $\begin{bmatrix} b \\ 0 \end{bmatrix}$: $x \rightarrow x - b$

so parabola becomes $y^2 = 4a(x - b)$

tangent to P_2 : equate y 's so $m^2x^2 = 4a(x - b)$

$$m^2x^2 - 4ax + 4ab = 0$$

tangent \Rightarrow only one solution to above

equation \Rightarrow repeated roots

$$\therefore \Delta = 0 : 16a^2 - 4m^2(4ab) = 0$$

$$\underbrace{16a^2 - 4m^2(4ab)}_{b^2 - 4ac} = 0 \quad 16a^2 = 16m^2ab$$

$$m = \pm\sqrt{\frac{a}{b}}$$



8 (b) The line $y = \sqrt{\frac{a}{b}}x$ meets P_2 at the point D .

The finite region R is bounded by the x -axis, P_2 and a line through D perpendicular to the x -axis.

The region R is rotated through 2π radians about the x -axis to form a solid.

Find, in terms of a and b , the volume of this solid.

Fully justify your answer.

first sketch problem

then find coord of D

tangent meets P_2 :

$$\left(\sqrt{\frac{a}{b}}x\right)^2 = 4a(x-b)$$

$$ax^2 = 4ab(x-b)$$


$$x^2 - 4bx + 4b^2 = 0 \Rightarrow (x-2b)^2 = 0$$

$$\Rightarrow x = 2b$$

find where P_2 meets x -axis $0 = 4a(x-b) \therefore x = b$

So integration bounds for volume of revolution are $[b, 2b]$

$$\begin{aligned} V &= \pi \int_b^{2b} 4a(x-b) dx = 4a\pi \left[\frac{x^2}{2} - bx \right]_b^{2b} \\ &= 4a\pi \left\{ \left(\frac{4b^2}{2} - 2b^2 \right) - \left(\frac{b^2}{2} - b^2 \right) \right\} \\ &= 4a\pi \left\{ 0 - \left(-\frac{b^2}{2} \right) \right\} \\ &= 2ab^2\pi \end{aligned}$$



Turn over ►



- 9 (a) Find the eigenvalues and corresponding eigenvectors of the matrix

$$M = \begin{bmatrix} \frac{1}{5} & \frac{2}{5} \\ -\frac{3}{5} & \frac{13}{10} \end{bmatrix}$$

[5 marks]

eigenvalues: $\det(M - \lambda I) = 0$

$$\begin{vmatrix} \frac{1}{5} - \lambda & \frac{2}{5} \\ -\frac{3}{5} & \frac{13}{10} - \lambda \end{vmatrix} = \left(\frac{1}{5} - \lambda\right)\left(\frac{13}{10} - \lambda\right) + \frac{6}{25} \stackrel{\text{required}}{=} 0$$

$$\rightarrow \frac{13}{50} - \frac{1}{5}\lambda - \frac{13}{10}\lambda + \lambda^2 + \frac{6}{25} = 0$$

$$\frac{25}{50} - \frac{15}{10}\lambda + \lambda^2 = 0$$

$$\Rightarrow 0.5 - 1.5\lambda + \lambda^2 = 0$$

$$2\lambda^2 - 3\lambda + 1 = 0$$

$$(2\lambda - 1)(\lambda - 1) = 0 \therefore \lambda = 1 \text{ \& } \lambda = 0.5$$

eigenvectors: $M\underline{v} = \lambda\underline{v} \Rightarrow (M - \lambda)\underline{v} = 0$

$$\lambda = 1: \begin{pmatrix} -\frac{4}{5} & \frac{2}{5} \\ -\frac{3}{5} & \frac{3}{10} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 0 \Rightarrow 2y - 4x = 0 \quad (x \cdot 5)$$

$$y = 2x$$

$$\lambda = \frac{1}{2}: \begin{pmatrix} -\frac{3}{10} & \frac{2}{5} \\ -\frac{3}{5} & \frac{4}{5} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 0 \Rightarrow 8y - 6x = 0 \quad (x \cdot 10)$$

$$4y = 3x$$

$$\text{So } \lambda = 1, \underline{v} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}; \lambda = 0.5, \underline{v} = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$$

- 9 (b) Find matrices U and D such that D is a diagonal matrix and $M = UDU^{-1}$

[2 marks]

U is made of eigenvectors, $D = \text{diag}(\lambda_1, \lambda_2)$



$$\lambda=1 \quad \lambda=0.5$$

$$\text{So } U = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} \rightarrow D = \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{2} \end{pmatrix}$$

eigenvalues in order
corresponding to their eigenvectors given in U

Do not write
outside the
box

9 (c) Given that $M^n \rightarrow L$ as $n \rightarrow \infty$, find the matrix L .

$$\text{find } U^{-1} \text{ det } U = 3 - 8 = -5 \Rightarrow U^{-1} = \frac{-1}{5} \begin{pmatrix} 3 & -4 \\ -2 & 1 \end{pmatrix} \quad [4 \text{ marks}]$$

$$M = UDU^{-1} \Rightarrow M^n = U D^n U^{-1} \\ = \frac{-1}{5} \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 1^n & 0 \\ 0 & (\frac{1}{2})^n \end{pmatrix} \begin{pmatrix} 3 & -4 \\ -2 & 1 \end{pmatrix}$$

$$\therefore L = \lim_{n \rightarrow \infty} M^n = \frac{-1}{5} \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 3 & -4 \\ -2 & 1 \end{pmatrix} \\ = \frac{-1}{5} \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 3 & -4 \\ 0 & 0 \end{pmatrix} \\ = \frac{-1}{5} \begin{pmatrix} 3 & -4 \\ 6 & -8 \end{pmatrix} = \begin{pmatrix} -0.6 & 0.8 \\ -1.2 & 1.6 \end{pmatrix}$$

9 (d) The transformation represented by L maps all points onto a line.

Find the equation of this line.

[2 marks]

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = L \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\Rightarrow x' = -\frac{1}{5}(3x - 4y), \quad y' = -\frac{1}{5}(6x - 8y)$$

$$\therefore y = 2x$$

Turn over ►



10

Prove by induction that $f(n) = n^3 + 3n^2 + 8n$ is divisible by 6 for all integers $n \geq 1$ [7 marks]

$$\text{for } n=1, f(1) = 1^3 + 3(1)^2 + 8(1) = 12 = 2 \times 6$$

$f(1)$ is divisible by 6, so statement is true for $n=1$.

assume the statement is true for $n=k$, so $f(k) = 6m$ for some integer m .

$$\begin{aligned} \text{for } n=k+1: f(k+1) &= (k+1)^3 + 3(k+1)^2 + 8(k+1) \\ &= k^3 + 3k^2 + 3k + 1 + 3(k^2 + 2k + 1) \\ &\quad + 8(k+1) \\ &= k^3 + 6k^2 + 17k + 12 \end{aligned}$$

$$f(k+1) - f(k) = 3k^2 + 9k + 12$$

$$\therefore f(k+1) = 6m + 3k^2 + 9k + 12$$

$$3k^2 + 9k + 12 = 3(k^2 + 3k + 4) \therefore \text{is a multiple of 3.}$$

$$3k^2 + 9k + 12 = 3(k^2 + 3k) + 12$$

$$k^2 + 3k = k(k+3) \rightarrow \text{either } k \text{ or } k+3 \text{ is even}$$

(difference is 3 so can't both be even).

even \times odd = even, so $k^2 + 3k$ is even, 4 is even

$$\therefore 3(k^2 + 3k + 4) \text{ is an even multiple of 3}$$

(divisible by 3 & 2) so is divisible by 6. thus, if $f(k)$ is divisible by 6, $f(k+1)$ is a sum of multiples of 6, hence is also a multiple of 6.

so, if we assume the statement is true for $f(k)$, it's true for $f(k+1)$. as it is true for $f(1)$, so by induction, we conclude it is true for all $k \in \mathbb{Z}^+$



Turn over for the next question

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Turn over ►



11

The line L_1 has equation

$$\frac{x-2}{3} = \frac{y+4}{8} = \frac{4z-5}{5}$$

The line L_2 has equation

$$\left(\mathbf{r} - \begin{bmatrix} -2 \\ 0 \\ 3 \end{bmatrix} \right) \times \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \mathbf{0}$$

Find the shortest distance between the two lines, giving your answer to three significant figures.

[8 marks]

$$\lambda = \frac{x-2}{3} = \frac{y+4}{8} = \frac{4z-5}{5} \Rightarrow \text{Vector for } L_1$$

$$\text{given by } \mathbf{r}_1 = \begin{pmatrix} 2 \\ -4 \\ 5/4 \end{pmatrix} + \lambda \begin{pmatrix} 3 \\ 8 \\ 5/4 \end{pmatrix} = \begin{pmatrix} 2 \\ -4 \\ 5/4 \end{pmatrix} + \lambda \begin{pmatrix} 12 \\ 32 \\ 5 \end{pmatrix}$$

$$L_2 \text{ vector: } \mathbf{r}_2 = \begin{pmatrix} -2 \\ 0 \\ 3 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}$$

now we have vector equations for both lines, we find the vector between them & the vector \perp to both

$$L_1 \rightarrow L_2: \begin{pmatrix} 2 \\ -4 \\ 5/4 \end{pmatrix} - \begin{pmatrix} -2 \\ 0 \\ 3 \end{pmatrix} = \begin{pmatrix} 4 \\ -4 \\ -7/4 \end{pmatrix}$$

$$\begin{aligned} \underline{n} &= \begin{pmatrix} 12 \\ 32 \\ 5 \end{pmatrix} \times \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 12 & 32 & 5 \\ 2 & 1 & 3 \end{vmatrix} = \mathbf{i}(96-5) - \mathbf{j}(36-10) + \mathbf{k}(12-64) \\ &= \begin{pmatrix} 91 \\ -26 \\ -52 \end{pmatrix} \end{aligned}$$



$$|n| = \sqrt{91^2 + 26^2 + 52^2} = \sqrt{11661} = 13\sqrt{69}$$

project ($L_1 \rightarrow L_2$) on $|n|$ & divide by length of n

$$\frac{\begin{pmatrix} 4 \\ -4 \\ -\frac{7}{4} \\ 4 \end{pmatrix} \cdot \begin{pmatrix} 91 \\ -26 \\ -52 \end{pmatrix}}{13\sqrt{69}} = \frac{43}{\sqrt{69}} = 5.17659\dots = 5.18 \text{ (3sf.)}$$

Turn over for the next question

Turn over ►



12 Abel and Bonnie are trying to solve this mathematical problem:

$$z = 2 - 3i \text{ is a root of the equation}$$

$$2z^3 + mz^2 + pz + 91 = 0$$

Find the value of m and the value of p .

Abel says he has solved the problem.

Bonnie says there is not enough information to solve the problem.

12 (a) Abel's solution begins as follows:

Since $z = 2 - 3i$ is a root of the equation,
 $z = 2 + 3i$ is another root.

State **one extra** piece of information about m and p which could be added to the problem to make the beginning of Abel's solution correct.

[1 mark]

m & p are real numbers



12 (b)

Prove that Bonnie is right.

[4 marks]

$$z^3 + \frac{m}{2}z^2 + \frac{p}{2}z + \frac{q}{2} = 0$$

$$\text{So product of roots} = -\frac{q}{2}$$

$$\text{as } (2-3i) \text{ is one root; } (2-3i)\beta\gamma = -\frac{q}{2}$$

$$\begin{aligned} \frac{-q}{2} \div (2-3i) &= \frac{-q}{2(2-3i)} \times \frac{2+3i}{2+3i} = \frac{-q(2+3i)}{2(4+9)} \\ &= \frac{-q(2+3i)}{26} \\ &= \frac{-7(2+3i)}{2} \end{aligned}$$

so roots could be $(2+3i)$ & $-\frac{7}{2}$,

$$\text{giving } m = -2 \times \left(2+3i + 2-3i - \frac{7}{2}\right)^2 = -1 \text{ \&}$$

$$p = +2 \times \left(4+9 + (2+3i)\frac{2-\frac{7}{2}}{2} + (2-3i)\frac{-\frac{7}{2}}{2}\right) = -2$$

but roots could also be $(4+6i)$ & $-\frac{7}{4}$,

$$\text{giving } m = -2 \times \left(2-3i + 4+6i - \frac{7}{4}\right)^2 \neq -1$$

↳ different values for m & p .

there is more than one set of possible values for m & p , so we don't have enough info to solve the problem.
∴ Bonnie is right.

Turn over ►



- 13 (a) Explain why $\int_3^{\infty} x^2 e^{-2x} dx$ is an improper integral.

[1 mark]

the upper limit is infinity, making it improper.

- 13 (b) Evaluate $\int_3^{\infty} x^2 e^{-2x} dx$

Show the limiting process.

[9 marks]

start by calculating the integral without the limits.
use integration by parts: $u = x^2$ $v' = e^{-2x}$
 $u' = 2x$ $v = -\frac{e^{-2x}}{2}$

$$\rightarrow \int x^2 e^{-2x} dx = -\frac{x^2 e^{-2x}}{2} + \int x e^{-2x} dx$$

if we differentiate this, we get 1, leaving us with an integral we can compute

integrate by parts again:

$$u = x \quad v' = e^{-2x}$$

$$u' = 1 \quad v = -\frac{e^{-2x}}{2}$$

$$\int x e^{-2x} dx = \left(-\frac{x e^{-2x}}{2} \right) + \int \frac{e^{-2x}}{2} dx = -\frac{x e^{-2x}}{2} - \frac{e^{-2x}}{4}$$

$$\therefore \int x^2 e^{-2x} dx = -\frac{x^2 e^{-2x}}{2} - \frac{x e^{-2x}}{2} - \frac{e^{-2x}}{4}$$

$$\int_3^{\infty} x^2 e^{-2x} dx = \lim_{n \rightarrow \infty} \int_3^n x^2 e^{-2x} dx$$

change to finite limit for proper integral



$$= \lim_{n \rightarrow \infty} \left\{ \left[\frac{-x^2 e^{-2x}}{2} - \frac{x e^{-2x}}{2} - \frac{e^{-2x}}{4} \right]_3^n \right\}$$

$n \rightarrow \infty: e^{-2n} \rightarrow 0$, this dominates over the increase of $x \rightarrow \infty$.

$$\hookrightarrow \int_3^{\infty} x^2 e^{-2x} dx = \frac{9e^{-6}}{2} + \frac{3e^{-6}}{2} + \frac{e^{-6}}{4} = \frac{25e^{-6}}{4}$$

Turn over for the next question

Turn over ►



14 Let

$$S_n = \sum_{r=1}^n \frac{1}{(r+1)(r+3)}$$

where $n \geq 1$

14 (a) Use the method of differences to show that

$$S_n = \frac{5n^2 + an}{12(n+b)(n+c)}$$

where a , b and c are integers.

[6 marks]

use partial fractions: $\frac{1}{(r+1)(r+3)} = \frac{A}{r+1} + \frac{B}{r+3}$

$$1 = A(r+3) + B(r+1)$$

$$\left. \begin{array}{l} r=-3: 1 = -2B \Rightarrow B = -\frac{1}{2} \\ r=-1: 1 = 2A \Rightarrow A = \frac{1}{2} \end{array} \right\} \frac{1}{2(r+1)} - \frac{1}{2(r+3)} = \frac{1}{(r+1)(r+3)}$$

$$\therefore 2S_n = \sum_{r=1}^n \frac{1}{r+1} - \frac{1}{r+3} = \frac{1}{2} - \frac{1}{4} + \frac{1}{3} - \frac{1}{5} + \frac{1}{4} - \frac{1}{6} + \dots$$

$$+ \frac{1}{n-1} - \frac{1}{n+1} + \frac{1}{n} - \frac{1}{n+2} + \frac{1}{n+1}$$

$$+ \frac{1}{n+3}$$

$$= \frac{1}{2} + \frac{1}{3} - \frac{1}{n+2} - \frac{1}{n+3}$$

$$= \frac{3(n+2)(n+3) + 2(n+2)(n+3)}{2 \times 3 \times (n+2)(n+3)} - \frac{3 \times 2(n+3)}{2 \times 3 \times (n+2)(n+3)} - \frac{2 \times 3(n+2)}{2 \times 3 \times (n+2)(n+3)}$$

$$= \frac{5(n+2)(n+3) - 6(n+3) - 6(n+2)}{6(n+2)(n+3)}$$



$$\Rightarrow S_n = \frac{5(n^2 + 5n + 6) - 6(2n + 5)}{12(n+3)(n+2)}$$
$$= \frac{5n^2 + 13n}{12(n+2)(n+3)}$$

Question 14 continues on the next page

Turn over ►



- 14 (b) Show that, for any number k greater than $\frac{12}{5}$, if the difference between $\frac{5}{12}$ and S_n is less than $\frac{1}{k}$, then

$$n > \frac{k - 5 + \sqrt{k^2 + 1}}{2}$$

difference between $\frac{5}{12}$ & $S_n < \frac{1}{k}$: $\frac{5}{12} - \frac{(5n^2 + 13n)}{12(n+2)(n+3)} < \frac{1}{k}$ [6 marks]

$$\Rightarrow \frac{5(n+2)(n+3) - (5n^2 + 13n)}{12(n+2)(n+3)} < \frac{1}{k}$$

$$\frac{12n + 30}{12(n+2)(n+3)} < \frac{1}{k}$$

both denominators > 0 so $k(12n + 30) < 12(n+2)(n+3)$

$$\Rightarrow 12n^2 + (60 - 12k)n + (72 - 30k) > 0$$

$$\div 6: 2n^2 + (10 - 2k)n + (12 - 5k) > 0$$

$k > \frac{12}{5}$ so $(12 - 5k) < 0 \Rightarrow$ above equation has a +ve & -ve root

$$\text{+ve root: } \frac{(2k - 10) + \sqrt{(10 - 2k)^2 - 8(12 - 5k)}}{4} =$$


$$\frac{(2k - 10) + \sqrt{(100 + 4k^2 - 40) - 96 + 40k}}{4} =$$

$$\frac{(2k - 10) + \sqrt{4k^2 + 4}}{4} = \frac{2(k - 5) + 2\sqrt{k^2 + 1}}{4}$$



$$= \frac{(k-5) + \sqrt{k^2+1}}{2}$$

as the root is +ve, if $12n^2 + (60-12k)n + (72-30k)$
is > 0 , then $n > \frac{(k-5) + \sqrt{k^2+1}}{2}$




Turn over for the next question

Turn over ►



15



Two tanks, A and B , each have a capacity of 800 litres.

At time $t = 0$ both tanks are full of pure water.

When $t > 0$, water flows in the following ways:

- Water with a salt concentration of μ grams per litre flows into tank A at a constant rate
- Water flows from tank A to tank B at a rate of 16 litres per minute
- Water flows from tank B to tank A at a rate of r litres per minute (b)
- Water flows out of tank B through a waste pipe
- The amount of water in each tank remains at 800 litres.

At time t minutes ($t \geq 0$) there are x grams of salt in tank A and y grams of salt in tank B .

This system is represented by the coupled differential equations

$$\frac{dx}{dt} = \underline{36} - 0.02x + 0.005y \quad (1)$$

$$\frac{dy}{dt} = 0.02x - 0.02y \quad (2)$$

15 (a) Find the value of r .

[2 marks]

from B to A \Rightarrow +ve y coefficient in $\frac{dx}{dt}$

$$r = 0.005 \times 800 = 4$$

\uparrow capacity of tanks



15 (b) Show that $\mu = 3$

[3 marks]

constant rate : flow into A = flow out of A

$$16 - 4 = 12 \text{ L/min} \Rightarrow \text{carries } 12 \mu \text{g/min}$$

36 is constant value in $\frac{dx}{dt}$ so $12\mu = 36$

$$\therefore \mu = 3$$

15 (c) Solve the coupled differential equations to find both x and y in terms of t .

[9 marks]

differentiate (2): $\frac{d^2y}{dt^2} = 0.02\dot{x} - 0.02\dot{y}$

$$0.02\dot{x} = \ddot{y} + 0.02\dot{y}$$

$$x = 50\dot{y} + y \quad \& \quad \dot{x} = 50\ddot{y} + \dot{y}$$

sub in (1): $50\ddot{y} + \dot{y} = 36 - 0.02(50\dot{y} + y) + 0.005y$

$$50\ddot{y} + 2\dot{y} + 0.015y = 36$$

$$\text{CF: } 50m^2 + 2m + 0.015 = 0 \Rightarrow m = \frac{-2 \pm \sqrt{2^2 - 4(50)(0.015)}}{2(50)}$$

$$= \frac{-2 \pm \sqrt{1}}{100}$$

$$\Rightarrow m = -\frac{1}{100} \quad \& \quad m = -\frac{3}{100}$$

Turn over ►



PI: let $y = \lambda \Rightarrow \dot{y} = \ddot{y} = 0$. sub in: $0.015\lambda = 36$
 $\lambda = 2400$
same form as RHS

$$\therefore y = Ae^{-0.03t} + Be^{-0.01t} + 2400$$

$x = 50\dot{y} + y$ so we need \dot{y} :

$$\dot{y} = -0.03Ae^{-0.03t} - 0.01Be^{-0.01t}$$

$$\Rightarrow x = -0.5Ae^{-0.03t} + 0.5Be^{-0.01t} + 2400$$

initial conditions: @ $t=0$, $x=0$ & $y=0$

$$\Rightarrow A + B + 2400 = 0 \quad (y) \quad \textcircled{1}$$

$$-0.5A + 0.5B + 2400 = 0 \quad (x) \quad \textcircled{2}$$

$$\textcircled{1} + 2 \times \textcircled{2}: 2B + 7200 = 0$$

$$B = -3600$$

$$A - 3600 + 2400 = 0$$

$$A = 1200$$

$$\text{so } x = -600e^{-0.03t} - 1800e^{-0.01t} + 2400$$

$$y = 1200e^{-0.03t} - 3600e^{-0.01t} + 2400$$

END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



Do not write
outside the
box

Question number	Additional page, if required. Write the question numbers in the left-hand margin.
	<hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/> <hr/>
	<p>Copyright information</p> <p>For confidentiality purposes, from the November 2015 examination series, acknowledgements of third-party copyright material are published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.</p> <p>Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.</p> <p>Copyright © 2019 AQA and its licensors. All rights reserved.</p>

