www.mymathscloud.com

General Certificate of Education January 2008 Advanced Level Examination

## ASSESSMENT and QUALIFICATIONS ALLIANCE

## MATHEMATICS Unit Pure Core 3

MPC3

Thursday 17 January 2008 1.30 pm to 3.00 pm

## For this paper you must have:

- an 8-page answer book
- the blue AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed: 1 hour 30 minutes

## **Instructions**

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MPC3.
- Answer all questions.
- Show all necessary working; otherwise marks for method may be lost.

### **Information**

- The maximum mark for this paper is 75.
- The marks for questions are shown in brackets.

### Advice

• Unless stated otherwise, you may quote formulae, without proof, from the booklet.

P2192/Jan08/MPC3 6/6/6/6/ MPC3

# www.my.mainscloud.com

## Answer all questions.

1 (a) Find  $\frac{dy}{dx}$  when:

(i) 
$$y = (2x^2 - 5x + 1)^{20}$$
;

(2 marks)

(ii) 
$$y = x \cos x$$
.

(2 marks)

(b) Given that

$$y = \frac{x^3}{x - 2}$$

show that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{kx^2(x-3)}{(x-2)^2}$$

where k is a positive integer.

(3 marks)

- 2 (a) Solve the equation  $\cot x = 2$ , giving all values of x in the interval  $0 \le x \le 2\pi$  in radians to two decimal places. (2 marks)
  - (b) Show that the equation  $\csc^2 x = \frac{3 \cot x + 4}{2}$  can be written as

$$2\cot^2 x - 3\cot x - 2 = 0 \tag{2 marks}$$

(c) Solve the equation  $\csc^2 x = \frac{3 \cot x + 4}{2}$ , giving all values of x in the interval  $0 \le x \le 2\pi$  in radians to two decimal places. (4 marks)

# MANN. MY MAINS COUNT. COM

## 3 The equation

$$x + (1 + 3x)^{\frac{1}{4}} = 0$$

has a single root,  $\alpha$ .

- (a) Show that  $\alpha$  lies between -0.33 and -0.32. (2 marks)
- (b) Show that the equation  $x + (1 + 3x)^{\frac{1}{4}} = 0$  can be rearranged into the form

$$x = \frac{1}{3}(x^4 - 1)$$
 (2 marks)

- (c) Use the iteration  $x_{n+1} = \frac{(x_n^4 1)}{3}$  with  $x_1 = -0.3$  to find  $x_4$ , giving your answer to three significant figures. (3 marks)
- 4 The functions f and g are defined with their respective domains by

$$f(x) = x^3$$
, for all real values of  $x$   
 $g(x) = \frac{1}{x-3}$ , for real values of  $x, x \neq 3$ 

- (a) State the range of f. (1 mark)
- (b) (i) Find fg(x). (1 mark)
  - (ii) Solve the equation fg(x) = 64. (3 marks)
- (c) (i) The inverse of g is  $g^{-1}$ . Find  $g^{-1}(x)$ . (3 marks)
  - (ii) State the range of  $g^{-1}$ . (1 mark)
- 5 (a) (i) Given that  $y = 2x^2 8x + 3$ , find  $\frac{dy}{dx}$ . (1 mark)
  - (ii) Hence, or otherwise, find

$$\int_{4}^{6} \frac{x-2}{2x^2 - 8x + 3} \, \mathrm{d}x$$

giving your answer in the form  $k \ln 3$ , where k is a rational number. (4 marks)

(b) Use the substitution u = 3x - 1 to find  $\int x\sqrt{3x - 1} \, dx$ , giving your answer in terms of x.

www.mymathscloud.com

Sketch the curve with equation  $y = \csc x$  for  $0 < x < \pi$ . 6

- Use the mid-ordinate rule with four strips to find an estimate for  $\int_{0.1}^{0.5} \csc x \, dx$ , giving your answer to three significant figures. (4 marks)
- Describe a sequence of **two** geometrical transformations that maps the graph of  $y = x^2$ 7 onto the graph of  $v = 4x^2 - 5$ . (4 marks)
  - Sketch the graph of  $y = |4x^2 5|$ , indicating the coordinates of the point where the curve crosses the *y*-axis. (3 marks)
  - Solve the equation  $|4x^2 5| = 4$ . (3 marks) (c)
    - Hence, or otherwise, solve the inequality  $|4x^2 5| \ge 4$ . (2 marks)
- Given that  $e^{-2x} = 3$ , find the exact value of x. 8 (2 marks)
  - Use integration by parts to find  $\int xe^{-2x} dx$ . (4 marks)
  - A curve has equation  $y = e^{-2x} + 6x$ .
    - Find the exact values of the coordinates of the stationary point of the curve. (4 marks)
    - (2 marks) Determine the nature of the stationary point.
    - The region R is bounded by the curve  $y = e^{-2x} + 6x$ , the x-axis and the lines x = 0 and x = 1.

Find the volume of the solid formed when R is rotated through  $2\pi$  radians about the x-axis, giving your answer to three significant figures. (5 marks)

## **END OF QUESTIONS**