

Centre Number						Candidate Number					
Surname											
Other Names											
Candidate Signature											

For Examiner's Use	
Examiner's Initials	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
TOTAL	



General Certificate of Education
Advanced Level Examination
June 2014

Mathematics

MPC3

Unit Pure Core 3

Tuesday 10 June 2014 9.00 am to 10.30 am

For this paper you must have:

- the blue AQA booklet of formulae and statistical tables.
- You may use a graphics calculator.

Time allowed

- 1 hour 30 minutes

- Instructions**
- Use black ink or black ball-point pen. Pencil should only be used for drawing.
 - Fill in the boxes at the top of this page.
 - Answer **all** questions.
 - Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
 - You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
 - Do not write outside the box around each page.
 - Show all necessary working; otherwise marks for method may be lost.
 - Do all rough work in this book. Cross through any work that you do not want to be marked.

- Information**
- The marks for questions are shown in brackets.
 - The maximum mark for this paper is 75.

- Advice**
- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
 - You do not necessarily need to use all the space provided.



Answer **all** questions.

Answer each question in the space provided for that question.

- 1 Use Simpson's rule, with five ordinates (four strips), to calculate an estimate for

$$\int_0^{\pi} x^{\frac{1}{2}} \sin x \, dx$$

Give your answer to four significant figures.

[4 marks]

QUESTION
PART
REFERENCE

Answer space for question 1



QUESTION
PART
REFERENCE

Answer space for question 1

A large rectangular area with horizontal dotted lines for writing the answer.

Turn over ►



2 A curve has equation $y = 2 \ln(2e - x)$.

(a) Find $\frac{dy}{dx}$. [2 marks]

(b) Find an equation of the normal to the curve $y = 2 \ln(2e - x)$ at the point on the curve where $x = e$. [4 marks]

(c) The curve $y = 2 \ln(2e - x)$ intersects the line $y = x$ at a single point, where $x = \alpha$.

(i) Show that α lies between 1 and 3. [2 marks]

(ii) Use the recurrence relation

$$x_{n+1} = 2 \ln(2e - x_n)$$

with $x_1 = 1$ to find the values of x_2 and x_3 , giving your answers to three decimal places. [2 marks]

(iii) **Figure 1**, on the opposite page, shows a sketch of parts of the graphs of $y = 2 \ln(2e - x)$ and $y = x$, and the position of x_1 .

On **Figure 1**, draw a cobweb or staircase diagram to show how convergence takes place, indicating the positions of x_2 and x_3 on the x -axis. [2 marks]

QUESTION
PART
REFERENCE

Answer space for question 2

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....




QUESTION
PART
REFERENCE

Answer space for question 2

(c)(iii)

Figure 1



Turn over ►



3 (a) (i) Differentiate $(x^2 + 1)^{\frac{5}{2}}$ with respect to x .

[2 marks]

(ii) Given that $y = e^{2x}(x^2 + 1)^{\frac{5}{2}}$, find the value of $\frac{dy}{dx}$ when $x = 0$.

[3 marks]

(b) A curve has equation $y = \frac{4x - 3}{x^2 + 1}$. Use the quotient rule to find the x -coordinates of the stationary points of the curve.

[5 marks]

QUESTION
PART
REFERENCE

Answer space for question 3



QUESTION
PART
REFERENCE


Answer space for question 3

A large rectangular area with horizontal dotted lines for writing the answer.



Turn over ►

- 4 The sketch shows part of the curve with equation $y = f(x)$.




- (a) On **Figure 2** below, sketch the curve with equation $y = -|f(x)|$. **[3 marks]**
- (b) On **Figure 3** on the page opposite, sketch the curve with equation $y = f(|2x|)$. **[2 marks]**
- (c) (i) Describe a sequence of two geometrical transformations that maps the graph of $y = f(x)$ onto the graph of $y = f(2x + 2)$. **[4 marks]**
- (ii) Find the coordinates of the image of the point $P(4, -3)$ under the sequence of transformations given in part (c)(i). **[2 marks]**

QUESTION
PART
REFERENCE

Answer space for question 4

(a)

Figure 2




QUESTION
PART
REFERENCE

Answer space for question 4

(b)

Figure 3



Area with horizontal dotted lines for writing answers.

Turn over ►



5 The functions f and g are defined with their respective domains by

$$f(x) = x^2 - 6x + 5, \quad \text{for } x \geq 3$$

$$g(x) = |x - 6|, \quad \text{for all real values of } x$$

(a) Find the range of f . **[2 marks]**

(b) The inverse of f is f^{-1} .
 Find $f^{-1}(x)$. Give your answer in its simplest form. **[4 marks]**

(c) (i) Find $gf(x)$. **[1 mark]**

(ii) Solve the equation $gf(x) = 6$. **[4 marks]**

QUESTION
PART
REFERENCE

Answer space for question 5



QUESTION
PART
REFERENCE

Answer space for question 5

A large rectangular area with horizontal dotted lines for writing the answer to question 5.

Turn over ►



6 (a) By using integration by parts twice, find

$$\int x^2 \sin 2x \, dx$$

[6 marks]

(b) A curve has equation $y = x\sqrt{\sin 2x}$, for $0 \leq x \leq \frac{\pi}{2}$.

The region bounded by the curve and the x -axis is rotated through 2π radians about the x -axis to generate a solid.

Find the exact value of the volume of the solid generated.

[3 marks]

QUESTION
PART
REFERENCE

Answer space for question 6



QUESTION
PART
REFERENCE

Answer space for question 6

Area with horizontal dotted lines for writing the answer.

Turn over ►



7 Use the substitution $u = 3 - x^3$ to find the exact value of $\int_0^1 \frac{x^5}{3 - x^3} dx$.

[6 marks]

QUESTION
PART
REFERENCE

Answer space for question 7



QUESTION
PART
REFERENCE

Answer space for question 7

A large rectangular area with horizontal dotted lines for writing the answer to question 7.

Turn over ►



8 (a) Show that the expression $\frac{1 - \sin x}{\cos x} + \frac{\cos x}{1 - \sin x}$ can be written as $2 \sec x$. [4 marks]

(b) Hence solve the equation

$$\frac{1 - \sin x}{\cos x} + \frac{\cos x}{1 - \sin x} = \tan^2 x - 2$$

giving the values of x to the nearest degree in the interval $0^\circ \leq x < 360^\circ$. [6 marks]

(c) Hence solve the equation

$$\frac{1 - \sin(2\theta - 30^\circ)}{\cos(2\theta - 30^\circ)} + \frac{\cos(2\theta - 30^\circ)}{1 - \sin(2\theta - 30^\circ)} = \tan^2(2\theta - 30^\circ) - 2$$

giving the values of θ to the nearest degree in the interval $0^\circ \leq \theta \leq 180^\circ$. [2 marks]

QUESTION
PART
REFERENCE

Answer space for question 8

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



QUESTION
PART
REFERENCE

Answer space for question 8

A large rectangular area with horizontal dotted lines for writing the answer to question 8.



Turn over ►

QUESTION
PART
REFERENCE

Answer space for question 8

A large rectangular area with horizontal dotted lines for writing the answer to question 8.



QUESTION
PART
REFERENCE

Answer space for question 8

A large rectangular area with horizontal dotted lines for writing the answer to question 8.

END OF QUESTIONS



There are no questions printed on this page

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

