

Further Pure 1 - January 2008

① $z = 2 + i$ $z^x = 2 - i$
 $xc + 3iy = 2 + i + 4i(2 - i)$
 $= 2 + i + 8i + 4$
 $xc + 3iy = 6 + 9i$

Real: $xc = 6$

Imaginary: $3y = 9 \Rightarrow y = 3$

② $d^2y/dx^2 = 2^x$ $y_{n+1} = y_n + h\beta(x_n)$
 $\beta(x) = 2^x$
 $x_1 = 1$ $y_2 = 4 + 0.01 \times 2^1$
 $y_1 = 4$ $= 4.02$
 $h = 0.01$

$x_2 = 1.01$ $y_3 = 4.02 + 0.01 \times 2^{1.01}$
 $y_2 = 4.02$ $= 4.04014$
 $h = 0.01$

③ Radians: $\theta = n\pi + a$
 Key angle: $\tan^{-1}(1) = \pi/4 = a$

So: $4(x - \pi/8) = n\pi + \pi/4$
 $x - \pi/8 = n\pi/4 + \pi/16$
 $x = n\pi/4 + 3\pi/16$

④ a) Key formula: $r = \frac{1}{2}n(n+1)$ $r^3 = \frac{1}{4}n^2(n+1)^2$

$\sum r^3 - br$
 $= \sum r^3 - b \sum r$
 $= \frac{1}{4}n^2(n+1)^2 - 3n(n+1)$
 $= \frac{1}{4}n(n+1) [n(n+1) - 12]$
 $= \frac{1}{4}n(n+1) [n^2 + n - 12]$
 $= \frac{1}{4}n(n+1)(n+4)(n-3)$

b) If $n = 1000$

$$\rightarrow S = 250(1001)(1004)(997)$$

$$2008 = 251 \times 2^3$$

Need to show that 251 and $2 \times 2 \times 2$ are factors of S

$$1004 = 251 \times 4 = 251 \times 2 \times 2$$

$$250 = 125 \times 2$$

\therefore Factors of 2008 are in S , so S is a multiple of 2008.

(5) a) Equation is: $\frac{x^2}{2^2} - \frac{y^2}{1^2} = 1$

Asymptotes at $y = b/a x$ & $y = -b/a x$

so $y = 1/2 x$ and $y = -1/2 x$

b) $xc = 4 \rightarrow 10/4 - y^2 = 1$

$$4 - y^2 = 1$$

$$\rightarrow y^2 = 3 \rightarrow y = \pm \sqrt{3}$$

c) i) Translation $\uparrow 2 \rightarrow 2 + \sqrt{3}$ and $2 - \sqrt{3}$

ii) Equation: $\frac{x^2}{4} - (y-2)^2 = 1$

Asymptotes: $y = 1/2 x + 2$ and $y = -1/2 x + 2$

(6) a) i) $\left(\begin{array}{c|c} \sqrt{3} & 3 \\ \hline 3 & -\sqrt{3} \end{array} \right)$

$$\left(\frac{\sqrt{3}}{3} \quad \frac{3}{-\sqrt{3}} \right) = \left(\frac{12}{0} \quad \frac{0}{12} \right) = 12 I$$

ii) $M = \begin{pmatrix} \sqrt{3} & 3 \\ 3 & -\sqrt{3} \end{pmatrix} \rightarrow$ Need: $\begin{pmatrix} 1/2 & \sqrt{3}/2 \\ \sqrt{3}/2 & -1/2 \end{pmatrix}$

so, take out factor of $2\sqrt{3} \rightarrow 2\sqrt{3} \begin{pmatrix} 1/2 & \sqrt{3}/2 \\ \sqrt{3}/2 & -1/2 \end{pmatrix}$

$$\rightarrow d_1 = 2\sqrt{3}$$

b) i) SF = $2\sqrt{3}$ or $\sqrt{12}$

ii) Reflections look like: $\begin{pmatrix} \cos(2\theta) & \sin(2\theta) \\ \sin(2\theta) & -\cos(2\theta) \end{pmatrix}$

$\therefore \theta = 30$

\Rightarrow Reflection in $y = \tan(30)$ or

$\rightarrow y = \sqrt{3}/3$ or

c) $M^4 = M^2 \times M^2$

$= 12I \times 12I = 144I$

= Enlargement, scale factor 144.

(7) a) i) $y = x^3 - x + 1$ $x = (-1 + h)$

$\rightarrow y = (-1 + h)^3 - (-1 + h) + 1$

$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$

$\rightarrow y = (-1)^3 + 3(-1)^2h + 3(-1)h^2 + h^3 + 1 - h + 1$

$y = -1 + 3h - 3h^2 + h^3 + 1 - h + 1$

$\rightarrow y = 1 + 2h - 3h^2 + h^3$

ii) 

$x_1 = -1$

$y_1 = (-1)^3 - (-1) + 1 = 1$

$x_2 = -1 + h$

$y_2 = 1 + 2h - 3h^2 + h^3$

(x_1, y_1) h

Gradient = $\frac{y_2 - y_1}{h}$

$= \frac{1 + 2h - 3h^2 + h^3 - 1}{h}$

$= \frac{2h - 3h^2 + h^3}{h} = 2 - 3h + h^2$

iii) As $h \rightarrow 0$, gradient $\rightarrow 2 - 3(0) + (0)^2 = 2$

gradient of chord \rightarrow gradient of tangent

b) i) $x^3 - x + 1 = 0$

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

$$f(x) = x^3 - x + 1$$


$$f'(x) = 3x^2 - 1$$

$$x_1 = -1$$

$$x_2 = -1 - \frac{(-1)^3 - (-1) + 1}{3(-1)^2 - 1}$$

$$= -1 - \frac{1}{2} = -\frac{3}{2}$$

ii)



8) a) i) $x^2 - 2x + 4 = 0$

$$\alpha + \beta = 2$$

$$\alpha\beta = 4$$

SUM $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$
 $= 2^3 - 3 \times 4 \times 2 = -16$

PRODUCT $\alpha^3\beta^3 = (\alpha\beta)^3$
 $= 4^3 = 64$

$$x^2 - \text{[SUM]}x + \text{[PRODUCT]} = 0$$

$$\rightarrow x^2 + 16x + 64 = 0$$

ii) Discriminant: $b^2 - 4ac$

$$= 16^2 - 4 \times 1 \times 64 = 0$$

\therefore Roots are real and equal

b) use formula: $\frac{2 \pm \sqrt{(-2)^2 - 4 \times 1 \times 4}}{2}$

$$= \frac{2 \pm \sqrt{-12}}{2} = \frac{2 \pm 2\sqrt{-3}}{2}$$

$$= 1 \pm \sqrt{-3} = 1 \pm i\sqrt{3}$$

c) From b) α and $\beta = 1 + i\sqrt{3}$ and $1 - i\sqrt{3}$

From a) α^3 and β^3 are equal

Therefore $(1 + i\sqrt{3})^3 = (1 - i\sqrt{3})^3$

9) a) $y = \frac{2}{x(x-4)}$

Asymptote: $x = 0$ and $x = 4$

As $x \rightarrow \infty$, $y \rightarrow \frac{2}{\infty} \rightarrow 0$

\therefore Asymptote at $y = 0$

b) Let $y = k$

$$k = \frac{2}{x(x-4)}$$

$$kx(x-4) = 2$$

$$kx^2 - 4kx = 2$$

$$kx^2 - 4kx - 2 = 0$$

At stationary points, roots are equal, so $b^2 - 4ac = 0$

$$(-4k)^2 - 4 \times k \times (-2) = 0$$

$$16k^2 + 8k = 0$$

$$2k^2 + k = 0$$

$$k(2k + 1) = 0$$

\downarrow

$$k = 0$$

\downarrow

$$k = -1/2$$

$$\boxed{k = 0}$$

$$0 = \frac{2}{x(x-4)} \rightarrow \text{NO SOLUTION}$$

$$\boxed{k = -1/2}$$

$$-1/2 = \frac{2}{x(x-4)} \rightarrow -1 = \frac{4}{x(x-4)}$$

$$\rightarrow -x(x-4) = 4$$

$$\rightarrow -x^2 + 4x = 4$$

$$\rightarrow x^2 - 4x + 4 = 0$$

$$\rightarrow (x-2)(x-2) = 0$$

$$\rightarrow x = 2 \quad \text{repeated solution} \quad \checkmark \quad \text{😊}$$

$$y = \frac{2}{2x-2} = -\frac{1}{2}$$

$$\therefore \text{co-ordinates} = (2, -\frac{1}{2})$$

c)

