

(1) $y_{n+1} = y_n + h f(x)$

$$\begin{aligned} x_1 &= 9 & \rightarrow & 6 + 0.25 \times \left[\frac{1}{2} + \sqrt{9} \right] \\ y_1 &= 6 & & = 6.05 \\ h &= 0.25 \end{aligned}$$

$$\begin{aligned} x_2 &= 9.25 & \rightarrow & 6.05 + 0.25 \times \left[\frac{1}{2} + \sqrt{9.25} \right] \\ y_2 &= 6.05 & & = 6.0995895... \\ & & & = 6.0996 \text{ (4dp)} \end{aligned}$$

(2) a) $\alpha + \beta = -\frac{8}{2} = -4$ $\alpha\beta = \frac{1}{2}$

b) i) $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$
 $= (-4)^2 - 2\left(\frac{1}{2}\right)$
 $= 16 - 1 = 15$

ii) $\alpha^4 + \beta^4 = (\alpha^2 + \beta^2)^2 - 2\alpha^2\beta^2$
 $= 15^2 - 2(\alpha\beta)^2$
 $= 225 - 2\left(\frac{1}{4}\right) = 44\frac{9}{2}$

c) sum $2\alpha^4 + \frac{1}{\beta^2} + 2\beta^4 + \frac{1}{\alpha^2}$
 $= 2\alpha^4 + 2\beta^4 + \frac{1}{\alpha^2} + \frac{1}{\beta^2}$
 $= 2(\alpha^4 + \beta^4) + \frac{\beta^2}{\alpha\beta^2} + \frac{\alpha^2}{\alpha\beta^2}$
 $= 2(\alpha^4 + \beta^4) + \frac{\alpha^2 + \beta^2}{(\alpha\beta)^2}$
 $= 2\left(44\frac{9}{2}\right) + \frac{15}{\left(\frac{1}{2}\right)^2}$
 $= 449 + 60 = 509.$

PRODUCT $(2\alpha^4 + 1/\beta^2)(2\beta^4 + 1/\alpha^2)$

$$\begin{aligned}
 &= 4\alpha^4\beta^4 + 2\alpha^2 + 2\beta^4 + 1/\alpha^2\beta^2 \\
 &= 4(\alpha\beta)^4 + 2(\alpha^2 + \beta^2) + 1/(\alpha\beta)^2 \\
 &= 4(1/2)^4 + 2(15) + 1/(1/2)^2 \\
 &= 1/4 + 30 + 4 = 137/4
 \end{aligned}$$

$\rightarrow x^2 - (\text{Sum})x + (\text{PRODUCT}) = 0$

$\rightarrow x^2 - 509x + 137/4 = 0$

$\rightarrow 4x^2 - 2036x + 137 = 0$

③ $\sum r^2(r-b) = \sum r^3 - br^2 = \sum r^3 - b\sum r^2$

Need: $\sum_1^{60} r^3 - b\sum_1^{60} r^2 = \left(\sum_1^{60} r^3 - b\sum_1^{60} r^2 \right)$

$$\begin{aligned}
 &= 1/4(60)^2(61)^2 - b \cdot 1/6(60)(61)(2 \times 60 + 1) - ((1+8) - (6+24)) \\
 &= 3,348,900 - 442,860 - (-21) \\
 &= 2,906,061
 \end{aligned}$$

④ Let $z = a + ib$

$\rightarrow 5i(a + ib) + 3(a - ib) + 1b = 8i$

$5ai + 5i^2ab + 3a - 3bi + 1b = 8i$

$5ai - 5ab + 3a - 3bi + 1b = 8i$

REAL $-5b + 3a + 1b = 0 \quad \text{①}$

IMAG $5a - 3b = 8 \quad \text{②}$


$$\begin{aligned} \textcircled{1} \quad 3a - 5b &= -16 & \times 5 \rightarrow & 15a - 25b = \\ \textcircled{2} \quad 5a - 3b &= 8 & \times 3 \rightarrow & 15a - 9b = 24 \end{aligned}$$

$$\textcircled{2} - \textcircled{1} \rightarrow 16b = 104 \rightarrow b = 6.5$$

$$\begin{aligned} \textcircled{1} \quad 3a - 5(6.5) &= -16 \\ 3a - 32.5 &= -16 \\ 3a &= 16.5 \\ a &= 5.5 \end{aligned}$$

$$\rightarrow 2 = 5.5 + 6.5i$$

5) a)



$$\begin{aligned} \textcircled{*} \quad y &= (-5+h)(-5+h+3) \\ &= (-5+h)(-2+h) \\ &= 10 - 5h - 2h + h^2 \\ &= h^2 - 7h + 10 \end{aligned}$$

$$\text{Gradient} = \frac{h^2 - 7h + 10 - 10}{-5 + h - (-5)}$$

$$= \frac{h^2 - 7h + \cancel{10}}{h} = h - 7$$

b) As $h \rightarrow 0$, gradient $\rightarrow 0 - 7 = -7$

This is a good approximation of gradient of curve at $(-5, 10)$


(6) a) $y = \frac{1}{x(x+2)}$

$[x=0], [x=-2]$

As $x \rightarrow \infty, y \rightarrow \frac{1}{\infty} \rightarrow [y=0]$

b) i) $x = -1 \rightarrow y = \frac{1}{-1(-1+2)} = \frac{1}{-1} = -1$

ii)



c) Mark $y = 1/8$ on sketch

Solve:

$$\frac{1}{x(x+2)} = \frac{1}{8}$$

$$8 = x(x+2)$$

$$8 = x^2 + 2x$$

$$x^2 + 2x - 8 = 0$$

$$(x+4)(x-2) = 0$$



$$x = -4$$




$$x = 2$$

From sketch, we need curve below $y = 1/8$

$$\rightarrow x \leq -4, \quad x \geq 2$$


$$\text{or } -2 < x < 0$$

7 a) i)



$$\begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

ii)



$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

b) Change order: \rightarrow

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

c) i)

$$\begin{bmatrix} -3 & -\sqrt{3} \\ -\sqrt{3} & 3 \end{bmatrix} \begin{bmatrix} 12 & 0 \\ 0 & 12 \end{bmatrix} = 12 I$$

ii) Scale Factor must be $\sqrt{12} = 2\sqrt{3}$

Take $2\sqrt{3}$ out of matrix:

$$2\sqrt{3} \begin{bmatrix} \sqrt{3}/2 & -1/2 \\ -1/2 & \sqrt{3}/2 \end{bmatrix}$$

Reflection \rightarrow $\begin{bmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{bmatrix}$

$$\sin^{-1}(-1/2) = -30 \rightarrow \theta = -15^\circ \text{ or } -75^\circ$$

or
-150

Check in cos: $\cos(2\theta)$ must = $-\sqrt{3}/2$
 $\cos(-30) = \sqrt{3}/2$ ✗
 $\cos(-150) = -\sqrt{3}/2$ ✓

∴ $\theta = -75$
 → Line of reflection: $y = [\tan(-75)]x$
 or $y = [-\tan(75)]x$
 or $y = [\tan(105)]x$

(8) a) $\theta = 2n\pi \pm a$

Key angle: $\cos^{-1}(\sqrt{3}/2) = \pi/4$

→ $5/4 x - \pi/3 = 2n\pi \pm \pi/4$
 $5/4 x = 2n\pi + \pi/3 \pm \pi/4$
 $5x = 8n\pi + 4\pi/3 \pm \pi$
 $x = 8/5 n\pi + 4\pi/15 \pm \pi/5$

~~$x = 8/5 n\pi + 20\pi/3 + 5\pi$, $x = 8/5 n\pi + 20\pi/3 - 5\pi$~~

$x = \frac{24n\pi}{15} + \frac{4\pi}{15} + \frac{3\pi}{15}$, $x = \frac{24n\pi}{15} + \frac{4\pi}{15} - \frac{3\pi}{15}$

$x = \frac{24n\pi + 7\pi}{15}$ $x = \frac{24n\pi + \pi}{15}$

- b) Smallest value of $n = 0 \rightarrow \frac{7\pi}{15} \approx \frac{7\pi}{15}$
 Largest value of $n = 12 \rightarrow 10\frac{2}{3}\pi \approx 19.266... \pi$

$$\text{Need } \sum_0^{12} \frac{24n\pi + 7\pi}{15} + \sum_0^{12} \frac{24n\pi + \pi}{15}$$

$$= \sum_0^{12} \frac{24n\pi}{15} + \sum_0^{12} \frac{7\pi}{15} + \sum_0^{12} \frac{24n\pi}{15} + \sum_0^{12} \frac{\pi}{15}$$

$$= \text{use } \sum_0^n r = \frac{1}{2} n(n+1) \quad \& \quad \sum_0^n 1 = n+1 \quad \leftarrow \begin{array}{l} \text{As if} \\ \text{Start at} \\ 0. \end{array}$$


$$\rightarrow \frac{24\pi}{15} \sum_0^{12} n + \frac{7\pi}{15} \sum_0^{12} 1 + \frac{24\pi}{15} \sum_0^{12} n + \frac{\pi}{15} \sum_0^{12} 1$$

$$= \left[\frac{24\pi}{15} \times \frac{1}{2} \times 12 \times 13 \right] + \frac{7\pi}{15} (13) + \left[\frac{24\pi}{15} \times \frac{1}{2} \times 12 \times 13 \right] + \frac{\pi}{15} \times 13$$

$$= \frac{\pi}{15} [1872 + 91 + 1872 + 13]$$

$$= 3848/15 \pi \quad \rightarrow \quad k = 3848/15$$

(9) c)



b) $y = cx + k$

$$\rightarrow \frac{x^2}{16} + \frac{(cx+k)^2}{9} = 1$$

$$\rightarrow 9x^2 + 16(cx+k)^2 = 144$$

$$9x^2 + 16[x^2 + 2cxk + k^2] = 144$$

$$9x^2 + 16x^2 + 32cxk + 16k^2 - 144 = 0$$

$$25x^2 + 32kx + (16k^2 - 144) = 0$$

For 2 distinct solutions, $b^2 - 4ac > 0$

$$\rightarrow (32k)^2 - 4 \times 25 \times (16k^2 - 144) > 0$$


$$1024k^2 - 1600k^2 + 14400 > 0$$

$$~~1024k^2 - 1600k^2~~$$

$$-576k^2 + 14400 > 0$$

$$14400 > 576k^2$$

$$k^2 < 25$$



c) Translation $\rightarrow \frac{(x-a)^2}{16} + \frac{(y-b)^2}{9} = 1$

$$\rightarrow 9(x-a)^2 + 16(y-b)^2 = 144$$

$$\rightarrow 9(x^2 - 2ax + a^2) + 16(y^2 - 2by + b^2) = 144$$

$$\rightarrow 9x^2 - 18ax + 9a^2 + 16y^2 - 32by + 16b^2 = 144$$

NEED:

$$9x^2 + 16y^2 \quad \boxed{-18ax + 18x} \quad \boxed{-32by - 64y} = \boxed{144 - 9a^2 - 16b^2}$$

① ② ③

① $a = -1$

② $b = 2$

③ $c = 144 - 9(-1)^2 - 16(2)^2 = 71$

d) From part b, we know that $b^2 - 4ac = 0$
when $k = 5$ or $k = -5$

→ $y = x + 5$ or $y = x - 5$ are tangents
to original curve E

E has been translated $\begin{pmatrix} -1 \\ 2 \end{pmatrix}$

∴ New tangents are:

$$(y - 2) = (x + 1) + 5$$

$$y - 2 = x + 6$$

$$y = x + 8$$

$$\left\{ \begin{array}{l} (y - 2) = (x + 1) - 5 \\ y - 2 = x - 4 \end{array} \right.$$

$$y - 2 = x - 4$$

$$y = x - 2$$