

Stats 1 - January 2005

- ① a) 1, weak positive correlation: takings seem to increase as temperature increases
2, one outlier / unusual result

b) Monday 10: (4, 312)

c) From calculator: $\Sigma x = 71$

$$r = 0.9177070\dots$$

d) The weather: i.e. if it is raining or not.

- ② a) From calculator:

$$\bar{x} = 152.5 \quad \sigma = 3.5 \quad n = 12$$

$$98\% \text{ z multiplier (2 tailed)} = 2.3263$$

$$\therefore \mu = \bar{x} \pm z_m \times \frac{\sigma}{\sqrt{n}}$$

$$\mu = 152.5 \pm 2.3263 \times \frac{3.5}{\sqrt{12}}$$

$$= 152.5 \pm 2.3504\dots$$

$$= (150.15, 154.85)$$

b) Confidence Interval: 150 ml is below the range of the CI \rightarrow mean volume is above 150 ml

Sample: $\frac{4}{12}$ or 25% have volumes below 150 ml

\therefore claim does not appear to be valid.

c) we know the volume is normally distributed.

- ③ a) See Mark Scheme

b) From calculator: $a = 14.336283\dots$ (intercept)

$b = 7.500894\dots$ (gradient)

$$\rightarrow y = 14.34 + 7.50x$$

x	0	25
y	14.3	201.9

See mark scheme bar graph

c) i) $x = 15 \rightarrow y = 14.34 + 7.5(15)$
 $= 126.9$ (10p)

reliable as data in observed range.

ii) $x = 35 \rightarrow y = 14.34 + 7.5(35)$
 $= 276.9$ (10p)


unreliable as 35 is outside observed range.

d) a = intercept = time to travel from area to/from depot


b = gradient = time to deliver each/one parcel

4) a) $X \sim N(106, 4^2)$

i) $P(X < 110)$
 $= P(Z < \frac{110 - 106}{4})$
 $= P(Z < 1)$
 $= 0.84134$




ii) $= P(X < 100)$
 $= P(Z < \frac{100 - 106}{4})$
 $= P(Z < -1.5)$
 $= P(Z > 1.5)$
 $= 1 - P(Z < 1.5)$
 $= 1 - 0.93319$
 $= 0.06681$



b) $X \sim N(\mu, 4^2)$

$P(X < 100) = 0.02$

Look up 0.98 \rightarrow Z value of 2.0537
 $\rightarrow -2.0537$



Standardize:

$$P\left(Z < \frac{100 - \mu}{4}\right) = 0.02$$

$$\rightarrow \frac{100 - \mu}{4} = -2.0537$$

$$100 - \mu = -8.2148$$

$$\mu = 100 + 8.2148$$

$$= 108.2148$$

c) $\bar{X} \sim N(108.5, 4^2/10)$

i) Mean = 108.5

Variance = $4^2/10 = 1.6$


ii) $P(\bar{X} > 110)$

$$= P\left(Z > \frac{110 - 108.5}{\frac{4}{\sqrt{10}}}\right)$$

$$= P(Z > 1.19)$$

$$= 1 - P(Z < 1.19)$$

$$= 1 - 0.88298 = 0.11702$$



5) a) $W \sim B(7, 0.4)$

i) $P(W \leq 2) = 0.4199$ (from tables)

ii) $P(1 < W < 5)$

can be: 2, 3, 4

$$= P(W \leq 4) - P(W \leq 1)$$

$$= 0.9037 - 0.1586 = 0.7451$$

b) $W \sim B(28, 0.4)$

$$P(W = 7) = {}^{28}C_7 \times 0.4^7 \times 0.6^{21}$$

$$= 0.042556 \dots$$

c) Mean = $np = 7 \times 0.4 = 2.8$
 Variance = $np(1-p) = 7 \times 0.4 \times 0.6 = 1.68$
 $\rightarrow SD = \sqrt{1.68} = 1.2961\dots$

d) i) From calculator:

$\bar{x} = 2.8$

$s = 2.267786\dots$

$\sum x = 140$
 $\sum x^2 = 644$

ii) Means are the same \rightarrow support believe that probability = 0.4

SD are very different \rightarrow does not support believe that probabilities are independent.

(b) a) A Totals: 64, 644, 946, 710, 944 : Grand = 1654

i) $P(F) = 944/1654$

ii) $P(F \cap A) = 275/1654$

iii) $P(F \cup A) = \frac{944 + 369}{1654} = \frac{1313}{1654}$

iv) $P(F/A) = \frac{P(F \cap A)}{P(A)} = \frac{275}{1654} = \frac{644}{1654}$
 $= \frac{275}{644}$

b) i) $P(3M) = \frac{710}{1654} \times \frac{709}{1653} \times \frac{708}{1652} = 0.07890\dots$

ii) $P(FFM) = \frac{944}{1654} \times \frac{943}{1653} \times \frac{710}{1652}$
 $= 0.13993\dots$

There are 3 arrangements of FFM

$\rightarrow 3 \times 0.13993 = 0.41980\dots$

c) i) $(F \cap A)$ Female and Academic

$(F \cup A)$ Male, or Academic or both.