

Stats 1 - January 2011

- ① a) i) strong positive correlation $\rightarrow r = 0.8$
ii) weak negative correlation $\rightarrow r = -0.3$

b) i) From calculator: $\sum x^2 = 6142.97$
 $r = 0.7570847\dots$

ii) Strong, positive, linear correlation between the circumference and weight of cricket balls.

② a) i) $175/645$

ii) $519/645$

iii) $63/645$

iv) $P(L|F) = \frac{P(L \cap F)}{P(F)} = \frac{94}{126}$

v) $P(M|L') = \frac{P(M \cap L')}{P(L')} = \frac{175 + 54 + 35}{296} = \frac{264}{296}$

b) $P(F, F) = \frac{94}{126} \times \frac{93}{125} = \frac{8742}{15750}$ or 0.555

c) $P(L, C, \text{NEITHER}) = \frac{349}{645} \times \frac{193}{644} \times \frac{103}{643} = 0.02597$

6 ways of arranging 3 objects

$$\rightarrow 6 \times 0.02597 = 0.15585\dots$$

③ a) i) True boundaries = $0.975 - 1.005$

$$\text{Midpoint} = \frac{0.975 + 1.005}{2} = 0.99$$

ii) Minimum = 0.975 , Maximum = 1.005

b) From calculator, and using midpoints for x :

$$\sum x^2 = 112.9662 \quad \sum x = 106.2$$

$$\rightarrow \text{mean}(\bar{x}) = 1.062$$

$$\text{Sample standard deviation (s)} = 0.04285$$

c) i) $\bar{x} = 1.062$ $s = 0.043$ $n = 100$

99% multiplier (2 tailed) $z = 2.5758$

$$\begin{aligned}
 \text{99\% CI for } \mu &= \bar{x} \pm z \times \frac{s}{\sqrt{n}} \\
 &= 1.062 \pm 2.5758 \times \frac{0.043}{\sqrt{100}} \\
 &= 1.062 \pm 0.1107 \dots \\
 &= (1.051, 1.073)
 \end{aligned}$$

- ii) we know the population is normally distributed
- iii) we have used midpoints as the data is in groups

- d) i) The lower bound of the confidence interval (1.051) is greater than 1
- ii) 99/100 of the pieces of data in the sample are in this range.

④ $X \sim B(15, 0.45)$ $X = \text{number of times target hit}$

a) i) $P(X \leq 5) = 0.2608$ (tables)

ii) $P(X > 10) = 1 - P(X \leq 10)$
 $= 1 - 0.9745 = 0.0255$

iii) $P(X = 6) = {}^{15}C_6 \times 0.45^6 \times 0.55^9$
 $= 0.1914$

iv) $P(5 \leq X \leq 10)$
 CAN BE: 5, 6 ... 10
 $\rightarrow P(X \leq 10) - P(X \leq 4)$
 $= 0.9745 - 0.1204 = 0.8541$

b) i) HITS = $P(H) = 0.85$
 $P(H^c, H) = 0.15 \times 0.8 = 0.12$ } +
0.97

ii) Let $M =$ number of times the target is missed
 $M \sim B(50, 0.03)$ $P(M) = 1 - 0.97$

$P(\text{HITS} \geq 48)$

HITS : 48 49 50

MISSSES : 2 1 0

$= P(M \leq 2) = 0.9108$ (from tables)

iii) Probability she shoots with 2nd = Probability she misses with 1st

$= 1 - 0.85 = 0.15$

$\therefore \text{Mean} = np = 80 \times 0.15 = 12 \text{ times}$

5) a) Time taken is dependent on leaving time

b) From calculator: $a = 29.9545...$ (intercept)

$b = 1.281818...$ (gradient)

$\rightarrow y = 29.95 + 1.28x$

c) 7.45am $\rightarrow x = 15$

$\rightarrow y = 29.95 + 1.28(15)$

$= 49.15 \text{ minutes}$

\rightarrow arrival time of 8.34am

$= 26 \text{ minutes before}$

d) i) $x = 85 \rightarrow y = 29.95 + 1.28(85)$

$= 138.75$


ii) You shouldn't use x values outside of the range of data to make predictions.

6) a) $V \sim N(412, 8^2)$

i) $P(V < 400)$


$= P(Z < \frac{400 - 412}{8})$

$= P(Z < -1.5)$



$$\begin{aligned}
 &= P(Z > 1.5) \\
 &= 1 - P(Z < 1.5) \\
 &= 1 - 0.93319 = 0.06681
 \end{aligned}$$

$$\begin{aligned}
 \text{ii) } P(V > 420) \\
 &= P(Z > \frac{420 - 412}{8}) \\
 &= P(Z > 1) \\
 &= 1 - P(Z < 1) \\
 &= 1 - 0.84134 \\
 &= 0.15866
 \end{aligned}$$




$$\text{iii) } P(V = 410) = 0$$

$$\text{b) i) } P(V < 400) = 0.05$$

$$\text{Z value for } 0.95 = 1.6449$$

$$\therefore \text{Z value for } V = 400$$

$$= -1.6449$$



Standardize:

$$\frac{400 - \mu}{\sigma} = -1.6449$$

$$\rightarrow 400 - \mu = -1.6449\sigma$$

$$P(V > 420) = 0.01$$

$$\text{Z value for } 0.99 = 2.3263$$

→ Standardize:

$$\frac{420 - \mu}{\sigma} = 2.3263$$

$$\rightarrow 420 - \mu = 2.3263\sigma$$



$$ii) \quad ① \quad 400 - \mu = -1.6449\sigma$$

$$② \quad 420 - \mu = 2.3263\sigma$$

$$② - ① \rightarrow 20 = 3.9712\sigma$$

$$\rightarrow \sigma = \frac{20}{3.9712} = 5.036\dots$$

$$\boxed{iv} \quad ① \quad 400 - \mu = -1.6449 (5.036\dots)$$

$$\rightarrow 400 - \mu = -8.2841\dots$$

$$\rightarrow 400 + 8.2841\dots = \mu$$

$$\rightarrow \mu = 408.28\dots$$