

Stats 1 - May / June 2006

① a) i) From calculator: $r = 0.14308\dots$

ii) Very weak positive linear correlation.

Suggests there is no relationship between the price and the number of pages.

iii) The author / popularity

b) Very strong positive linear correlation.

The sale price appears to be determined by the number of pages.


② $H \sim N(185, 10^2)$

a) i) $P(H < 200)$

$$= P\left(Z < \frac{200 - 185}{10}\right)$$

$$= P(Z < 1.5)$$

$$= 0.93319$$




ii) $P(H > 175)$

$$= P\left(Z > \frac{175 - 185}{10}\right)$$

$$= P(Z > -1)$$

$$= P(Z < 1)$$

$$= 0.84134$$



iii) $P(175 < H < 200)$


$$= P(H < 200) - P(H < 175)$$

using previous answers

$$= 0.93319 - [1 - 0.84134] = 0.77453$$

$$b) \bar{H} \sim N\left(185, \frac{10^2}{4}\right)$$

$$\begin{aligned} & P(\bar{H} > 190) \\ &= P\left(Z > \frac{190 - 185}{\frac{10}{\sqrt{4}}}\right) \\ &= P(Z > 1) \\ &= 1 - P(Z < 1) \\ &= 1 - 0.84134 \\ &= 0.15866 \end{aligned}$$



③ a) i) From Calculator: $a = 262.888\dots$ (intercept)
 $b = -3.25$ (gradient)
 $\rightarrow y = 262.8 - 3.25x$

ii) $b = -3.25 \rightarrow$ Each extra month (x) has a decrease in pressure of 3.25 kPa

iii) $a = 262.8 \rightarrow$ The initial pressure of the tyre is 262.8 kPa

b) i) $2x - 3.25 = -6.5$ ($2 \times$ gradient)

ii) $x = 8 \rightarrow y = 265 + -6.5(8)$
 $= 213$, which < 220

④ a) i) From Calculator: $\bar{x} = 505.2$ $n = 10$
 $s = 5.95411\dots$

99% CI (2 tailed) $\rightarrow z = 2.5758$

99% CI $\rightarrow N = \bar{x} \pm z \times \frac{s}{\sqrt{n}}$

$$N = 505.2 \pm 2.5758 \times \frac{5.95411}{\sqrt{10}}$$

$\rightarrow \mu = 505.2 \pm 4.854$
 $\rightarrow \mu = (500.3, 510.054)$ (1dp)

- ii) The weights of packets were normally distributed.
- iii) **EXAMPLE** $3/10$ or 30% of packets below 500g
CI 500g outside (below) 99% confidence interval
 \therefore weight seems to be more than 500g
 Claim does not seem justified

b) 1% (100% - 99%)

5) a) $K \sim P(15, 0.3)$

i) $P(K=5) \rightarrow {}^{15}C_5 \times 0.3^5 \times 0.7^{10}$
 $= 0.20613\dots$

ii) $P(K \leq 7) = 0.95$ (from table)

iii) $P(2 < K < 7)$

CAN BE: 3, 4, 5, 6

$\rightarrow P(K \leq 6) - P(K \leq 2)$
 $= \cancel{0.9884} - \cancel{0.1268} =$
 $0.8684 - 0.1268 = 0.7421$

b) i) MEAN = $np = 15 \times 0.4 = 6$

$SD = \sqrt{np(1-p)} = \sqrt{15 \times 0.4 \times 0.6} = \sqrt{3.6}$
 $= 1.897\dots$

ii) From calculator: $\bar{x} = 6$
 $s = 2.9814$

iii) Means are the same
 Standard deviations are different
 We have reasons to doubt Kirk's claims

6) a) i) $P(D) = \frac{120}{320}$

ii) $P(D \cap R) = \frac{24}{320}$

iii) $P(D \cup T) = \frac{120 + 88}{320} = \frac{208}{320}$

iv) $P(D | R) = \frac{24}{64}$

64 = house with 0 kids

v) $P(R | D') = \frac{40}{200}$

200 = houses not detached

b) i) $R \cap S$, $R \cap T$ or $S \cap T$

ii) If independent $P(D | R) = P(D)$

$$P(D | R) = \frac{24}{64} = 0.375$$

$$P(D) = \frac{120}{320} = 0.375$$

$\therefore D$ & R are Independent

c) i) $D' \cup T$ = A semi-detached house, or a house with 2 children, or both

ii) $D \cap (R \cup S)$ = A detached house that has less than 2 children.