www.mymathscloud.com
My Maths Cloud

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	

CENTRE NUMBER						CANDIDATE NUMBER					

* 9 5 4 9 3 2 4 7 3 1 *

FURTHER MATHEMATICS **9231/12**

Paper 1 Further Pure Mathematics 1 **May/June 2021**

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Any blank pages are indicated.

DC (PQ) 199003/2
© UCLES 2021

BLANK PAGE

www.mymathscloud.com
My Maths Cloud

www.mymathscloud.com
My Maths Cloud

1 Prove by mathematical induction that $2^{4n} + 31^n - 2$ is divisible by 15 for all positive integers n. [6]

www.mymathscloud.com
My Maths Cloud

2 **(a)** Use standard results from the List of formulae (MF19) to find $\sum_{r=1}^{n}\left(1-r-r^2\right)$ in terms of n, simplifying your answer. [3]

www.mymathscloud.com
My Maths Cloud

(b) Show that

$$\frac{1-r-r^2}{(r^2+2r+2)(r^2+1)} = \frac{r+1}{(r+1)^2+1} - \frac{r}{r^2+1}$$

and hence use the method of differences to find $\sum_{r=1}^{n} \frac{1-r-r^2}{(r^2+2r+2)(r^2+1)}$. [5]

(c) Deduce the value of $\sum_{r=1}^{\infty} \frac{1-r-r^2}{(r^2+2r+2)(r^2+1)}$. [1]

www.mymathscloud.com
My Maths Cloud

3 The equation $x^4 - 2x^3 - 1 = 0$ has roots α, β, γ, δ.

(a) Find a quartic equation whose roots are α^3, β^3, γ^3, δ^3 and state the value of $\alpha^3 + \beta^3 + \gamma^3 + \delta^3$. [4]

www.mymathscloud.com My Maths Cloud

(b) Find the value of $\frac{1}{\alpha^3}+\frac{1}{\beta^3}+\frac{1}{\gamma^3}+\frac{1}{\delta^3}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find the value of $\alpha^4+\beta^4+\gamma^4+\delta^4$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

www.mymathscloud.com
My Maths Cloud

4 The matrix **M** represents the sequence of two transformations in the x-y plane given by a rotation of 60° anticlockwise about the origin followed by a one-way stretch in the x-direction, scale factor d ($d \neq 0$).

(a) Find **M** in terms of d. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The unit square in the x-y plane is transformed by **M** onto a parallelogram of area $\frac{1}{2}d^2$ units^2.

Show that $d = 2$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

www.mymathscloud.com
My Maths Cloud

The matrix **N** is such that $\mathbf{MN} = \begin{pmatrix} 1 & 1 \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix}$.

(c) Find **N**. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Find the equations of the invariant lines, through the origin, of the transformation in the x-y plane represented by **MN**. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

www.mymathscloud.com
My Maths Cloud

5 The curve C has polar equation $r = a\cot\left(\frac{1}{3}\pi - \theta\right)$, where a is a positive constant and $0 \leqslant \theta \leqslant \frac{1}{6}\pi$.

It is given that the greatest distance of a point on C from the pole is $2\sqrt{3}$.

(a) Sketch C and show that $a = 2$. [3]

.....

.....

(b) Find the exact value of the area of the region bounded by C, the initial line and the half-line $\theta = \frac{1}{6}\pi$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

www.mymathscloud.com My Maths Cloud

(c) Show that C has Cartesian equation $2\left(x+y\sqrt{3}\right)=\left(x\sqrt{3}-y\right)\sqrt{x^2+y^2}$. [3]

www.mymathscloud.com
My Maths Cloud

6 Let t be a positive constant.

The line l_1 passes through the point with position vector $t\mathbf{i}+\mathbf{j}$ and is parallel to the vector $-2\mathbf{i}-\mathbf{j}$. The line l_2 passes through the point with position vector $\mathbf{j}+t\mathbf{k}$ and is parallel to the vector $-2\mathbf{j}+\mathbf{k}$.

It is given that the shortest distance between the lines l_1 and l_2 is $\sqrt{21}$.

(a) Find the value of t. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The plane Π_1 contains l_1 and is parallel to l_2.

(b) Write down an equation of Π_1, giving your answer in the form $\mathbf{r}=\mathbf{a}+\lambda\mathbf{b}+\mu\mathbf{c}$. [1]

.....

.....

.....

.....

.....

www.mymathscloud.com
My Maths Cloud

The plane Π_2 has Cartesian equation $5x - 6y + 7z = 0$.

(c) Find the acute angle between l_2 and Π_2. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Find the acute angle between Π_1 and Π_2. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

www.mymathscloud.com
My Maths Cloud

7 The curve C has equation $y = \dfrac{x^2 + x + 9}{x + 1}$.

(a) Find the equations of the asymptotes of C. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the coordinates of the stationary points on C. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

www.mymathscloud.com
My Maths Cloud

(c) Sketch C, stating the coordinates of any intersections with the axes. [3]

.....

(d) Sketch the curve with equation $y = \left|\dfrac{x^2+x+9}{x+1}\right|$ and find the set of values of x for which $2\left|x^2+x+9\right| > 13\left|x+1\right|$. [5]

www.mymathscloud.com
My Maths Cloud

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2021