

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
GCE Advanced Subsidiary Level and GCE Advanced Level

**MARK SCHEME for the October/November 2010 question paper
for the guidance of teachers**

9709 MATHEMATICS	
9709/11	Paper 1, maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	Page
	GCE A/AS LEVEL – October/November 2010	9709	11

Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol \surd implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2010	9709	11

The following abbreviations may be used in a mark scheme or used on the scripts:

- AEF Any Equivalent Form (of answer is equally acceptable)
- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
- CWO Correct Working Only – often written by a ‘fortuitous’ answer
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)
- SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through $\sqrt{}$ ” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper 11
	GCE AS/A LEVEL – October/November 2010	9709	11
1	$\int \left(x + \frac{1}{x}\right)^2 dx$ $= \frac{x^3}{3} - \frac{1}{x} + 2x + (c)$	B1 × 3	co. Omission of middle term of expansion can still get 2/3. [3]
2	$(1 + ax)^6$ Term in $x = 6ax$ Equate with $-30 \rightarrow a = -5$ Term in $x^3 = \frac{6 \cdot 5 \cdot 4}{3!} a^3$ \rightarrow coefficient of -2500	B1 B1√ B1 B1√	co √ from his answer for $6ax$ co For $20 \times a^3$ [4]
3	$f: x \mapsto 2x + 3,$ $g: x \mapsto x^2 - 2x,$ $gf(x) = (2x + 3)^2 - 2(2x + 3)$ $= 4x^2 + 8x + 3$ $= 4(x + 1)^2 - 1$	M1 A1 $3 \times B1\sqrt$	Must be f into g, not g into f. co Allow all these as √ for either fg or gf. [5]
4	(i) $\frac{\sin x \tan x}{1 - \cos x} = \frac{\sin^2 x}{\cos x(1 - \cos x)}$ $= \frac{1 - \cos^2 x}{\cos x(1 - \cos x)}$ $= \frac{(1 - \cos x)(1 + \cos x)}{\cos x(1 - \cos x)} = \frac{1}{\cos x} + 1$ (ii) $\frac{1}{\cos x} + 1 + 2 = 0$ $\rightarrow \cos x = -\frac{1}{3}$ $\rightarrow x = 109.5^\circ$ or 250.5°	M1 M1 M1 M1 A1 A1√	Use of $\tan x = \sin x \div \cos x$ Use of $\sin^2 x = 1 - \cos^2 x$ Realising the need to use difference of 2 squares. Answer given. [3] Uses part (i) with $\cos x$ as subject. co. √ for $360^\circ - 1^{\text{st}}$ answer. [3]
5	$\vec{AC} = -6\mathbf{i} + 10\mathbf{k}$ $\vec{BC} = -8\mathbf{j} + 10\mathbf{k}$ $\vec{AC} \cdot \vec{BC} = 100$ $\vec{AC} \cdot \vec{BC} = \sqrt{136}\sqrt{164} \cos ACB$ Angle $ACB = 48.0^\circ$	B1 B1 M1 M1 M1 A1	co (or \vec{CA}) co (or \vec{CB}) Must be scalar – available for any pair For modulus – available for any vector All linked correctly – for ACB only co [6]

6	(a) $a + 4d = 18$ $\frac{5}{2}(2a + 4d) = 75$ Solution $\rightarrow a = 12, d = 1\frac{1}{2}$	B1 B1 M1 A1	[4]	co or $75 = 5/2(a + 18) \rightarrow a = 12$ etc co Solution of sim equations co for both
	(b) $a = 16$ and $ar^3 = \frac{27}{4}$ $r = \frac{3}{4}$ Sum to infinity = 64	B1 M1 A1		[3]
7	$x \mapsto 3 - 2 \tan(\frac{1}{2}x)$ (i) Range of $f \leq 3$	B1	[1]	co. Allow <
	(ii) $f(\frac{2}{3}\pi) = 3 - 2\sqrt{3}$	B1	[1]	co
	(iii)	B2, 1, 0 Indep.	[2]	Starting at $y = 3$ Shape correct – no turning points. Tending tangentially towards $x = \pi$
	(iv) $y = 3 - 2 \tan(\frac{x}{2})$ $\rightarrow f^{-1}(x) = 2 \tan^{-1}(\frac{3-x}{2})$	M1 M1 A1	[3]	Attempt at making x the subject. Order of operations all ok. co – but with x , not y .
8	(i) $2x + 2y + \frac{\pi x}{2} = 60$ $\rightarrow y = 30 - x - \frac{\pi x}{4}$	M1 A1	[2]	Linking 60 with sum of at least 4 sides and use of radians co
	(ii) $A = xy + \frac{\pi x^2}{4}$ $= x(30 - x - \frac{\pi x}{4}) + \frac{\pi x^2}{4}$ $= 30x - x^2$	M1 A1	[2]	Subs “y” into area eqn and use $\frac{1}{2}r^2\theta$ co.
	(iii) $\frac{dA}{dx} = 30 - 2x$ $= 0$ when $x = 15$ cm	M1 A1	[2]	Knowing to differentiate Sets differential to 0 + solution. co.
	(iv) Max.	M1 A1	[2]	Any valid method. co.

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2010	9709	11

9	(i) $RS^2 = 10^2 - 6^2$ $\rightarrow RS = 8 \text{ cm.}$	M1 A1	[2]	Use of Pythagoras (or other) Answer given.
	(ii) $\sin \theta = 8/10$ oe \rightarrow angle $RPQ = 0.9273$ radians	M1 A1	[2]	Use of trig – even if with degrees. co in radians. (Accept 0.927)
	(iii) Region = trapezium – 2 sectors Area of trapezium = 40 cm^2 Large sector = $\frac{1}{2} \times 8^2 \times 0.9273$ Small sector angle = $(\pi - 0.9273)$ Small sector = $\frac{1}{2} \times 2^2 \times 2.214$ $\rightarrow 5.90 \text{ cm}^2$	B1 M1 M1 A1	 [4]	co Use of $\frac{1}{2}r^2\theta$. Use of $\frac{1}{2}r^2\theta$ with angle = $\pi -$ (ii) co
10	$y = 4x - x^2 + 3$			
	(i) $\frac{dy}{dx} = 4 - 2x$ At $x = 3$, $m = -2$ Gradient of normal = $\frac{1}{2}$ Eqn of normal $y - 6 = \frac{1}{2}(x - 3)$ $\rightarrow 2y = x + 9$	B1 M1 M1 A1	 [4]	co Use of $m_1m_2 = -1$ Use of $y - k = m(x - h)$ or $y = mx + c$ (where m is gradient of normal)
	(ii) Meets axes at $(0, \frac{9}{2})$ and $(-9, 0)$ Mid-point is $(\frac{-9}{2}, \frac{9}{4})$	M1 A1	 [2]	Sets x and y to 0 + midpoint formula. co.
(iii) $2y = x + 9$, $y = 4x - x^2 + 3$ $\rightarrow 2x^2 - 7x + 3 = 0$ oe $\rightarrow (\frac{1}{2}, \frac{3}{4})$	M1 A1 M1 A1	 [4]	Eliminates x completely. Correct eqn. Solution of quadratic. co	

11	$y = \frac{9}{2-x}$			
	<p>(i) $\frac{dy}{dx} = -9(2-x)^{-2} \times -1$</p> <p>$\frac{9}{(2-x)^2} \neq 0$. No turning points.</p>	<p>B1 B1 B1√</p>	<p>[3]</p>	<p>Without the “× -1” Indep. With the “×-1”. Indep. √ provided of form $k \div (2-x)^2$.</p>
	<p>(ii) $V = \pi \int \frac{81}{(2-x)^2} dx$</p> <p>$\int y^2 dx = -81(2-x)^{-1} \div (-1)$</p> <p>Use of limits 0 to 1</p> <p>$\rightarrow \frac{81\pi}{2}$ (or 127)</p>	<p>B1 B1 M1 A1</p>	<p>[4]</p>	<p>Answer without the “÷ -1 including π For “÷ -1”. Uses both limits in an integral of y^2 – if “0” ignored, M0. co (If π omitted – max 3/4)</p>
<p>(iii) $\frac{9}{2-x} = x+k$</p> <p>$\rightarrow x^2 - 2x + kx - 2k + 9 = 0$</p> <p>Uses $b - 4ac$</p> <p>$\rightarrow k^2 + 4k - 32$</p> <p>$\rightarrow$ end-points of 4 and -8</p> <p>Range for 2 points of intersection</p> <p>$\rightarrow k < -8, k > 4$.</p>	<p>M1 M1 A1 A1</p>	<p>[4]</p>	<p>Elimination of y Uses discriminant End-values correct. Accept \leq, \geq.</p>	