

MARK SCHEME for the October/November 2011 question paper

www.mymathscloud.com

for the guidance of teachers

9709 MATHEMATICS

9709/21

Paper 2, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



			hun n.
Page 2	Mark Scheme: Teachers' version	Syllabus	Papto
	GCE AS/A LEVEL – October/November 2011	9709	21 41

Cloud.com

Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Mark for a correct result or statement independent of method marks.
- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol √ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0. B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking *g* equal to 9.8 or 9.81 instead of 10.

			www.p.
Page 3	Mark Scheme: Teachers' version	Syllabus	Pap natura
	GCE AS/A LEVEL – October/November 2011	9709	
The following abbreviations may be used in a mark scheme or used on the scripts:			21 "Scioud.com

- AEF Any Equivalent Form (of answer is equally acceptable)
- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
- CWO Correct Working Only – often written by a 'fortuitous' answer
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)
- SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{}$ " marks. MR is not applied when the candidate misreads his own figures - this is regarded as an error in accuracy. An MR-2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA –1 This is deducted from A or B marks in the case of premature approximation. The PA -1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Pat	2 4
Tage 4	GCE AS/A LEVEL – October/November 2011	9709	21	1ªths
EITHER	State or imply non-modular inequality $(4 - 5x)^2 < 3^2$, or conor pair of linear equations	rresponding equat	Pap 21 ion M1	Clot
	Obtain critical values $\frac{1}{5}$ and $\frac{7}{5}$		A1	
	State correct answer $\frac{1}{5} < x < \frac{7}{5}$		A1	
OR	State one critical value, e.g. $x = \frac{1}{5}$, by solving a linear equ	uation (or inequali	ity)	
	or from a graphical method or by inspection State the other critical value correctly		B1 B1	
	State correct answer $\frac{1}{5} < x < \frac{7}{5}$		B1	[3]
Integrate a	nd obtain term of the form $k \ln(4x + 1)$		M1	
State corre	ct term $\frac{1}{2}\ln(4x+1)$		A1	
	limits correctly		M1	
	r the logarithm of a quotient or a power en answer correctly		M1 A1	[5]
Obtain der	ivative of the form $k \sec^2 2x$, where $k = 1$ or $k = \frac{1}{2}$		M1	
Use correc	rect derivative $\sec^2 2x$ t method for solving $\sec^2 2x = 4$		A1 M1	
Obtain ans	wer $x = \frac{1}{6}\pi$ (or 0.524 radians)		A1	
Obtain ans	wer $x = \frac{1}{3}\pi$ (or 1.05 radians) and no others in range		A1	[5]
	ecognizable solution method for quadratic in 3^x		M1	
	= 5 and $3^x = 2$ hmic method to solve an equation of the form $3^x = k$, where $k > 2$	0	A1 M1	
State answ State answ			A1 A1	[5]
State answ			AI	[9]
(i) Substi	tute $x = \frac{1}{2}$ and equate to 10		M1	
Obtain	answer $a = -16$		A1	
Eithe	r show that $f(3) = 0$ or divide by $(x - 3)$ obtaining a remainder of	of zero	B1	[3]
	y stage state that $x = 3$ is a solution		B1	
Attem Obtaiı	pt division by $(x - 3)$ reaching a partial quotient of $4x^2 + kx$ n quadratic factor $4x^2 - 4x - 3$		M1 A1	
	n solutions $x = \frac{3}{2}$ and $x = -\frac{1}{2}$		A1	
	2 2			

					mm.	3	
	Pa	ige 5	Mark Scheme: Teachers' version	Syllabus	Pap	2.8	Max ()
		900	GCE AS/A LEVEL – October/November 2011	9709	21	ath	Contraction of the second seco
6	(i)		sign of $x^3 - 2x^2 + 5x - 3$ at $x = 0.7$ and $x = 0.8$ the argument correctly with appropriate calculations		M1 A1	[2]	AND COULD COLL
	(ii)	Rearrange a State $a = 2$	equation to given equation or vice versa a and $b = 5$		B1 B1	[2]	
	(iii)	Obtain fina Show suffi	erative formula correctly at least once al answer 0.74 icient iterations to justify its accuracy to 2 d.p. or show there al (0.735, 0.745)	e is a sign change	M1 A1 in B1	[3]	
7	(i)	Use product Obtain corr Use $\frac{dy}{dx} = \frac{dy}{dx}$	ct rule to differentiate y rect derivative in any form in t for y		M1 A1 M1 A1	[4]	
	(ii)		$\frac{dy}{dx} = 0$ and make correct conclusion		M1 A1	[2]	
	(iii)	Substitute <i>t</i> Obtain (e^{-6})	t = -2 into equation for x or y ⁶ , $4e^{-2} + 3$)		M1 A1	[2]	
8	(i)	Make relev Obtain a co Use $\sin^2 x =$	vant use of the $cos(A + B)$ formula vant use of the $cos 2A$ and $sin 2A$ formulae orrect expression in terms of $cos x$ and $sin x$ = $1 - cos^2 x$ to obtain an expression in terms of $cos x$ ren answer correctly	Ν	M1* M1* A1 (1(dep*) A1	[5]	
	(ii)	Replace int	tegrand by $\frac{1}{2}\cos 3x + \frac{1}{2}\cos x$, or equivalent		B1		
		Integrate, c	obtaining $\frac{1}{6}\sin 3x + \frac{1}{2}\sin x$, or equivalent	В	B1 + B1		
		Use limits o Obtain give	correctly		M1 A1	[5]	