

**CAMBRIDGE**  
INTERNATIONAL EXAMINATIONS

**JUNE 2002**

**GCE Advanced Level  
GCE Advanced Subsidiary Level**

<b>MARK SCHEME</b>
<b>MAXIMUM MARK : 75</b>
<b>SYLLABUS/COMPONENT : 9709 /3, 8719 /3</b> <b>MATHEMATICS</b> <b>(Pure 3)</b>

Page 1	Mark Scheme	Syllabus	Paper
	A & AS Level Examinations – June 2002	9709, 8719	3

- 1 *EITHER:* Express LHS in terms of  $\cos\theta$  and  $\sin\theta$  or in terms of  $\tan\theta$  M1  
 Make sufficient relevant use of double-angle formula(e) M1  
 Complete proof of the result A1  
*OR:* Express RHS in terms of  $\cos\theta$  and  $\sin\theta$  or in terms of  $\tan\theta$  M1  
 Express RHS as the difference (or sum) of two fractions M1  
 Complete proof of the result A1 **3**
- [SR: an attempt ending with  $\frac{1 \cdot \tan^2\theta}{\tan\theta} = \cot\theta - \tan\theta$  earns M1 B1 only.]
- 2 *EITHER:* Show correct (unsimplified) version of the  $x$  or the  $x^2$  or the  $x^3$  term M1  
 Obtain correct first two terms  $1 + x$  A1  
 Obtain correct quadratic term  $2x^2$  A1  
 Obtain correct cubic term  $\frac{14}{3}x^3$  (allow  $\frac{28}{6}$ , 4.67, 4.66 for the coefficient) A1  
 [The M mark may be implied by correct simplified terms, if no working is shown. It is not earned by unexpanded binomial coefficients involving  $-\frac{1}{3}$ , e.g.  ${}^{-\frac{1}{3}}C_1$  or  $\binom{-\frac{1}{3}}{2}$ .]  
 [An attempt to divide 1 by the expansion of  $(1 - 3x)^{\frac{1}{3}}$  earns M1 if the expansion has a correct (unsimplified)  $x$ ,  $x^2$ , or  $x^3$  term and if the partial quotient contains a term in  $x$ . The remaining A marks are awarded as above.]
- OR:* Differentiate and calculate  $f(0)$ ,  $f'(0)$ , where  $f(x) = k(1 - 3x)^{-\frac{1}{3}}$  M1  
 Obtain correct first two terms  $1 + x$  A1  
 Obtain correct quadratic term  $2x^2$  A1  
 Obtain correct cubic term  $\frac{14}{3}x^3$  (allow  $\frac{28}{6}$ , 4.67, 4.66 for the coefficient) A1 **4**
- 3 Attempt to find  $a$  and/or quadratic factor by division or by inspection M1  
 Obtain partial quotient or factor  $x^2 - x$  A1  
 State answer  $a = 6$  B1  
 State or imply the other factor is  $x^2 - x + 3$  A1 **4**
- [The M1 is earned if division has produced a partial quotient  $x^2 + bx$ , or if inspection has an unknown factor  $x^2 + bx + c$  and has reached an equation in  $b$  and/or  $c$ .]  
 [SR: a correct division with unresolved constant remainder can earn M1A1B0A1.]  
 [NB: successive division by a pair of incorrect linear factors, e.g.  $x - 1$  and  $x + 2$  or  $x + 1$  and  $x + 2$ , can earn M1A0 or M1A1 (if their product is of the form  $x^2 + x + k$ ).]

Page 2	Mark Scheme	Syllabus	Paper
	A & AS Level Examinations – June 2002	9709, 8719	3

- 4 (i) Use the formula correctly at least once M1  
 State  $\alpha = 1.26$  as final answer A1  
 Show sufficient iterations to justify  $\alpha = 1.26$  to 2d.p., or show there is a sign change in the interval (1.255, 1.265) A1 3
- (ii) State any suitable equation in one unknown e.g.  $x = \frac{2}{3} \left( x + \frac{1}{x^2} \right)$  B1  
 State exact value of  $\alpha$  (or  $x$ ) is  $\sqrt[3]{2}$  or  $2^{\frac{1}{3}}$  B1 2
- 5 Obtain derivative  $\pm 2\sin x + k \cos 2x$  or  $\pm 2\sin x + k(\cos^2 x \pm \sin^2 x)$  M1  
 Equate derivative to zero and use trig formula to obtain an equation involving only one trig function M1  
 Obtain a correct equation of this type e.g.  $2\sin^2 x + \sin x - 1 = 0$  or  $\cos 2x = \cos \left( \frac{1}{2} \pi - x \right)$  A1  
 Obtain value  $x = \frac{1}{6} \pi$  (allow 0.524 radians or  $30^\circ$ ) A1  
 Show by any method that the corresponding point is a maximum point A1  
 Obtain second value  $x = \frac{5}{6} \pi$  (allow 2.62 radians or  $150^\circ$ ), and no others in range A1 ✓  
 Determine that it corresponds to a minimum point A1 7
- 6 (i) State or imply  $f(x) = \frac{A}{(3x+1)} + \frac{B}{(x+1)^2} + \frac{C}{(x+1)}$  B1  
 State or obtain  $A = -3$  B1  
 State or obtain  $B = 2$  B1  
 Use any relevant method to find  $C$  M1  
 Obtain  $C = 1$  A1 5  
 [Special case: allow the form  $\frac{A}{(3x+1)} + \frac{Dx+E}{(x+1)^2}$  and apply the above scheme ( $A = -3, D = 1, E = 3$ ).]  
 {SR: if  $f(x)$  is given an incomplete form of partial fractions, give B1 for a form equivalent to the omission of  $C$ , or  $E$ , or  $B$  in the above, and M1 for finding one coefficient.]
- (ii) Integrate and obtain terms  $-\ln(3x+1) - \frac{2}{(x+1)} + \ln(x+1)$  B1 + B1 + B1 ✓  
 Use limits correctly M1  
 Obtain the given answer correctly A1 5

Page 3	Mark Scheme	Syllabus	Paper
	A & AS Level Examinations – June 2002	9709, 8719	3

7	(i) State that $\frac{dm}{dt} = k(50 - m)^2$ Justify $k = 0.002$	B1 B1	2
	(ii) Separate variables and attempt to integrate $\frac{1}{(50 - m)^2}$ Obtain $\pm \frac{1}{(50 - m)}$ and $0.002t$ , or equivalent Evaluate a constant or use limits $t = 0, m = 0$ Obtain any correct form of solution e.g. $\frac{1}{(50 - m)} = 0.002t + \frac{1}{50}$ Obtain given answer correctly	M1 A1 M1 A1	
	(iii) Obtain answer $m = 25$ when $t = 10$ Obtain answer $t = 90$ when $m = 45$	A1 B1 B1	5 2
	(iv) State that $m$ approaches 50	B1	1
8	(i) State or imply a simplified direction vector of $l$ is $3i - j + 2k$ , or equivalent State equation of $l$ is $r = i + k + \lambda(3i - j + 2k)$ , or $\frac{x-1}{3} = \frac{y}{-1} = \frac{z-1}{2}$ , or equivalent Substitute in equation of $p$ and solve for $\lambda$ , or one of $x, y$ , or $z$ Obtain point of intersection $-2i + j - k$ [Any notation is acceptable.]	B1 B1 ✓ M1 A1	4
	(ii) State or imply a normal vector of $p$ is $i + 3j - 2k$ <i>EITHER:</i> Use scalar product to obtain $a + 3b - 2c = 0$ Use points on $l$ to obtain two equations in $a, b, c$ e.g. $a + c = 1, 4a - b + 3c = 1$ Solve simultaneous equations, obtaining one unknown Obtain one correct unknown e.g. $a = -\frac{2}{3}$ Obtain the other unknowns e.g. $b = \frac{4}{3}, c = \frac{5}{3}$ <i>OR:</i> Use scalar product to obtain $a + 3b - 2c = 0$ Use scalar product to obtain $3a - b + 2c = 0$ Solve simultaneous equations to obtain one ratio e.g. $a : b$ Obtain $a : b : c = 2 : -4 : -5$ , or equivalent Obtain $a = -\frac{2}{3}, b = \frac{4}{3}, c = \frac{5}{3}$	B1 M1 B1 ✓ M1 A1 A1 M1 B1 ✓ M1 A1 A1	
	[NB: candidates may transfer from the <i>EITHER</i> to <i>OR</i> scheme by subtracting the two "point" equations, or transfer from <i>OR</i> to <i>EITHER</i> by finding one of the "point" equations.]		
	<i>OR:</i> Calculate the vector product $(3i - j + 2k) \times (i + 3j - 2k)$ Obtain answer $-4i + 8j + 10k$ , or equivalent Substitute in $-4x + 8y + 10z = d$ to find $d$ , or equivalent Obtain $d = 6$ , or equivalent Obtain $a = -\frac{2}{3}, b = \frac{4}{3}, c = \frac{5}{3}$	M1 A1 ✓ M1 A1 A1	
	<i>OR:</i> State or imply a correct equation of the plane e.g. $r = \lambda(3i - j + 2k) + \mu(i + 3j - 2k) + i + k$ State 3 equations in $x, y, z, \lambda$ , and $\mu$ , e.g. $x = 3\lambda + \mu + 1, y = -\lambda + 3\mu, z = 2\lambda - 2\mu + 1$ Eliminate $\lambda$ and $\mu$ Obtain equation $-4x + 8y + 10z = 6$ , or equivalent Obtain $a = -\frac{2}{3}, b = \frac{4}{3}, c = \frac{5}{3}$	M1 A1 ✓ M1 A1 A1	6
	[SR: condone the use of $xi + yj + zk$ for $ai + bj + ck$ in the <i>EITHER</i> scheme and the first <i>OR</i> scheme.]		

Page 4	Mark Scheme	Syllabus	Paper
	A & AS Level Examinations – June 2002	9709, 8719	3

9	(i) State or imply that $r = 2$	B1	
	State or imply that $\theta = \frac{1}{3}\pi$ (allow 1.05 radians or $60^\circ$ )	B1	
	Obtain modulus 4, and argument $\frac{2}{3}\pi$ of $u^2$ (allow $2^2$ ; 2.09 or 2.10 radians or $120^\circ$ )	B1 + B1✓	
	Obtain modulus 8 and argument $\pi$ of $u^3$ (allow $2^3$ ; 3.14 or 3.15 radians or $180^\circ$ )	B1✓	5
	[Follow through on wrong $r$ and $\theta$ ]		
	[SR: if $u^2$ and $u^3$ are only given in polar form, give B1✓ for $u^2$ and B1✓ for $u^3$ .]		
(ii)	EITHER: Deduce that $u^2 - 2u + 4 = 0$ from $u^3 + 8 = 0$		
	OR: Verify that $u^2 - 2u + 4 = 0$ by calculation	B1	
	State that the other root is $1 - i\sqrt{3}$ , or equivalent	B1	2
	[NB: stating that the roots are $1 \pm i\sqrt{3}$ is sufficient for both B marks.]		
(iii)	Show both points correctly on an Argand diagram	B1	
	Show the correct relevant circle	B1	
	Show line (segment) correctly	B1	
	Shade the correct region	B1	4
	[SR: allow work on separate diagrams to be eligible for the first three B marks.]		
10	(i) State at any stage that the $x$ -coordinate of $A$ is equal to 1, or that $A$ is the point (1,0)	B1	1
	(ii) State $f'(x) = 2 \frac{\ln x}{x}$ , or equivalent	B1	
	Use product or quotient rule for the next differentiation	M1	
	Obtain $2 \cdot \frac{1}{x} \cdot \frac{1}{x} + 2 \ln x \cdot \left(\frac{-1}{x^2}\right)$ , or any equivalent correct unsimplified form	A1	
	Verify that $f''(e) = 0$	A1	4
(iii)	State or imply area is $\int_1^e (\ln x)^2 dx$	B1	
	Use $\frac{dx}{du} = e^u$ , or equivalent, in substituting for $x$ throughout	M1	
	Obtain given answer correctly (allow change of limits to be done mentally)	A1	3
(iv)	Attempt the first integration by parts, going the correct way	M1	
	Obtain $(u^2 - 2u \pm 2)e^u$ , or equivalent, after two applications of the rule	A1	
	Obtain exact answer in terms of $e$ , in any correct form, e.g. $(e - 2e + 2e) - 2$ , or $e - 2$	A1	3
	[The substitution in (iii) may be done in reverse i.e. starting with the $u$ integral and obtaining the $x$ integral. The M1A1 scheme applies, but only an explicit statement will earn the B1.]		
	[The M1A1A1 in (iv) applies to those working in terms of $x$ and obtaining $x((\ln x)^2 - 2 \ln x \pm 2)$ , or equivalent.]		