

Write your name here

Surname	Other names
---------	-------------

**Pearson Edexcel**  
**Level 3 GCE**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

# Further Mathematics

**Advanced Subsidiary**

**Further Mathematics options**

**Paper 2A: Further Pure Mathematics 1 and Further Pure Mathematics 2**

Sample Assessment Material for first teaching September 2017

**Time: 1 hour 40 minutes**

Paper Reference

**8FM0/2A**

**You must have:**

Mathematical Formulae and Statistical Tables, calculator

Total Marks

--

**Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

## Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- There are **two** sections in this question paper. Answer **all** the questions in Section A and **all** the questions in Section B.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Answers should be given to three significant figures unless otherwise stated.

## Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 80.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

## Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

S58538A

©2017 Pearson Education Ltd.

1/1/1/1



Pearson

### SECTION A

Answer ALL questions. Write your answers in the spaces provided.

1. (a) Use the substitution  $t = \tan\left(\frac{x}{2}\right)$  to show that

$$\sec x - \tan x \equiv \frac{1-t}{1+t} \quad x \neq (2n+1)\frac{\pi}{2}, n \in \mathbb{Z} \qquad (3)$$

- (b) Use the substitution  $t = \tan\left(\frac{x}{2}\right)$  and the answer to part (a) to prove that

$$\frac{1-\sin x}{1+\sin x} \equiv (\sec x - \tan x)^2 \quad x \neq (2n+1)\frac{\pi}{2}, n \in \mathbb{Z} \qquad (3)$$

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



2. The value,  $V$  hundred pounds, of a particular stock  $t$  hours after the opening of trading on a given day is modelled by the differential equation

$$\frac{dV}{dt} = \frac{V^2 - t}{t^2 + tV} \quad 0 < t < 8.5$$

A trader purchases £300 of the stock one hour after the opening of trading.

Use two iterations of the approximation formula  $\left(\frac{dy}{dx}\right)_0 \approx \frac{y_1 - y_0}{h}$  to estimate, to the nearest £, the value of the trader's stock half an hour after it was purchased.

(6)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



3. Use algebra to find the set of values of  $x$  for which

$$\frac{1}{x} < \frac{x}{x+2}$$

(6)


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



4.



**Figure 1**

Figure 1 shows a sketch of a solid sculpture made of glass and concrete. The sculpture is modelled as a parallelepiped.

The sculpture is made up of a concrete solid in the shape of a tetrahedron, shown shaded in Figure 1, whose vertices are  $O(0, 0, 0)$ ,  $A(2, 0, 0)$ ,  $B(0, 3, 1)$  and  $C(1, 1, 2)$ , where the units are in metres. The rest of the solid parallelepiped is made of glass which is glued to the concrete tetrahedron.

- (a) Find the surface area of the glued face of the tetrahedron. (4)
- (b) Find the volume of glass contained in this parallelepiped. (5)
- (c) Give a reason why the volume of concrete predicted by this model may not be an accurate value for the volume of concrete that was used to make the sculpture. (1)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





5.




Diagram not drawn to scale

Figure 2

[ You may quote without proof that for the general parabola  $y^2 = 4ax$ ,  $\frac{dy}{dx} = \frac{2a}{y}$  ]

The parabola  $C$  has equation  $y^2 = 16x$ .

(a) Deduce that the point  $P(4p^2, 8p)$  is a general point on  $C$ . (1)

The line  $l$  is the tangent to  $C$  at the point  $P$ .

(b) Show that an equation for  $l$  is  $py = x + 4p^2$  (3)

The finite region  $R$ , shown shaded in Figure 2, is bounded by the line  $l$ , the  $x$ -axis and the parabola  $C$ .

The line  $l$  intersects the directrix of  $C$  at the point  $B$ , where the  $y$  coordinate of  $B$  is  $\frac{10}{3}$

Given that  $p > 0$

(c) show that the area of  $R$  is 36 (8)

---



---



---



---



---





### SECTION B

Answer ALL questions. Write your answers in the spaces provided.

6. Given that

$$\mathbf{A} = \begin{pmatrix} 3 & 1 \\ 6 & 4 \end{pmatrix}$$

- (a) find the characteristic equation of the matrix  $\mathbf{A}$ . (2)
  
- (b) Hence show that  $\mathbf{A}^3 = 43\mathbf{A} - 42\mathbf{I}$ . (3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



7. (i) Without performing any division, explain why 8184 is divisible by 6 (2)
- (ii) Use the Euclidean algorithm to find integers  $a$  and  $b$  such that

$$27a + 31b = 1 \tag{4}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

8. A curve  $C$  is described by the equation

$$|z - 9 + 12i| = 2|z|$$

(a) Show that  $C$  is a circle, and find its centre and radius.

(4)

(b) Sketch  $C$  on an Argand diagram.

(2)

Given that  $w$  lies on  $C$ ,

(c) find the largest value of  $a$  and the smallest value of  $b$  that must satisfy

$$a \leq \text{Re}(w) \leq b$$

(2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

Question 8 continued

Lined writing area for the question.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 8 is 8 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

9. The operation \* is defined on the set  $S = \{0, 2, 3, 4, 5, 6\}$  by  $x*y = x + y = xy \pmod{7}$

*	0	2	3	4	5	6
0						
2		0				
3						5
4						
5		4				
6						

- (a) (i) Complete the Cayley table shown above
- (ii) Show that  $S$  is a group under the operation \*
- (You may assume the associative law is satisfied.)
- (6)
  
- (b) Show that the element 4 has order 3
- (2)
  
- (c) Find an element which generates the group and express each of the elements in terms of this generator.
- (3)

---

---

---

---

---

---

---

---

---

---

---

---

**Question 9 continued**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**(Total for Question 9 is 11 marks)**

10. A population of deer on a large estate is assumed to increase by 10% during each year due to natural causes.

The population is controlled by removing a constant number,  $Q$ , of the deer from the estate at the end of each year.

At the start of the first year there are 5000 deer on the estate.

Let  $P_n$  be the population of deer at the end of year  $n$ .

(a) Explain, in the context of the problem, the reason that the deer population is modelled by the recurrence relation

$$P_n = 1.1P_{n-1} - Q, \quad P_0 = 5000, \quad n \in \mathbb{Z}^+ \tag{3}$$

(b) Prove by induction that  $P_n = (1.1)^n (5000 - 10Q) + 10Q, \quad n \geq 0$  (5)

(c) Explain how the long term behaviour of this population varies for different values of  $Q$ . (2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

Question 10 continued

Lined area for writing answers to Question 10.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA







## Paper 2 Option A

### Further Pure Mathematics 1 Mark Scheme (Section A)

Question	Scheme	Marks	AOs
<b>1(a)</b>	$\sec x - \tan x = \frac{1}{1-t^2} - \frac{2t}{1-t^2}$	M1	2.1
	$= \frac{1+t^2}{1-t^2} - \frac{2t}{1-t^2} = \frac{1-2t+t^2}{1-t^2}$	M1	1.1b
	$= \frac{(1-t)^2}{(1-t)(1+t)} = \frac{1-t}{1+t} *$	A1*	2.1
		<b>(3)</b>	
<b>(b)</b>	$\frac{1-\sin x}{1+\sin x} = \frac{1-\frac{2t}{1+t^2}}{1+\frac{2t}{1+t^2}}$	M1	1.1a
	$= \frac{1+t^2-2t}{1+t^2+2t}$	M1	1.1b
	$= \frac{(1-t)^2}{(1+t)^2} = \left(\frac{1-t}{1+t}\right)^2 = (\sec x - \tan x)^2 *$	A1*	2.1
		<b>(3)</b>	

**(6 marks)**

**Notes:**

**(a)**

**M1:** Uses  $\sec x = \frac{1}{\cos x}$  and the  $t$ -substitutions for both  $\cos x$  and  $\tan x$  to obtain an expression in terms of  $t$

**M1:** Sorts out the  $\sec x$  term, and puts over a common denominator of  $1-t^2$

**A1\*:** Factorises both numerator and denominator (must be seen) and cancels the  $(1+t)$  term to achieve the answer

**(b)**

**M1:** Uses the  $t$ -substitution for  $\sin x$  in both numerator and denominator

**M1:** Multiplies through by  $1+t^2$  in numerator and denominator

**A1\*:** Factorises both numerator and denominator and makes the connection with part (a) to achieve the given result

Question	Scheme	Marks	AOs
2	£300 purchased one hour after opening $\Rightarrow V_0 = 3$ and $t_0 = 1$ ; half an hour after purchase $\Rightarrow t_2 = 1.5$ , so step $h$ required is 0.25	B1	3.3
	$t_0 = 1, V_0 = 3, \left(\frac{dV}{dt}\right)_0 \approx \frac{3^2 - 1}{1^2 + 3} = 2$	M1	3.4
	$V_1 \approx V_0 + h\left(\frac{dV}{dt}\right)_0 = 3 + 0.25 \times 2 = \dots$	M1	1.1b
	$= 3.5$	A1ft	1.1b
	$\left(\frac{dV}{dt}\right)_1 \approx \frac{3.5^2 - 1.25}{1.25^2 + 1.25 \times 3.5} \left(= \frac{176}{95}\right)$	M1	1.1b
	$V_2 \approx V_1 + h\left(\frac{dV}{dt}\right)_1 = 3.5 + 0.25 \times \frac{176}{95} = 3.963\dots$ , so £396 (nearest £)	A1	3.2a
		(6)	

(6 marks)

**Notes:**

- B1:** Identifies the correct initial conditions and requirement for  $h$
- M1:** Uses the model to evaluate  $\frac{dV}{dt}$  at  $t_0$ , using their  $t_0$  and  $V_0$
- M1:** Applies the approximation formula with their values
- A1ft:** 3.5 or exact equivalent. Follow through their step value
- M1:** Attempt to find  $\left(\frac{dV}{dt}\right)_1$  with their 3.5
- A1:** Applies the approximation and interprets the result to give £396

Question	Scheme	Marks	AOs
3	$\frac{1}{x} < \frac{x}{x+2}$		
	$\frac{(x+2)-x^2}{x(x+2)} < 0$ or $x(x+2)^2 - x^3(x+2) < 0$	M1	2.1
	$\frac{x^2-x-2}{x(x+2)} > 0 \Rightarrow \frac{(x-2)(x+1)}{x(x+2)} > 0$ or $x(x+2)(2-x)(x+1) < 0$	M1	1.1b
	At least two correct critical values from $-2, -1, 0, 2$	A1	1.1b
	All four correct critical values $-2, -1, 0, 2$	A1	1.1b
	$\{x \in \mathbb{R} : x < -2\} \cup \{x \in \mathbb{R} : -1 < x < 0\} \cup \{x \in \mathbb{R} : x > 2\}$	M1 A1	2.2a 2.5
<b>(6)</b>			
<b>(6 marks)</b>			
<b>Notes:</b>			
<p><b>M1:</b> Gathers terms on one side and puts over common denominator, or multiply by <math>x^2(x+2)^2</math> and then gather terms on one side</p> <p><b>M1:</b> Factorise numerator or find roots of numerator or factorise resulting in equation into 4 factors</p> <p><b>A1:</b> At least 2 correct critical values found</p> <p><b>A1:</b> Exactly 4 correct critical values</p> <p><b>M1:</b> Deduces that the 2 “outsides” and the “middle interval” are required. May be by sketch, number line or any other means</p> <p><b>A1:</b> Exactly 3 correct intervals, accept equivalent set notations, but must be given as a set e.g. accept <math>\mathbb{R} - ([-2, -1] \cup [0, 2])</math> or <math>\{x \in \mathbb{R} : x &lt; -2 \text{ or } -1 &lt; x &lt; 0 \text{ or } x &gt; 2\}</math></p>			

Question	Scheme	Marks	AOs
<b>4(a)</b>	Identifies glued face is triangle $ABC$ and attempts to find the area, e.g. evidences by use of $\frac{1}{2} \mathbf{AB} \times \mathbf{AC} $	M1	3.1a
	$\frac{1}{2} \mathbf{AB} \times \mathbf{AC}  = \frac{1}{2} (-2\mathbf{i} + 3\mathbf{j} + \mathbf{k}) \times (-\mathbf{i} + \mathbf{j} + 2\mathbf{k}) $	M1	1.1b
	$= \frac{1}{2} 5\mathbf{i} + 3\mathbf{j} + \mathbf{k} $	M1	1.1b
	$= \frac{1}{2}\sqrt{35}(\text{m}^2)$	A1	1.1b
		<b>(4)</b>	
	<b>Alternative</b>		
	Identifies glued face is triangle $ABC$ and attempts to find the area, e.g. evidences by use of $\frac{1}{2}\sqrt{ \mathbf{AB} ^2 \mathbf{AC} ^2 - (\mathbf{AB} \cdot \mathbf{AC})^2}$	M1	3.1a
	$ \mathbf{AB} ^2 = 4 + 9 + 1 = 14$ , $ \mathbf{AC} ^2 = 1 + 1 + 4 = 6$ and $\mathbf{AB} \cdot \mathbf{AC} = 2 + 3 + 2 = 7$	M1	1.1b
	So area of glue is $= \frac{1}{2}\sqrt{(14)(6) - (7)^2}$	M1	1.1b
	$= \frac{1}{2}\sqrt{35} (\text{m}^2)$	A1	1.1b
		<b>(4)</b>	
	<b>(b)</b>	Volume of parallelepiped taken up by concrete is e.g. $\frac{1}{6}(\mathbf{OC} \cdot (\mathbf{OA} \times \mathbf{OB}))$	M1
$= \frac{1}{6}(\mathbf{i} + \mathbf{j} + 2\mathbf{k}) \cdot (2\mathbf{i} \times (3\mathbf{j} + \mathbf{k}))$		M1	1.1b
$= \frac{10}{6} = \frac{5}{3}$		A1	1.1b
Volume of parallelepiped is $6 \times$ volume of tetrahedron ( $= 10$ ), so volume of glass is difference between these, viz. $10 - \frac{5}{3} = \dots$		M1	3.1a
Volume of glass $= \frac{25}{3}(\text{m}^3)$		A1	1.1b
		<b>(5)</b>	

Question	Scheme	Marks	AOs
	<b>4(b) Alternative</b>		
	$-\mathbf{j} + 3\mathbf{k}$ is perpendicular to both $\mathbf{OA} = 2\mathbf{i}$ and $\mathbf{OB} = 3\mathbf{j} + \mathbf{k}$	M1	3.1a
	Area $AOB = \frac{1}{2} \times  \mathbf{OA}  \times  \mathbf{OB}  = \frac{1}{2} \times 2 \times \sqrt{10} = \sqrt{10}$	A1	1.1b
	$\mathbf{i} + \mathbf{j} + 2\mathbf{k} - p(-\mathbf{j} + 3\mathbf{k}) = \mu(2\mathbf{i}) + \lambda(3\mathbf{j} + \mathbf{k}) \Rightarrow p = \frac{1}{2}$ and so height of tetrahedron is $h = \frac{1}{2}  -\mathbf{j} + 3\mathbf{k}  = \frac{1}{2} \sqrt{10}$	M1	3.1a
	Volume of glass is $V = 5 \times$ Volume of tetrahedron $= 5 \times \frac{1}{3} \sqrt{10} \times \frac{1}{2} \sqrt{10}$	M1	1.1b
	$= \frac{25}{3} (\text{m}^3)$	A1	1.1b
		<b>(5)</b>	
<b>(c)</b>	The glued surfaces may distort the shapes / reduce the volume of concrete Measurements in m may not be accurate The surface of the concrete tetrahedron may not be smooth Pockets of air may form when the concrete is being poured	B1	3.2b
		<b>(1)</b>	
<b>(10 marks)</b>			
<b>Question 4 notes:</b>			
Accept use of column vectors throughout			
<b>(a)</b>			
<b>M1:</b> Shows an understanding of what is required via an attempt at finding the area of triangle $ABC$			
<b>M1:</b> Any correct method for the triangle area is fine			
<b>M1:</b> Finds $\mathbf{AB}$ and $\mathbf{AC}$ or any other appropriate pair of vectors to use in the vector product and attempts to use them			
<b>A1:</b> Correct procedure for the vector product with at least 1 correct term $\frac{1}{2}\sqrt{35}$ or exact equivalent			
<b>(a) Alternative</b>			
<b>M1:</b> Finds two appropriate sides and attempts the scalar product and magnitudes of two of the sides			
<b>M1:</b> May use different sides to those shown			
<b>M1:</b> Correct full method to find the area of the triangle using their two sides			
<b>A1:</b> $\frac{1}{2}\sqrt{35}$ or exact equivalent			

<b>Question 4 notes continued:</b>	
<b>(b)</b>	
<b>M1:</b>	Attempts volume of concrete by finding volume of tetrahedron with appropriate method
<b>M1:</b>	Uses the formula with correct set of vectors substituted (may not be the ones shown) and vector product attempted
<b>A1:</b>	Correct value for the volume of concrete
<b>M1:</b>	Attempt to find total volume of glass by multiplying their volume of concrete by 6 and subtracting their volume of concrete. May restart to find the volume of parallelepiped
<b>A1:</b>	$\frac{25}{3}$ only, ignore reference to units
<b>(b) Alternative</b>	
<b>M1:</b>	Notes (or works out using scalar products) that $-\mathbf{j} + 3\mathbf{k}$ is a vector perpendicular to both $\mathbf{OA} = 2\mathbf{i}$ and $\mathbf{OB} = 3\mathbf{j} + \mathbf{k}$
<b>A1:</b>	Finds (using that $\mathbf{OA}$ and $\mathbf{OB}$ are perpendicular), area of $AOB = \sqrt{10}$
<b>M1:</b>	Solves $\mathbf{i} + \mathbf{j} + 2\mathbf{k} - p(-\mathbf{j} + 3\mathbf{k}) = \mu(2\mathbf{i}) + \lambda(3\mathbf{j} + \mathbf{k})$ to get the height of the tetrahedron
	$\left[ (\mu = \lambda =) p = \frac{1}{2}, \text{ so } h = \frac{1}{2}  -\mathbf{j} + 3\mathbf{k}  = \frac{1}{2} \sqrt{10} \right]$
<b>M1:</b>	Identifies the correct area as 5 times the volume of the tetrahedron (may be done as in main scheme via the difference)
<b>A1:</b>	$\frac{25}{3}$ only, ignore reference to units
<b>(c)</b>	
<b>B1:</b>	Any acceptable reason in context

Question	Scheme	Marks	AOs
<b>5(a)</b>	$y^2 = (8p)^2 = 64p^2$ and $16x = 16(4p^2) = 64p^2$ $\Rightarrow P(4p^2, 8p)$ is a general point on $C$	B1	2.2a
		<b>(1)</b>	
<b>(b)</b>	$y^2 = 16x$ gives $a = 4$ , or $2y \frac{dy}{dx} = 16$ so $\frac{dy}{dx} = \frac{8}{y}$	M1	2.2a
	$l: y - 8p = \left(\frac{8}{8p}\right)(x - 4p^2)$	M1	1.1b
	leading to $py = x + 4p^2$ *	A1*	2.1
		<b>(3)</b>	
<b>(c)</b>	$B\left(-4, \frac{10}{3}\right)$ into $l \Rightarrow \frac{10p}{3} = -4 + 4p^2$	M1	3.1a
	$6p^2 - 5p - 6 = 0 \Rightarrow (2p - 3)(3p + 2) = 0 \Rightarrow p = \dots$	M1	1.1b
	$p = \frac{3}{2}$ and $l$ cuts $x$ -axis when $\frac{3}{2}(0) = x + 4\left(\frac{3}{2}\right)^2 \Rightarrow x = \dots$	M1	2.1
	$x = -9$	A1	1.1b
	$p = \frac{3}{2} \Rightarrow P(9, 12) \Rightarrow \text{Area}(R) = \frac{1}{2}(9 - -9)(12) - \int_0^9 4x^{\frac{1}{2}} dx$	M1	2.1
	$\int 4x^{\frac{1}{2}} dx = \frac{4x^{\frac{3}{2}}}{\left(\frac{3}{2}\right)} (+c)$ or $\frac{8}{3}x^{\frac{3}{2}} (+c)$	M1	1.1b
		A1	1.1b
	$\text{Area}(R) = \frac{1}{2}(18)(12) - \frac{8}{3}\left(9^{\frac{3}{2}} - 0\right) = 108 - 72 = 36$ *	A1*	1.1b
	<b>(8)</b>		

Question	Scheme	Marks	AOs
	<b>5(c) Alternative 1</b>		
	$B\left(-4, \frac{10}{3}\right)$ into $l \Rightarrow \frac{10p}{3} = -4 + 4p^2$	M1	3.1a
	$6p^2 - 5p - 6 = 0 \Rightarrow (2p - 3)(3p + 2) = 0 \Rightarrow p = \dots$	M1	1.1b
	$p = \frac{3}{2}$ into $l$ gives $\frac{3}{2}y = x + 4\left(\frac{3}{2}\right)^2 \Rightarrow x = \dots$	M1	2.1
	$x = \frac{3}{2}y - 9$	A1	1.1b
	$p = \frac{3}{2} \Rightarrow P(9, 12) \Rightarrow \text{Area}(R) = \int_0^{12} \left( \frac{1}{16}y^2 - \left(\frac{3}{2}y - 9\right) \right) dy$	M1	2.1
	$\int \left( \frac{1}{16}y^2 - \frac{3}{2}y + 9 \right) dy = \frac{1}{48}y^3 - \frac{3}{4}y^2 + 9y (+c)$	M1	1.1b
		A1	1.1b
	$\text{Area}(R) = \left( \frac{1}{48}(12)^3 - \frac{3}{4}(12)^2 + 9(12) \right) - (0)$ $= 36 - 108 + 108 = 36 *$	A1*	1.1b
		<b>(8)</b>	
	<b>5(c) Alternative 2</b>		
	$B\left(-4, \frac{10}{3}\right)$ into $l \Rightarrow \frac{10p}{3} = -4 + 4p^2$	M1	3.1a
	$6p^2 - 5p - 6 = 0 \Rightarrow (2p - 3)(3p + 2) = 0 \Rightarrow p = \dots$	M1	1.1b
	$p = \frac{3}{2}$ and $l$ cuts $px$ -axis when $\frac{3}{2}(0) = x + 4\left(\frac{3}{2}\right)^2 \Rightarrow x = \dots$	M1	2.1
	$x = -9$	A1	1.1b
	$p = \frac{3}{2} \Rightarrow P(9, 12)$ and $x = 0$ in $l: y = \frac{2}{3}x + 6$ gives $y = 6$ $\Rightarrow \text{Area}(R) = \frac{1}{2}(9)(6) + \int_0^9 \left( \left(\frac{2}{3}x + 6\right) - \left(4x^{\frac{1}{2}}\right) \right) dx$	M1	2.1
	$\int \left( \frac{2}{3}x + 6 - 4x^{\frac{1}{2}} \right) dx = \frac{1}{3}x^2 + 6x - \frac{8}{3}x^{\frac{3}{2}} (+c)$	M1	1.1b
		A1	1.1b
	$\text{Area}(R) = 27 + \left( \left(\frac{1}{3}(9)^2 + 6(9) - \frac{8}{3}(9^{\frac{3}{2}})\right) - (0) \right)$ $= 27 + (27 + 54 - 72) = 27 + 9 = 36 *$	A1*	1.1b
		<b>(8)</b>	
<b>(12 marks)</b>			



<b>Question 5 notes:</b>	
<b>(a)</b>	<b>B1:</b> Substitutes $y_p = 8p$ into $y^2$ to obtain $64p^2$ and substitutes $x_p = 4p^2$ into $16x$ to obtain $64p^2$ and concludes that $P$ lies on $C$
<b>(b)</b>	<p><b>M1:</b> Uses the given formula to deduce the derivative. Alternatively, may differentiate using chain rule to deduce it</p> <p><b>M1:</b> Applies <math>y - 8p = m(x - 4p^2)</math>, with their tangent gradient <math>m</math>, which is in terms of <math>p</math>. Accept use of <math>8p = m(4p^2) + c</math> with a clear attempt to find <math>c</math></p> <p><b>A1*:</b> Obtains <math>py = x + 4p^2</math> by <b>cso</b></p>
<b>(c)</b>	<p><b>M1:</b> Substitutes their <math>x = "-a"</math> and <math>y = \frac{10}{3}</math> into <math>l</math></p> <p><b>M1:</b> Obtains a 3 term quadratic and solves (using the usual rules) to give <math>p = \dots</math></p> <p><b>M1:</b> Substitutes their <math>p</math> (which must be positive) and <math>y = 0</math> into <math>l</math> and solves to give <math>x = \dots</math></p> <p><b>A1:</b> Finds that <math>l</math> cuts the <math>x</math>-axis at <math>x = -9</math></p> <p><b>M1:</b> Fully correct method for finding the area of <math>R</math> i.e. <math>\frac{1}{2}(\text{their } x_p - "-9")(\text{their } y_p) - \int_0^{\text{their } x_p} 4x^2 dx</math></p> <p><b>M1:</b> Integrates <math>\pm \lambda x^{\frac{1}{2}}</math> to give <math>\pm \mu x^{\frac{3}{2}}</math>, where <math>\lambda, \mu \neq 0</math></p> <p><b>A1:</b> Integrates <math>4x^{\frac{1}{2}}</math> to give <math>\frac{8}{3}x^{\frac{3}{2}}</math>, simplified or un-simplified</p> <p><b>A1*:</b> Fully correct proof leading to a correct answer of 36</p>
<b>(c)</b>	<p><b>Alternative 1</b></p> <p><b>M1:</b> Substitutes their <math>x = "-a"</math> and <math>y = \frac{10}{3}</math> into <math>l</math></p> <p><b>M1:</b> Obtains a 3 term quadratic and solves (using the usual rules) to give <math>p = \dots</math> Substitutes their <math>p</math> (which must be positive) into <math>l</math> and rearranges to give <math>x = \dots</math></p> <p><b>M1:</b> Finds <math>l</math> as <math>x = \frac{3}{2}y - 9</math></p> <p><b>A1:</b> Fully correct method for finding the area of <math>R</math></p> <p><b>M1:</b> i.e. <math>\int_0^{\text{their } y_p} \left( \frac{1}{16}y^2 - \text{their} \left( \frac{3}{2}y - 9 \right) \right) dy</math></p> <p><b>M1:</b> Integrates <math>\pm \lambda y^2 \pm \mu y \pm \nu</math> to give <math>\pm \alpha y^3 \pm \beta y^2 \pm \nu y</math>, where <math>\lambda, \mu, \nu, \alpha, \beta \neq 0</math></p> <p><b>A1:</b> Integrates <math>\frac{1}{16}y^2 - \left( \frac{3}{2}y - 9 \right)</math> to give <math>\frac{1}{48}y^3 - \frac{3}{4}y^2 + 9y</math>, simplified or un-simplified</p> <p><b>A1*:</b> Fully correct proof leading to a correct answer of 36</p>

**Question 5 notes continued:**

(c) **Alternative 2**

**M1:** Substitutes their  $x = "-a"$  and  $y = \frac{10}{3}$  into  $l$

**M1:** Obtains a 3 term quadratic and solves (using the usual rules) to give  $p = \dots$

**M1:** Substitutes their  $p$  (which must be positive) and  $y = 0$  into  $l$  and solves to give  $x = \dots$

**A1:** Finds that  $l$  cuts the  $x$ -axis at  $x = -9$

**M1:** Fully correct method for finding the area of  $R$

$$\text{i.e. } \frac{1}{2}(\text{their } 9)(\text{their } 6) + \int_0^{\text{their } x_p} \left( \text{their } \left( \frac{2}{3}x + 6 \right) - \left( 4x^{\frac{1}{2}} \right) \right) dy$$

**M1:** Integrates  $\pm \lambda x \pm \mu \pm \nu x^{\frac{1}{2}}$  to give  $\pm \alpha x^2 \pm \mu x \pm \beta x^{\frac{3}{2}}$ , where  $\lambda, \mu, \nu, \alpha, \beta \neq 0$

**A1:** Integrates  $\left( \frac{2}{3}x + 6 \right) - \left( 4x^{\frac{1}{2}} \right)$  to give  $\frac{1}{3}x^2 + 6x - \frac{8}{3}x^{\frac{3}{2}}$ , simplified or un-simplified

**A1\*:** Fully correct proof leading to a correct answer of 36

**Further Pure Mathematics 2 Mark Scheme (Section B)**

Question	Scheme	Marks	AOs
<b>6(a)</b>	Consider $\det \begin{pmatrix} 3-\lambda & 1 \\ 6 & 4-\lambda \end{pmatrix} = (3-\lambda)(4-\lambda) - 6$	M1	1.1b
	So $\lambda^2 - 7\lambda + 6 = 0$ is characteristic equation	A1	1.1b
		<b>(2)</b>	
	So $\mathbf{A}^2 = 7\mathbf{A} - 6\mathbf{I}$	B1ft	1.1b
<b>(b)</b>	Multiplies both sides of their equation by $\mathbf{A}$ so $\mathbf{A}^3 = 7\mathbf{A}^2 - 6\mathbf{A}$	M1	3.1a
	Uses $\mathbf{A}^3 = 7(7\mathbf{A} - 6\mathbf{I}) - 6\mathbf{A}$ So $\mathbf{A}^3 = 43\mathbf{A} - 42\mathbf{I}$ *	A1*cso	1.1b
		<b>(3)</b>	
<b>(5 marks)</b>			
<b>Notes:</b>			
<b>(a)</b>			
<b>M1:</b> Complete method to find characteristic equation			
<b>A1:</b> Obtains a correct three term quadratic equation – may use variable other than $\lambda$			
<b>(b)</b>			
<b>B1ft:</b> Uses Cayley Hamilton Theorem to produce equation replacing $\lambda$ with $\mathbf{A}$ and constant term with constant multiple of identity matrix, $\mathbf{I}$			
<b>M1:</b> Multiplies equation by $\mathbf{A}$			
<b>A1*:</b> Replaces $\mathbf{A}^2$ by linear expression in $\mathbf{A}$ and achieves printed answer with no errors			

Question	Scheme	Marks	AOs
<b>7(i)</b>	Adding digits $8 + 1 + 8 + 4 = 21$ which is divisible by 3 ( or continues to add digits giving $2+1=3$ which is divisible by 3 ) so concludes that 8184 is divisible by 3	M1	1.1b
	8184 is even, so is divisible by 2 and as divisible by both 3 and 2, so it is divisible by 6	A1	1.1b
		<b>(2)</b>	
<b>(ii)</b>	Starts Euclidean algorithm $31=27 \times 1 + 4$ and $27 = 4 \times 6 + 3$	M1	1.2
	$4 = 3 \times 1 + 1$ ( so hcf = 1 )	A1	1.1b
	So $1 = 4 - 3 \times 1 = 4 - (27 - 4 \times 6) \times 1 = 4 \times 7 - 27 \times 1$	M1	1.1b
	$(31 - 27 \times 1) \times 7 - 27 \times 1 = 31 \times 7 - 27 \times 8$ $a = -8$ and $b = 7$	A1cso	1.1b
		<b>(4)</b>	
<b>(6 marks)</b>			
<b>Notes:</b>			
<b>(i)</b>			
<b>M1:</b> Explains divisibility by 3 rule in context of this number by adding digits			
<b>A1:</b> Explains divisibility by 2, giving last digit even as reason and makes conclusion that number is divisible by 6			
<b>(ii)</b>			
<b>M1:</b> Uses Euclidean algorithm showing two stages			
<b>A1:</b> Completes the algorithm. Does not need to state that hcf = 1			
<b>M1:</b> Starts reversal process, doing two stages and simplifying			
<b>A1cso:</b> Correct completion, giving clear answer following complete solution			

Question	Scheme	Marks	AOs
<b>8(a)</b>	$(x - 9)^2 + (y + 12)^2 = 4[x^2 + y^2]$	M1	2.1
	$3x^2 + 3y^2 + 18x - 24y - 225 = 0$ which is the equation of a circle	A1*	2.2a
	As $x^2 + y^2 + 6x - 8y - 75 = 0$ so $(x + 3)^2 + (y - 4)^2 = 10^2$	M1	1.1b
	Giving centre at $(-3, 4)$ and radius = 10	A1ft	1.1b
		<b>(4)</b>	
<b>(b)</b>		M1	1.1b
		A1	1.1b
		<b>(2)</b>	
<b>(c)</b>	Values range from <b>their</b> $-3 - 10$ to their $-3 + 10$	M1	3.1a
	So $-13 \leq \text{Re}(w) \leq 7$	A1ft	1.1b
		<b>(2)</b>	
<b>(8 marks)</b>			
<b>Notes:</b>			
<p><b>(a)</b>  <b>M1:</b> Obtains an equation in terms of <math>x</math> and <math>y</math> using the given information  <b>A1:</b> Expands and simplifies the algebra, collecting terms and obtains a circle equation correctly, deducing that this is a circle  <b>M1:</b> Completes the square for their equation to find centre and radius  <b>A1ft:</b> Both correct</p>			
<p><b>(b)</b>  <b>M1:</b> Draws a circle with centre and radius as given from <b>their</b> equation  <b>A1:</b> Correct circle drawn, as above, with centre at <math>-3 + 4i</math> and passing through all four quadrants</p>			
<p><b>(c)</b>  <b>M1:</b> Attempts to find where a line parallel to the real axis, passing through the centre of the circle, meets the circle so using “their <math>-3 - 10</math>” to “their <math>-3 + 10</math>”  <b>A1ft:</b> Correctly obtains the correct answer for their centre and radius</p>			

Question	Scheme	Marks	AOs																																																	
<b>9(a)(i)</b>	<table border="1"> <tr><td>*</td><td>0</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td></tr> <tr><td>0</td><td>0</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td></tr> <tr><td>2</td><td>2</td><td>0</td><td></td><td></td><td>4</td><td></td></tr> <tr><td>3</td><td>3</td><td></td><td></td><td></td><td></td><td>5</td></tr> <tr><td>4</td><td>4</td><td></td><td></td><td></td><td></td><td></td></tr> <tr><td>5</td><td>5</td><td>4</td><td></td><td></td><td></td><td></td></tr> <tr><td>6</td><td>6</td><td></td><td>5</td><td></td><td></td><td></td></tr> </table>	*	0	2	3	4	5	6	0	0	2	3	4	5	6	2	2	0			4		3	3					5	4	4						5	5	4					6	6		5				M1	1.1b
	*	0	2	3	4	5	6																																													
0	0	2	3	4	5	6																																														
2	2	0			4																																															
3	3					5																																														
4	4																																																			
5	5	4																																																		
6	6		5																																																	
<table border="1"> <tr><td>*</td><td>0</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td></tr> <tr><td>0</td><td>0</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td></tr> <tr><td>2</td><td>2</td><td>0</td><td>6</td><td>5</td><td>4</td><td>3</td></tr> <tr><td>3</td><td>3</td><td>6</td><td>4</td><td>2</td><td>0</td><td>5</td></tr> <tr><td>4</td><td>4</td><td>5</td><td>2</td><td>6</td><td>3</td><td>0</td></tr> <tr><td>5</td><td>5</td><td>4</td><td>0</td><td>3</td><td>6</td><td>2</td></tr> <tr><td>6</td><td>6</td><td>3</td><td>5</td><td>0</td><td>2</td><td>4</td></tr> </table>	*	0	2	3	4	5	6	0	0	2	3	4	5	6	2	2	0	6	5	4	3	3	3	6	4	2	0	5	4	4	5	2	6	3	0	5	5	4	0	3	6	2	6	6	3	5	0	2	4	M1 A1	1.1b 1.1b	
*	0	2	3	4	5	6																																														
0	0	2	3	4	5	6																																														
2	2	0	6	5	4	3																																														
3	3	6	4	2	0	5																																														
4	4	5	2	6	3	0																																														
5	5	4	0	3	6	2																																														
6	6	3	5	0	2	4																																														
<b>(ii)</b>	Identity is zero and there is closure as shown above	M1	2.1																																																	
	3 and 5 are inverses, 4 and 6 are inverses, 2 is self-inverse, 0 is identity so is self-inverse	M1	2.5																																																	
	Associative law may be assumed so $S$ forms a group	A1	1.1b																																																	
		<b>(6)</b>																																																		
<b>(b)</b>	$4*4*4 = 4*(4*4) = 4*6$ or $4*4*4 = (4*4)*4 = 6*4$	M1	2.1																																																	
	$= 0$ (the identity) so 4 has order 3	A1	2.2a																																																	
		<b>(2)</b>																																																		
<b>(c)</b>	3 and 5 each have order 6 so either generates the group	M1	3.1a																																																	
	<b>Either</b> $3^1 = 3, 3^2 = 4, 3^3 = 2, 3^4 = 6, 3^5 = 5, 3^6 = 0$	A1, A1	1.1b																																																	
	<b>Or</b> $5^1 = 5, 5^2 = 6, 5^3 = 2, 5^4 = 4, 5^5 = 3, 5^6 = 0$		1.1b																																																	
	<b>(3)</b>																																																			
<b>(11 marks)</b>																																																				

<b>Question 9 notes:</b>	
<b>(a)(i)</b>	<p><b>M1:</b> Begins completing the table – obtaining correct first row and first column and using symmetry</p> <p><b>M1:</b> Mostly correct – three rows or three columns correct (so demonstrates understanding of using *)</p> <p><b>A1:</b> Completely correct</p>
<b>(a)(ii)</b>	<p><b>M1:</b> States closure and identifies the identity as zero</p> <p><b>M1:</b> Finds inverses for each element</p> <p><b>A1:</b> States that associative law is satisfied and so all axioms satisfied and <math>S</math> is a group</p>
<b>(b)</b>	<p><b>M1:</b> Clearly begins process to find <math>4*4*4</math> reaching <math>6*4</math> or <math>4*6</math> with clear explanation</p> <p><b>A1:</b> Gives answer as zero, states identity and deduces that order is 3</p>
<b>(c)</b>	<p><b>M1:</b> Finds either 3 or 5 or both</p> <p><b>A1:</b> Expresses four of the six terms as powers of either generator correctly (may omit identity and generator itself)</p> <p><b>A1:</b> Expresses all six terms correctly in terms of either 3 or 5 (Do not need to give both)</p>

Question	Scheme	Marks	AOs
<b>10(a)</b>	$P_{n-1}$ is the population at the end of year $n - 1$ and this is increased by 10% by the end of year $n$ , so is multiplied by 110% = 1.1 to give $1.1 \times P_{n-1}$ as new population by natural causes	B1	3.3
	$Q$ is subtracted from $1.1 \times P_{n-1}$ as $Q$ is the number of deer removed from the estate	B1	3.4
	So $P_n = 1.1P_{n-1} - Q$ , $P_0 = 5000$ as population at start is 5000 and $n \in \mathbb{Z}^+$	B1	1.1b
		<b>(3)</b>	
<b>(b)</b>	Let $n = 0$ , then $P_0 = (5000 - 10Q)(1.1)^0 + 10Q = 5000$ so result is true when $n = 0$	B1	2.1
	Assume result is true for $n = k$ , $P_k = (1.1)^k (5000 - 10Q) + 10Q$ , then as $P_{k+1} = 1.1P_k - Q$ , so $P_{k+1} = \dots$	M1	2.4
	$P_{k+1} = 1.1 \times 1.1^k (5000 - 10Q) + 1.1 \times 10Q - Q$	A1	1.1b
	So $P_{k+1} = (5000 - 10Q)(1.1)^{k+1} + 10Q$ ,	A1	1.1b
	Implies result holds for $n = k + 1$ and so by induction $P_n = (5000 - 10Q)(1.1)^n + 10Q$ , is true for all integer $n$	B1	2.2a
		<b>(5)</b>	
<b>(c)</b>	For $Q < 500$ the population of deer will grow, for $Q > 500$ the population of deer will fall	B1	3.4
	For $Q = 500$ the population of deer remains steady at 5000,	B1	3.4
		<b>(2)</b>	
<b>(10 marks)</b>			
<b>Notes:</b>			
<b>(a)</b>			
<b>B1:</b> Need to see 10% increase linked to multiplication by scale factor 1.1			
<b>B1:</b> Needs to explain that subtraction of $Q$ indicates the removal of $Q$ deer from population			
<b>B1:</b> Needs complete explanation with mention of $P_n = 1.1P_{n-1} - Q$ , $P_0 = 5000$ being the initial number of deer			
<b>(b)</b>			
<b>B1:</b> Begins proof by induction by considering $n = 0$			
<b>M1:</b> Assumes result is true for $n = k$ and uses iterative formula to consider $n = k + 1$			
<b>A1:</b> Correct algebraic statement			
<b>A1:</b> Correct statement for $k + 1$ in required form			
<b>B1:</b> Completes the inductive argument			
<b>(c)</b>			
<b>B1:</b> Consideration of both possible ranges of values for $Q$ as listed in the scheme			
<b>B1:</b> Gives the condition for the steady state			