

Mark Scheme (Results)

Summer 2012

GCE Further Pure FP3
(6669) Paper 1

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at:

www.pearson.com/uk

Summer 2012

Publications Code UA032243

All the material in this publication is copyright

© Pearson Education Ltd 2012

June 2012
6669 Further Pure Maths FP3
Mark Scheme

Question Number	Scheme	Marks
1. (a)	Uses formula to obtain $e = \frac{5}{4}$	M1A1
	Uses ae formula	M1 (3)
(b)	Uses other formula $\frac{a}{e}$	M1
	Obtains both Foci are $(\pm 5, 0)$ and Directrices are $x = \pm \frac{16}{5}$ (needs both method marks)	A1 cso (2) (5 marks)

Notes

a1M1: Uses $b^2 = a^2(e^2 - 1)$ to get $e > 1$

a1A1: cao

a2M1: Uses ae

b1M1: Uses $\frac{a}{e}$

b1A1: cso for both foci and both directrices. Must have both of the 2 previous M marks may be implicit.

Question Number	Scheme	Marks
2.	$\frac{dy}{dx} = \sinh 3x$ $\text{so } s = \int \sqrt{1 + \sinh^2 3x} dx$ $\therefore s = \int \cosh 3x dx$ $= \left[\frac{1}{3} \sinh 3x \right]_b^{ln a}$ $= \frac{1}{3} \sinh 3 \ln a = \frac{1}{6} [e^{3 \ln a} - e^{-3 \ln a}]$ $= \frac{1}{6} \left(a^3 - \frac{1}{a^3} \right) \quad (\text{so } k = 1/6)$	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>DM1</p> <p>A1 (6 marks)</p>

Notes

1B1: cao

1M1: Use of arc length formula, need both $\sqrt{\quad}$ and $\left(\frac{dy}{dx}\right)^2$.

1A1: $\int \cosh 3x dx$ cao

2M1: Attempt to integrate, getting a hyperbolic function o.e.

3M1: depends on previous M mark. Correct use of $\ln a$ and 0 as limits. Must see some exponentials.

2A1: cao

Question Number	Scheme	Marks
3. (a)	$\vec{AC} = 3\mathbf{i} + 6\mathbf{j} + 2\mathbf{k}, \quad \vec{BC} = -3\mathbf{i} + 4\mathbf{j} + 3\mathbf{k}$ $\vec{AC} \times \vec{BC} = 10\mathbf{i} - 15\mathbf{j} + 30\mathbf{k}$	B1, B1 M1 A1 (4)
(b)	$\text{Area of triangle } ABC = \frac{1}{2} 10\mathbf{i} - 15\mathbf{j} + 30\mathbf{k} = \frac{1}{2} \sqrt{1225} = 17.5$	M1 A1 (2)
(c)	Equation of plane is $10x - 15y + 30z = -20$ or $2x - 3y + 6z = -4$ So $\mathbf{r} \cdot (2\mathbf{i} - 3\mathbf{j} + 6\mathbf{k}) = -4$ or correct multiple	M1 A1 (2) (8 marks)

Notes

a1B1: $\vec{AC} = 3\mathbf{i} + 6\mathbf{j} + 2\mathbf{k}$ cao, any form

a2B1: $\vec{BC} = -3\mathbf{i} + 4\mathbf{j} + 3\mathbf{k}$ cao, any form

a1M1: Attempt to find cross product, modulus of one term correct.

a1A1: cao, any form.

b1M1: modulus of their answer to (a) – condone missing $\frac{1}{2}$ here. To finding area of triangle by correct method.

b1A1: cao.

c1M1: [Using their answer to (a) to] find **equation** of plane. Look for **a.n** or **b.n** or **c.n** for p.

c1A1: cao

Question Number	Scheme	Marks
4(a)	$I_n = \left[x^n \left(-\frac{1}{2} \cos 2x \right) \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} -\frac{1}{2} n x^{n-1} \cos 2x dx$ <p>so</p> $I_n = \left\langle \left[x^n \left(-\frac{1}{2} \cos 2x \right) \right]_0^{\frac{\pi}{4}} \right\rangle + \left[\frac{1}{4} n x^{n-1} \sin 2x \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} \frac{1}{4} n(n-1) x^{n-2} \sin 2x dx$ <p>i.e. $I_n = \frac{1}{4} n \left(\frac{\pi}{4} \right)^{n-1} - \frac{1}{4} n(n-1) I_{n-2} *$</p>	<p>M1 A1</p> <p>M1 A1</p> <p>A1cso</p> <p>(5)</p>
(b)	$I_0 = \int_0^{\frac{\pi}{4}} \sin 2x dx = \left[-\frac{1}{2} \cos 2x \right]_0^{\frac{\pi}{4}} = \frac{1}{2}$ $I_2 = \frac{1}{4} \times 2 \times \left(\frac{\pi}{4} \right) - \frac{1}{4} \times 2 \times I_0, \text{ so } I_2 = \frac{\pi}{8} - \frac{1}{4}$	<p>M1 A1</p> <p>M1 A1</p> <p>(4)</p>
(c)	$I_4 = \left(\frac{\pi}{4} \right)^3 - \frac{1}{4} \times 4 \times 3 I_2 = \frac{\pi^3}{64} - 3 \left(\frac{\pi}{8} - \frac{1}{4} \right) = \frac{1}{64} (\pi^3 - 24\pi + 48) *$	<p>M1 A1cso</p> <p>(2)</p>

Notes

a1M1: Use of integration by parts, integrating $\sin 2x$, differentiating x^n .

a1A1: cao

a2M1: Second application of integration by parts, integrating $\cos 2x$, differentiating x^{n-1} .

a2A1: cao

a3A1: cso Including correct use of $\frac{\pi}{4}$ and 0 as limits.

b1M1: Integrating to find I_0 or setting up parts to find I_2 .

b1A1: cao (Accept $I_0 = \frac{1}{2}$ here for both marks)

b2M1: Finding I_2 in terms of π . If 'n's left in M0

b2A1: cao

c1M1: Finding I_4 in terms of I_2 then in terms of π . If 'n's left in M0

c1A1: cso

Question Number	Scheme	Marks
5. (a)	$\operatorname{ar sinh} 2x, +x \frac{2}{\sqrt{1+4x^2}}$	M1A1, A1 (3)
(b)	$\begin{aligned} \therefore \int_0^{\sqrt{2}} \operatorname{arsinh} 2x dx &= [x \operatorname{ar sinh} 2x]_0^{\sqrt{2}} - \int_0^{\sqrt{2}} \frac{2x}{\sqrt{1+4x^2}} dx \\ &= [x \operatorname{ar sinh} 2x]_0^{\sqrt{2}} - \left[\frac{1}{2} (1+4x^2)^{\frac{1}{2}} \right]_0^{\sqrt{2}} \\ &= \sqrt{2} \operatorname{arsinh} 2\sqrt{2} - \left[\frac{3}{2} - \frac{1}{2} \right] \\ &= \sqrt{2} \ln(3+2\sqrt{2}) - 1 \end{aligned}$	1M1 1A1ft 2M1 2A1 3DM1 4M1 3A1 (7) (10 marks)

Notes

a1M1: Differentiating getting an arsinh term **and** a term of the form $\frac{px}{\sqrt{1 \pm qx^2}}$

a1A1: cao $\operatorname{ar sinh} 2x$

a2A1: cao $+ \frac{2x}{\sqrt{1+4x^2}}$

b1M1: rearranging their answer to (a). **OR** setting up parts

b1A1: ft from their (a) **OR** setting up parts correctly

b2M1: Integrating getting an arsinh or arcosh term **and** a $(1 \pm ax^2)^{\frac{1}{2}}$ term o.e..

b2A1: cao

b3DM1: depends on previous M, correct use of $\sqrt{2}$ and 0 as limits.

b4M1: converting to log form.

b3A1: cao depends on all previous M marks.

Question Number	Scheme	Marks
6(a)	$\frac{2x}{a^2} + \frac{2y}{b^2} \frac{dy}{dx} = 0 \quad \text{and so} \quad \frac{dy}{dx} = -\frac{xb^2}{ya^2} = -\frac{b \cos \theta}{a \sin \theta}$ $\therefore y - b \sin \theta = -\frac{b \cos \theta}{a \sin \theta} (x - a \cos \theta)$ <p>Uses $\cos^2 \theta + \sin^2 \theta = 1$ to give $\frac{x \cos \theta}{a} + \frac{y \sin \theta}{b} = 1$ *</p>	<p>M1 A1</p> <p>M1</p> <p>A1cso</p> <p>(4)</p>
(b)	<p>Gradient of circle is $-\frac{\cos \theta}{\sin \theta}$ and equation of tangent is</p> $y - a \sin \theta = -\frac{\cos \theta}{\sin \theta} (x - a \cos \theta) \quad \text{or sets } a = b \text{ in previous answer}$ <p>So $y \sin \theta + x \cos \theta = a$</p>	<p>M1</p> <p>A1</p> <p>(2)</p>
(c)	<p>Eliminate x or y to give $y \sin \theta (\frac{a}{b} - 1) = 0$ or $x \cos \theta (\frac{b}{a} - 1) = b - a$</p> <p>$l_1$ and l_2 meet at $(\frac{a}{\cos \theta}, 0)$</p>	<p>M1</p> <p>A1, B1</p> <p>(3)</p>
(d)	<p>The locus of R is part of the line $y = 0$, such that $x \geq a$ and $x \leq -a$</p> <p>Or clearly labelled sketch.</p> <p>Accept "real axis"</p>	<p>B1, B1</p> <p>(2)</p> <p>(11 marks)</p>

Notes

a1M1: Finding gradient in terms of θ . Must use calculus.

a1A1: cao

a2M1: Finding equation of tangent

a2A1: cso (answer given). Need to get $\cos^2 \theta + \sin^2 \theta$ on the same side.

b1M1: Finding gradient and equation of tangent, **or** setting $a = b$.

b1A1: cao need not be simplified.

c1M1: As scheme

c1A1: $x = \frac{a}{\cos \theta}$, need not be simplified.

c1B1: $y = 0$, need not be simplified.

d1B1: Identifying locus as $y = 0$ or real/'x' axis.

d2B1: Depends on previous B mark, identifies correct parts of $y = 0$. Condone use of strict inequalities.

Question Number	Scheme	Marks
7(a)	$f(x) = 5\cosh x - 4\sinh x = 5 \times \frac{1}{2}(e^x + e^{-x}) - 4 \times \frac{1}{2}(e^x - e^{-x})$ $= \frac{1}{2}(e^x + 9e^{-x}) \quad *$	M1 A1cso (2)
(b)	$\frac{1}{2}(e^x + 9e^{-x}) = 5 \Rightarrow e^{2x} - 10e^x + 9 = 0$ So $e^x = 9$ or 1 and $x = \ln 9$ or 0	M1 A1 M1 A1 (4)
(c)	Integral may be written $\int \frac{2e^x}{e^{2x} + 9} dx$ This is $\frac{2}{3} \arctan\left(\frac{e^x}{3}\right)$ Uses limits to give $\left[\frac{2}{3} \arctan 1 - \frac{2}{3} \arctan\left(\frac{1}{\sqrt{3}}\right)\right] = \left[\frac{2}{3} \times \frac{\pi}{4} - \frac{2}{3} \times \frac{\pi}{6}\right] = \frac{\pi}{18} *$	B1 M1 A1 DM1 A1cso (5) (11 marks)

Notes

a1M1: Replacing both coshx and sinhx by terms in e^x and e^{-x} condone sign errors here.

a1A1: cso (answer given)

b1M1: Getting a three term quadratic in e^x

b1A1: cao

b2M1: solving to $x =$

b2A1: cao need $\ln 9$ (o.e) and 0 (not $\ln 1$)

c1B1: cao getting into suitable form, may substitute first.

c1M1: Integrating to give term in arctan

c1A1: cao

c2M1: Depends on previous M mark. Correct use of $\ln 3$ and $\frac{1}{2} \ln 3$ as limits.

c2A1: cso must see them subtracting two terms in π .

Question Number	Scheme	Marks
8. (a)	$\begin{vmatrix} 2-\lambda & 1 & 0 \\ 1 & 2-\lambda & 0 \\ -1 & 0 & 4-\lambda \end{vmatrix} = 0 \therefore (2-\lambda)(2-\lambda)(4-\lambda) - (4-\lambda) = 0$ <p>(4 - λ) = 0 verifies λ = 4 is an eigenvalue (can be seen anywhere)</p> <p>∴ (4 - λ){4 - 4λ + λ² - 1} = 0 ∴ (4 - λ){λ² - 4λ + 3} = 0</p> <p>∴ (4 - λ)(λ - 1)(λ - 3) = 0 and 3 and 1 are the other two eigenvalues</p>	M1 M1 A1 M1 A1 (5)
(b)	<p>Set $\begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ -1 & 0 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = 4 \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ or $\begin{pmatrix} -2 & 1 & 0 \\ 1 & -2 & 0 \\ -1 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$</p> <p>Solve -2x + y = 0 and x - 2y = 0 and -x = 0 to obtain x = 0, y = 0, z = k</p> <p>Obtain eigenvector as k (or multiple)</p>	M1 M1 A1 (3)
(c)	<p>l_1 has equation which may be written $\begin{pmatrix} 3 + \lambda \\ 2 - \lambda \\ -2 + 2\lambda \end{pmatrix}$</p> <p>So l_2 is given by $\mathbf{r} = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ -1 & 0 & 4 \end{pmatrix} \begin{pmatrix} 3 + \lambda \\ 2 - \lambda \\ -2 + 2\lambda \end{pmatrix}$</p> <p>i.e. $\mathbf{r} = \begin{pmatrix} 8 + \lambda \\ 7 - \lambda \\ -11 + 7\lambda \end{pmatrix}$</p> <p>So $(\mathbf{r} - \mathbf{c}) \times \mathbf{d} = \mathbf{0}$ where $\mathbf{c} = 8\mathbf{i} + 7\mathbf{j} - 11\mathbf{k}$ and $\mathbf{d} = \mathbf{i} - \mathbf{j} + 7\mathbf{k}$</p>	B1 M1 M1 A1 A1ft (5) (13 marks)

Notes

a1M1: Condone missing = 0. (They might expand the determinant using any row or column)

a2M1: Shows λ = 4 is an eigenvalue. Some working needed need to see = 0 at some stage.

a1A1: Three term quadratic factor cao, may be implicit (this A depends on 1st M only)

a2M1: Attempt at factorisation (usual rules), solving to λ = .

a2A1: cao. If they state λ = 1 and 3 please give the marks.

b1M1: Using $\mathbf{Ax} = 4\mathbf{x}$ o.e.

b2M1: Getting a pair of correct equations.

b1A1: cao

c1B1: Using **a** and **b**.

c1M1: Using $\mathbf{r} = \mathbf{Mx}$ their matrix in **a** and **b**.

c2M1: Getting an expression for l_2 with at least one component correct.

c1A1: cao all three components correct

c2A1ft: ft their vector, must have $\mathbf{r} = \text{or } (\mathbf{r} - \mathbf{c}) \times \mathbf{d} = \mathbf{0}$ need both equation and r.

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com
Order Code UA032243 Summer 2012

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual



Llywodraeth Cynulliad Cymru
Welsh Assembly Government

