



EDEXCEL STATISTICS S1 (6683) - NOVEMBER 2002  
PROVISIONAL MARK SCHEME

Question Number	Scheme	Marks
1.	<p>(a) Statistical models allow problems to be solved without the need to construct a full-scale physical model, saving time/expense. They allow parameters to be changed and refinements to be made quickly.</p> <p>(b) (i) Normal; (ii) Discrete uniform</p>	<p>B2, 1, 0 (2)</p> <p>B1, B1 (2)</p> <p><b>(4 marks)</b></p>
2.	<p>(a) 60A, 40S, 2M</p> $P(\text{all only arts}) = \frac{60}{125} \times \frac{59}{124} \times \frac{58}{123} = \frac{3422}{31775} = 0.10769\dots$ <p>(b) <math>P(\text{exactly one only science}) = 3 \times \frac{40}{125} \times \frac{85}{124} \times \frac{84}{123}</math></p> $= \frac{2856}{6355} = 0.44940\dots$	<p>B1</p> <p>M1 A1 A1 (4)</p> <p>B1</p> <p>M1 A1 (3)</p> <p><b>(7 marks)</b></p>
3.	<p>(a) <math>P(A \cap B) = P(A)P(B) = 0.25 \times 0.30 = 0.075</math></p> <p>(b) <math>P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.25 + 0.30 - 0.075 = 0.475</math></p> <p>(c) <math>P(A \cap B') = \frac{P(A \cap B')}{P(B')} = \frac{P(A) - P(A \cap B)}{1 - P(B)}</math></p> $= \frac{0.25 - 0.075}{1 - 0.3}$ $= 0.25$	<p>M1 A1 (2)</p> <p>M1</p> <p>A1 (2)</p> <p>M1</p> <p>M1 A1ft</p> <p>A1 (4)</p> <p><b>(8 marks)</b></p>

EDEXCEL STATISTICS S1 (6683) - Nov 2002  
PROVISIONAL MARK SCHEME

Question Number	Scheme	Marks
<p>4. (a)</p> 	<p> <math>P(L &gt; 50.98) = 0.025</math>  <math>P\left[Z &gt; \frac{50.98 - \alpha}{0.5}\right] = 0.025</math>  <math>\therefore \frac{50.98 - \alpha}{0.5} = 1.96</math>  <math>\therefore \alpha = 50 \text{ (*)}</math> </p> <p> <math>L \sim N(50, 0.5^2)</math> </p> <p>           (b) <math>P(49.25 &lt; L &lt; 50.75) = P\left[\frac{49.25 - 50}{0.5} &lt; Z &lt; \frac{50.75 - 50}{0.5}\right]</math>  <math>= P(-1.5 &lt; Z &lt; 1.5) \quad -1.5 \text{ \&amp; } +1.5</math>  <math>= 2\Phi(1.5) - 1</math>  <math>= 0.8664</math> </p> <p>           (c) <math>P(\text{Both}) = (1 - 0.8664)^2</math>  <math>= 0.01784\dots</math> </p>	<p>B1 M1 A1 M1 A1 (5)</p> <p>M1 A1 M1 A1 (4)</p> <p>M1 A1 (2)</p> <p>(11 marks)</p>
<p>5. (a)</p>	<p> <math>S_{ss} = 108.07875; S_{st} = 129.1675</math>  <math>q = \frac{S_{st}}{S_{ss}} = \frac{129.1675}{108.07875} = 1.1951239\dots</math>  <math>p = \frac{65.0}{8} - (1.951239\dots) \times \frac{48.5}{8} = 0.879561\dots</math>  <math>\therefore t = 0.879561\dots + 1.1951259\dots S</math> </p> <p>           (b) <math>y - 20 = 0.879561\dots + 1.1951239\dots(x - 6)</math>  <math>\therefore y = 13.709 + 1.195x</math> </p> <p>           (c) 0.943; the pmcc is an index (no units) and is not affected by linear transformations of either/both variables         </p>	<p>B1; B1 M1, A1 M1, A1 A1 ft (7) M1, A1 ft A1 (3) B1; B1 (2)</p> <p>(12 marks)</p>

EDEXCEL STATISTICS S1 (6683) - NOVEMBER 2002  
PROVISIONAL MARK SCHEME

Question Number	Scheme	Marks
6.	<p>(a) <math>\alpha + \beta = 0.5</math>  <math>-2\alpha + 2\beta = -0.2</math>  <math>\therefore \alpha = 0.3, \beta = 0.2</math></p> <p>(b) <math>F(0.8) = 0.6</math></p> <p>(c) <math>E(X^2) = (4 \times 0.3) + \dots + (4 \times 0.2), = 2.4</math>  <math>\therefore \text{Var}(X) = 2.4 - (-0.2)^2, = 2.36</math></p> <p>(d) <math>E(3X - 2) = 3E(X) - 2, = -2.6</math></p> <p>(e) <math>\text{Var}(2X + 6) = 4 \text{Var}(X), = 9.44</math></p>	<p>B1  M1  M1 A1; A1 <b>(6)</b>  B1 ft <b>(1)</b>  M1, A1  M1, A1 <b>(4)</b>  M1, A1 ft <b>(2)</b>  M1, A1 ft <b>(2)</b>  <b>(15 marks)</b></p>
7.	<p>(a) Mode = 78</p> <p>(b) <math>Q_1 = 56; Q_2 = 70; Q_3 = 78</math></p> <p>(c) <math>(Q_3 - Q_1) = 22</math>  <math>Q_1 - 1.0(Q_3 - Q_1) = 34 \Rightarrow 31 \text{ \&amp; } 31 \text{ are outliers}</math>  <math>Q_3 + 1.0(Q_3 - Q_1) = 100 \Rightarrow \text{no outliers}</math></p> <p>(d) <i>(accurate sketch on graph paper required)</i></p> <div style="display: flex; align-items: center; justify-content: center;">  <div style="margin-left: 20px;"> <p>boxplot</p> <p>scales and labels</p> <p><math>Q_1, Q_2, Q_3</math></p> <p>31, 32, 34 (39), 92</p> </div> </div>	<p>B1 <b>(1)</b>  B1; B1; B1 <b>(3)</b>  M1 A1  A1 <b>(3)</b>  M1  B1  A1  A1 <b>(4)</b>  B1  M1  A1 <b>(3)</b>  M1, A1  M1, A1 <b>(4)</b>  <b>(18 marks)</b></p>
(e)	$\alpha = \frac{3363}{50} = 67.26$ $\sigma^2 = \frac{238305}{50} - (67.26)^2 = 242.1924$ $\therefore \sigma = \sqrt{242.1924} = 15.56253\dots$	<p>B1  M1  A1 <b>(3)</b></p>
(f)	$(Q_3 - Q_2) < (Q_2 - Q_1)$ , i.e. $8 < 14 \Rightarrow \text{negative skew}$ Mean < Median < Mode, i.e. $67.26 < 70 < 78 \Rightarrow \text{negative skew}$	<p>M1, A1  M1, A1 <b>(4)</b>  <b>(18 marks)</b></p>