

52 512

1. A manufacturer produces sweets of length  $L$  mm where  $L$  has a continuous uniform distribution with range  $[15, 30]$ .


(a) Find the probability that a randomly selected sweet has a length greater than 24 mm. (2)

These sweets are randomly packed in bags of 20 sweets.

(b) Find the probability that a randomly selected bag will contain at least 8 sweets with length greater than 24 mm. (3)

(c) Find the probability that 2 randomly selected bags will both contain at least 8 sweets with length greater than 24 mm. (2)

$$a) L \sim U[15, 30] \quad P(X > 24) = \frac{6}{15} = 0.4$$



$$b) X \sim B(20, \frac{6}{15}) \quad P(X \geq 8) = 1 - P(X \leq 7) = 0.5841$$

$$b) 0.5841^2 = 0.341$$

Leave blank

2. A test statistic has a distribution  $B(25, p)$ .

Given that

$$H_0: p = 0.5 \quad H_1: p \neq 0.5$$

(a) find the critical region for the test statistic such that the probability in each tail is as close as possible to 2.5%.

(b) State the probability of incorrectly rejecting  $H_0$  using this critical region. (2)

$$P(X \leq L) \approx 0.025 \quad P(X \geq H) \approx 0.025$$

$$P(X \leq 7) = 0.0216 \quad P(X > H-1) \approx 0.025$$

$$P(X \leq 8) = 0.0539 \quad 1 - P(X \leq H-1) \approx 0.025$$

$$\therefore L = 7 \quad P(X \leq H-1) \approx 0.975$$

$$CR \{X \leq 7\} \cup \{X \geq 18\} \quad P(X \leq 17) = 0.9784$$

$$b) ASL = 0.0216 + 0.0216 = 0.0432$$

4.32% chance of incorrectly rejecting  $H_0$

3. (a) Write down two conditions needed to approximate the binomial distribution by the Poisson distribution.

(2)

A machine which manufactures bolts is known to produce 3% defective bolts. The machine breaks down and a new machine is installed. A random sample of 200 bolts is taken from those produced by the new machine and 12 bolts were defective.

- (b) Using a suitable approximation, test at the 5% level of significance whether or not the proportion of defective bolts is higher with the new machine than with the old machine. State your hypotheses clearly.

(7)

3a) large  $n$ , small  $p \approx np \leq 10$

$$x \sim B(200, 0.03) \quad np = 6 \approx x \sim P_0(6)$$

$$H_0: \lambda = 6 \quad P(x > 12) \quad P(x > 11) = 1 - P(x \leq 11)$$

$$H_1: \lambda > 6 \quad = 0.0201 < 0.05$$

$\therefore$  there is enough evidence to reject null hypothesis since result is significant  
 $\therefore$  evidence to suggest the proportion of faulty bolts has increased.

4. The number of houses sold by an estate agent follows a Poisson distribution with a mean of 2 per week.

- (a) Find the probability that in the next 4 weeks the estate agent sells

- (i) exactly 3 houses,  
 (ii) more than 5 houses.

The estate agent monitors sales in periods of 4 weeks.

- (b) Find the probability that in the next twelve of these 4 week periods there are exactly nine periods in which more than 5 houses are sold.

(3)

The estate agent will receive a bonus if he sells more than 25 houses in the next 10 weeks.

- (c) Use a suitable approximation to estimate the probability that the estate agent receives a bonus.

$$a) x \sim P_0(8) \quad P(x=3) = \frac{e^{-8} \times 8^3}{3!} = 0.0286$$

$$ii) P(x > 5) = 1 - P(x \leq 5) = 0.8088$$

$$b) y \sim B(12, 0.8088)$$

$$P(y=9) = \binom{12}{9} 0.8088^9 0.1912^3 = 0.2277$$

$$c) t \sim P_0(20) \quad \mu = 20 \quad \sigma^2 = 20 \approx N(20, 20)$$

$$P(t > 25) \approx P(t > 25.5) \approx P\left(z > \frac{25.5 - 20}{\sqrt{20}}\right)$$

$$\approx P(z > 1.23) = 1 - \Phi(1.23) = 0.1093$$

5. The queuing time,  $X$  minutes, of a customer at a till of a supermarket has probability density function

$$f(x) = \begin{cases} \frac{3}{32}x(k-x) & 0 \leq x \leq k \\ 0 & \text{otherwise} \end{cases}$$

- (a) Show that the value of  $k$  is 4 (4)  
 (b) Write down the value of  $E(X)$ . (1)  
 (c) Calculate  $\text{Var}(X)$ . (4)  
 (d) Find the probability that a randomly chosen customer's queuing time will differ from the mean by at least half a minute. (3)

$$\begin{aligned} \text{a) } \int f(x) dx &= 1 \Rightarrow \frac{3}{32} \int_0^k (kx - x^2) dx = \frac{3}{32} \left[ \frac{kx^2}{2} - \frac{x^3}{3} \right]_0^k \\ &= \frac{3}{32} \left( \frac{k^3}{6} \right) = 1 \Rightarrow k^3 = 64 \therefore k = 4 \end{aligned}$$

$$\begin{aligned} \text{b) } E(X) &= \int_0^4 x f(x) dx = \frac{3}{32} \int_0^4 (4x^2 - x^3) dx \\ &= \frac{3}{32} \left[ \frac{4}{3}x^3 - \frac{1}{4}x^4 \right]_0^4 = \frac{3}{32} \left( \frac{64}{3} \right) = 2. \end{aligned}$$

$$\begin{aligned} \text{c) } E(X^2) &= \int_0^4 x^2 f(x) dx = \frac{3}{32} \int_0^4 (4x^3 - x^4) dx \\ &= \frac{3}{32} \left[ x^4 - \frac{1}{5}x^5 \right]_0^4 = \frac{3}{32} \left( \frac{256}{5} \right) = \frac{24}{5} \end{aligned}$$

$$\text{Var}(X) = E(X^2) - E(X)^2 = \frac{24}{5} - 4 = \frac{4}{5}$$

$$\begin{aligned} \text{d) } P(1.5 < X < 2.5) &= \frac{3}{32} \int_{1.5}^{2.5} (4x - x^2) dx = \frac{3}{32} \left[ 2x^2 - \frac{x^3}{3} \right]_{1.5}^{2.5} = 0.633 \end{aligned}$$

Leave blank

6. A bag contains a large number of balls.

65% are numbered 1

35% are numbered 2

A random sample of 3 balls is taken from the bag.

Find the sampling distribution for the range of the numbers on the 3 selected.

$$\begin{aligned} \text{range} = 0 & \quad \begin{matrix} 1,1,1 \\ 2,2,2 \end{matrix} \quad \begin{matrix} P = 0.65^3 \\ P = 0.35^3 \end{matrix} \end{aligned}$$

$$\therefore P(\text{range} = 0) = 0.3175$$


$$\therefore$$

range	0	1
P	0.3175	0.6825

7. The continuous random variable  $X$  has probability density function  $f(x)$  given by

$$f(x) = \begin{cases} \frac{x^2}{45} & 0 \leq x \leq 3 \\ \frac{1}{5} & 3 < x < 4 \\ \frac{1}{3} - \frac{x}{30} & 4 \leq x \leq 10 \\ 0 & \text{otherwise} \end{cases}$$

- (a) Sketch  $f(x)$  for  $0 \leq x \leq 10$  (4)
- (b) Find the cumulative distribution function  $F(x)$  for all values of  $x$ . (8)
- (c) Find  $P(X \leq 8)$ . (2)



$$0 \leq x \leq 3 \quad F(x) = \int_0^x \frac{t^2}{45} dt = \left[ \frac{t^3}{135} \right]_0^x = \frac{x^3}{135}$$

$$3 < x < 4 \quad F(x) = \int_3^x \frac{1}{5} dt + F(3) = \left[ \frac{1}{5}t \right]_3^x + \frac{27}{135} = \frac{1}{5}x - \frac{2}{5}$$

$$4 \leq x \leq 10 \quad F(x) = \int_4^x \left( \frac{1}{3} - \frac{t}{30} \right) dt + F(4) = \left[ \frac{1}{3}t - \frac{t^2}{60} \right]_4^x + \frac{2}{5}$$

$$= -\frac{x^2}{60} + \frac{1}{3}x - \frac{2}{3}$$

$$F(x) = \begin{cases} 0 & x < 0 \\ \frac{x^3}{135} & 0 \leq x \leq 3 \\ \frac{1}{5}x - \frac{2}{5} & 3 < x < 4 \\ -\frac{x^2}{60} + \frac{1}{3}x - \frac{2}{3} & 4 \leq x \leq 10 \\ 1 & x > 10 \end{cases}$$

c)  $F(8) = \frac{-8^2}{60} + \frac{1}{3}(8) - \frac{2}{3} = \frac{14}{15}$

8. In a large restaurant an average of 3 out of every 5 customers ask for water with their meal. A random sample of 10 customers is selected.

- (a) Find the probability that
- exactly 6 ask for water with their meal,
  - less than 9 ask for water with their meal.
- (5)

A second random sample of 50 customers is selected.

- (b) Find the smallest value of  $n$  such that

$$P(X < n) \geq 0.9$$

where the random variable  $X$  represents the number of these customers who ask for water. (3)

a)  $X =$  Customer ask for water  $X \sim B(10, 0.6)$   
 $Y =$  Customer does not ask for water

$$Y \sim B(10, 0.4)$$

i)  $P(X=6) \Rightarrow P(Y=4) = \binom{10}{4} 0.4^4 0.6^6 = 0.251$

ii)  $P(X < 9) = P(X \leq 8) \Rightarrow P(Y \geq 2) = P(Y > 1)$   
 $= 1 - P(Y \leq 1) = 0.9536$

b)  $P(X < n) = P(Y > 50 - n) \geq 0.9 \quad X \sim B(50, 0.6)$   
 $\Rightarrow 1 - P(Y \leq 50 - n) \geq 0.9 \quad Y \sim B(50, 0.4)$   
 $\Rightarrow P(Y \leq 50 - n) \leq 0.1$

$$P(Y \leq 15) = 0.0955 < 0.10$$

$$P(Y \leq 16) = 0.1561 > 0.10$$

$$\therefore 50 - n = 15 \quad \therefore n = 35$$