

Mark Scheme (Results)

Summer 2013

GCE Statistics 4 (6686/01)

## Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at [www.edexcel.com](http://www.edexcel.com).

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

[www.edexcel.com/contactus](http://www.edexcel.com/contactus)

### **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2013

Publications Code UA037005

All the material in this publication is copyright

© Pearson Education Ltd 2013

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

# EDEXCEL GCE MATHEMATICS

## General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes:

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\checkmark$  will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper
  - $\square$  The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
  5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
  6. If a candidate makes more than one attempt at any question:
    - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
    - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
  7. Ignore wrong working or incorrect statements following a correct answer.
  8. In some instances, the mark distributions (e.g. M1, B1 and A1) printed on the candidate's response may differ from the final mark scheme

Question Number	Scheme	Marks
1.	$H_0 : \sigma^2 = 2.4^2 \quad H_1 : \sigma^2 \neq 2.4^2$ $s^2 = \frac{1414.08 - 10 \times \left(\frac{113.4}{10}\right)^2}{9} = 14.236$ $\chi^2 = \frac{9s^2}{\sigma^2} = \frac{9 \times 14.236}{2.4^2} = 22.24375$ <p>Critical Value <math>\chi_9^2(0.025) = 19.023</math></p> <p>Significant result, there is evidence of a change in standard deviation <u>or</u> the data do not support George's belief</p>	B1 M1 A1 M1 A1 B1 A1cso (7) [7]
<b>Notes</b>		
<p>1<sup>st</sup> B1 Both hypotheses, must use <math>\sigma</math>. Allow <math>H_0 : \sigma = 2.4 \quad H_1 : \sigma \neq 2.4</math></p> <p>1<sup>st</sup> M1 correct method used</p> <p>1<sup>st</sup> A1 awrt 14.2</p> <p>2<sup>nd</sup> M1 <math>\chi^2 = \frac{9 \times \text{"their } s^2 \text{"}}{2.4^2}</math></p> <p>2<sup>nd</sup> A1 awrt 22.2</p> <p>2<sup>nd</sup> B1 for critical value, this should be compatible with their alternative hypothesis ( 16.919 for one tail test)</p> <p>3<sup>rd</sup> A1ft fully correct solution only</p>		

Question Number	Scheme	Marks
<p>2. (a)</p>	<p><math>d = \text{Jan} - \text{June}: -2, 1, -3, 2, -2, 3, 2, 2</math>  <math>\bar{d} = 0.375, \sum d^2 = 39 \Rightarrow s^2 = 5.4107\dots</math> or <math>s = 2.326\dots</math>  <math>t_7(0.025) = 2.365</math>            Confidence Interval: <math>0.375 \pm 2.365 \times \frac{2.326\dots}{\sqrt{8}}</math>  <math>= \underline{(-1.57, 2.32)}</math> (o.e.)</p>	<p>M1 M1, M1 B1 M1 A1, A1 (7)</p>
	<b>Notes</b>	
<p>(b)</p>	<p><math>H_0 : \mu_D = 0</math>    <math>H_1 : \mu_D \neq 0</math>            Comment that 0 is in the interval            Not sig, no evidence of a change in <b>mean time to assemble component</b></p>	<p>B1 M1 A1ft (3) <b>[10]</b></p>
	<b>Notes</b>	
<p>(a)</p>	<p>1<sup>st</sup> M1 for attempting differences            2<sup>nd</sup> M1 for attempting <math>\bar{d}</math>            3<sup>rd</sup> M1 for attempting <math>s_d^2</math>, correct expression with their <math>\sum d^2</math> and <math>\bar{d}</math> or correct calculation (to 2 sf or better)            4<sup>th</sup> M1 for use of a correct CI formula, using a value for <math>t</math> and ft their values.            1<sup>st</sup> A1 for lower limit of -1.57 or -2.32            2<sup>nd</sup> A1 for corresponding upper limit</p>	
<p>S.C.</p>	<p>Allow A1A1 for (0, 2.32)</p>	
<p>(b)</p>	<p>B1 for both hypotheses using <math>\mu_D</math>            M1 for a comment about 0 being in (or out) of <u>their</u> interval            A1 contextual conclusion – must include assemble components</p>	
<p>S.C.</p>	<p>If they have used difference in means test in part (a) to get the confidence interval then award the B1 for <math>H_0 : \mu_x - \mu_y = 0</math>    <math>H_1 : \mu_x - \mu_y \neq 0</math> or the correct hypotheses.</p>	

Question Number	Scheme	Marks
<p><b>3. (a)</b></p> <p><math>H_0 : \sigma_A^2 = \sigma_B^2 \quad H_1 : \sigma_A^2 \neq \sigma_B^2</math></p> <p><math>F = \frac{s_B^2}{s_A^2} = \frac{4.37^2}{4.24^2} = 1.0622\dots</math></p> <p><math>F_{12,6}(0.01) = 7.72</math></p> <p>Not sig, so no evidence of a difference in variances</p> <p><b>(b)</b></p> <p><math>H_0 : \mu_A = \mu_B \quad H_1 : \mu_A &lt; \mu_B</math></p> <p><math>s_p^2 = \frac{6 \times 4.24^2 + 12 \times 4.37^2}{18} = 18.7238 \quad \text{or} \quad s_p = 4.327\dots</math></p> <p><math>t = \pm \frac{14.31 - 8.43}{s_p \sqrt{\frac{1}{7} + \frac{1}{13}}} = \pm 2.8985\dots</math> awrt 2.9</p> <p><math>t_{18}(0.01) = 2.552</math></p> <p>sig, there is evidence to support <b>archaeologist's</b> claim or there is evidence that bricks for site <i>B</i> have higher <b>mean compression</b> strength than those from site <i>A</i>.</p> <p><b>(c)</b></p> <p>The test in (b) requires <math>\sigma_A^2 = \sigma_B^2</math> and the test in part (a) shows that this is a reasonable assumption. (o.e.)</p>	<p>B1</p> <p>M1A1</p> <p>B1</p> <p>A1ft <b>(5)</b></p> <p>B1</p> <p>M1</p> <p>M1A1</p> <p>B1</p> <p>A1ft</p> <p><b>(6)</b></p> <p>B1</p> <p><b>(1)</b></p> <p><b>[12]</b></p>	
<b>Notes</b>		
<p><b>(a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p>	<p>M1 for use of a correct formula Allow <math>F = \frac{s_A^2}{s_B^2} = \frac{4.24^2}{4.37^2} = 0.941\dots</math> with 0.1295..</p> <p>B1 if <i>A</i> and <i>B</i> not used it must be clear which is <i>A</i> and which is <i>B</i></p> <p>1<sup>st</sup> M1 for attempt to calculate <math>s_p</math> or <math>s_p^2</math></p> <p>2<sup>nd</sup> M1 for attempt correct test statistic</p> <p>2<sup>nd</sup> A1 ft need <b>archaeologist's or compression</b></p> <p>Need to refer to 'allows us to assume variances the same' and this is needed in for test.</p> <p>oe</p>	

Question Number	Scheme	Marks
<p><b>4. (a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p> <p><b>(d)</b></p> <p><b>(e)</b></p> <p><b>(f)</b></p> <p><b>(g)</b></p>	<p><math>E(X) = \mu = \frac{2a-a}{2} = \frac{a}{2}; \quad E(\bar{X}) = \mu = \frac{a}{2}</math> so biased estimator for <math>a</math></p> <p>Bias = <math>\frac{a}{2} - a = -\frac{a}{2}</math></p> <p><math>k = 2</math></p> <p><math>\text{Var}(X) = \sigma^2 = \frac{(2a-a)^2}{12} = \frac{9a^2}{12} = \frac{3a^2}{4}; \quad \text{Var}(\bar{X}) = \frac{\sigma^2}{2}</math></p> <p><math>\text{Var}(Y) = k^2 \text{Var}(\bar{X}) = 4 \times \frac{\sigma^2}{2} = 4 \times \frac{3a^2}{4 \times 2} = \frac{3}{2}a^2</math></p> <p><math>E(M) = \int \frac{2x(x+a)}{9a^2} dx = \left[ \frac{2x^3}{27a^2} + \frac{ax^2}{9a^2} \right]_{-a}^{2a} = \left( \frac{16a}{27} + \frac{4a}{9} \right) - \left( -\frac{2a}{27} + \frac{a}{9} \right) [=a]</math></p> <p>So <math>E(M) = a</math> and therefore <math>M</math> is an unbiased estimator for <math>a</math></p> <p><math>\text{Var}(M) = \frac{3}{2}a^2 - a^2 = \frac{1}{2}a^2</math></p> <p><math>\text{Var}(M) &lt; \text{Var}(Y)</math>, so <math>M</math> is the better estimator of <math>a</math></p> <p>Maximum value = <u>5</u></p>	<p>M1;A1</p> <p>B1(accept <math>\pm</math>)</p> <p><b>(3)</b></p> <p>B1</p> <p><b>(1)</b></p> <p>B1;B1</p> <p>M1,A1</p> <p><b>(4)</b></p> <p>M1A1,M1d</p> <p>A1cso</p> <p><b>(4)</b></p> <p>B1</p> <p><b>(1)</b></p> <p>M1, A1</p> <p><b>(2)</b></p> <p>B1ft</p> <p><b>(1)</b></p> <p><b>[16]</b></p>
<b>Notes</b>		
<p><b>(a)</b></p> <p><b>(c)</b></p> <p><b>(d)</b></p> <p><b>(f)</b></p> <p><b>(g)</b></p>	<p>M1 for use of formula or integration or symmetry to find <math>E(X)</math></p> <p>1<sup>st</sup> B1 for use of formula for variance</p> <p>2<sup>nd</sup> B1 for use of <math>\frac{\sigma^2}{n}</math> formula</p> <p>M1 for <math>k^2 \text{Var}(\bar{X})</math> and ft their <math>k</math></p> <p>1<sup>st</sup> M1 for attempt at correct integration of correct expression</p> <p>1<sup>st</sup> A1 for correct integration</p> <p>2<sup>nd</sup> M1d dependent on previous M, for attempting to use correct limits</p> <p>2<sup>nd</sup> A1 need statement that <math>M</math> is therefore unbiased</p> <p>M1 for comparison of their <math>\text{Var}(Y)</math> and their <math>\text{Var}(M)</math></p> <p>B1ft for calculation of their estimate based on their choice in (f). If they choose <math>Y</math> answer is 4 (or twice their <math>k</math>)</p>	



Question Number	Scheme	Marks
<p>5. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p><math>Y = \text{no. of organisms in 20 ml. } Y \sim \text{Po}(2\lambda)</math>  <math>\text{Size} = P(Y \geq 4   Y \sim \text{Po}(2)) = 1 - P(Y \leq 3) = 1 - 0.8571 = \underline{\underline{0.1429}}</math></p> <p><math>P(\text{Type II error}) = 1 - P(Y \geq 4   Y \sim \text{Po}(5)) = P(Y \leq 3) = \underline{\underline{0.2650}}</math></p> <p><math>X = \text{no. of organisms in 10 ml. } X \sim \text{Po}(\lambda)</math>  <math>\text{Power} = P(X \geq 2) + P(X=1) \times P(X \geq 2)</math>  <math>= P(X \geq 2) [1 + P(X=1)] = [1 - e^{-\lambda} (1 + \lambda)] \times [1 + \lambda e^{-\lambda}]</math>  <math>= 1 - e^{-\lambda} - \lambda e^{-\lambda} + \lambda e^{-\lambda} - \lambda(1 + \lambda)e^{-2\lambda} = 1 - e^{-\lambda} - \lambda(1 + \lambda)e^{-2\lambda}</math></p> <p><math>r = 0.92</math></p> <p>See Graph paper</p> <div data-bbox="284 891 1236 1818" data-label="Figure"> </div>	<p>M1, A1 (2)</p> <p>M1, A1 (2)</p> <p>M1 M1A1 A1cso (4)</p> <p>B1 (1)</p> <p>B1B1 (2)</p>
<p>Question Number</p>	<p>Scheme</p>	<p>Marks</p>
<p>(f)</p>	<p>Expected time for statistician's test: <math>30 \times P(X=1) + 15 \times [1 - P(X=1)]</math></p>	<p>M1</p>

	$= 30\lambda e^{-\lambda} + 15(1 - \lambda e^{-\lambda}) = 15(1 + \lambda e^{-\lambda})$ slower if: $15(1 + \lambda e^{-\lambda}) > 20, \Rightarrow \lambda e^{-\lambda} > \frac{1}{3}$  $\lambda e^{-\lambda}$ with $\lambda = 1$ is 0.36..., with $\lambda = 2$ is 0.27...so second(statisticians) test is slower if $\lambda = 1$ but faster for $\lambda = 2$ . Second test is more powerful for all $\lambda$ Choose second test - more powerful and faster for $\lambda \geq 2$	A1 M1,A1cso (4)  B1  B1 (2) [17]
<b>Notes</b>		
(a)	M1 for correct expression for size using Po(2)	
(b)	M1 for correct expression using Po(5)	
(c)	1 <sup>st</sup> M1 for a correct expression in terms of probabilities Alternate answer $1 - [P(X=0) + P(X=1) \times P(X \leq 1)]$ 2 <sup>nd</sup> M1 for an attempt at a correct equation in $\lambda$ 1 <sup>st</sup> A1 for a correct expression in $\lambda$	
(e)	1 <sup>st</sup> B1 points 2 <sup>nd</sup> B1 curve (or straight lines)	
(f)	1 <sup>st</sup> M1 for an attempt to calculate expected time Alternate method $15 + 15 \times P(X=1)$ 1 <sup>st</sup> A1 for a correct expression in terms of $\lambda$ 2 <sup>nd</sup> M1 for attempt at correct inequality	
(g)	1 <sup>st</sup> B1 for a comment about power & timings 2 <sup>nd</sup> B1 for selecting second test	

Question Number	Scheme	Marks
6. (a)	$H_0 : \sigma_A^2 = \sigma_B^2 \quad H_1 : \sigma_A^2 \neq \sigma_B^2$ $s_A^2 = (0.25)^2 = 0.0625 \quad s_B^2 = (0.178885\dots)^2 = 0.032$	B1 B1B1

$$F = \frac{0.0625}{0.032} = 1.953\dots$$

Critical Value:  $F_{3,5} = 5.41$

not sig, samples come from populations with common variance

(b) 
$$s_p^2 = \frac{3 \times 0.25^2 + 5 \times 0.032}{8} = 0.04343\dots = (0.0284\dots)^2$$

Use  $\frac{8s_p^2}{\sigma^2} \sim \chi_8^2$

$$1.344 < \frac{8 \times 0.0434\dots}{\sigma^2} < 21.955$$

99% confidence interval is **(0.0158, 0.259)**

M1A1

B1

A1cso (7)

M1A1

M1

B1,B1

A1 (6)

[13]

Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)

Order Code UA037005 Summer 2013

For more information on Edexcel qualifications, please visit our website  
[www.edexcel.com](http://www.edexcel.com)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual



Llywodraeth Cymru  
Welsh Assembly Government

