

Mark Scheme (Results)

Summer 2014

Pearson Edexcel GCE in Statistics S4
(6686/01)

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2014

Publications Code UA040129

All the material in this publication is copyright

© Pearson Education Ltd 2014

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

General Instructions for Marking

1. The total number of marks for the paper is 75
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \surd will be used for correct ft
 - cao – correct answer only
 - cso – correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - d... or dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper or ag- answer given
 - \square or d... The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

Question Number	Scheme	Mark
1.	$H_0 : \mu = 100 \quad H_1 : \mu < 100$ $t = \frac{ \bar{x} - \mu }{s/\sqrt{n}} = \frac{ 92.875 - 100 }{8.3055/\sqrt{8}} = 2.4264... \quad \text{or} \quad \frac{c - 100}{8.3055/\sqrt{8}} = -1.895 \therefore \text{CR } c < 94.435$ $t_7(5\%) = \pm 1.895$ <p>There is evidence to reject H_0. <u>Malcolm's belief is supported</u> or there is evidence that the amount of <u>oil</u> placed in bottles is <u>less than 100mm</u></p>	B1 M1A1 B1 A1ft (5)
Notes		
	B1 both hypotheses M1 either $\frac{ 92.875 - 100 }{8.3055/\sqrt{8}}$ or $p = 0.0228$ or $\frac{c - 100}{8.3055/\sqrt{8}} = -(a \text{ } t \text{ value})$ A1 awrt 2.43 or awrt 94.4 or awrt 0.0228 B1 ± 1.895 or $0.0228 < 0.05$ (must have correct comparison for hypotheses) A1ft Do Not allow contradictions	


Question Number	Scheme	Marks
2(a)(i)	Type I - H_0 rejected when it is true	B1
(ii)	Type II - H_0 is accepted when it is false	B1 (2)
(b)	$P(X < c \mid \lambda = 6) \approx 0.05$ $P(X \leq 2) = 0.0620$ $P(X \leq 1) = 0.0174$ Critical region = $X \leq 2$ $P(\text{Type 1 error}) = P(X \leq 2 \mid \lambda = 6) = 0.062$	M1 A1 A1cao (3)
(c)	$P(\text{Type 2 error}) = P(X \geq 3 \mid \lambda = 4)$ $= 1 - 0.2381$ $= 0.7619$	M1 A1 (2)
Notes		
(b)	M1 use of Po(6) A1 correct CR. May be implied by correct probability. Allow if written as part of a 2 tailed CR A1 awrt 0.062	
(c)	M1 using Po(4) and $1 - P(X \leq 2)$, ft their CR in (b) if one tail A1awrt 0.762	

Question Number	Scheme	Marks
<p>3(a) (i)</p> <p>(ii)</p> <p>(b)</p>	<p>$\bar{x} = \frac{181}{9} = 20.111 \dots$</p> <p>$s_x^2 = \left(\frac{3913 - 9 \times \bar{x}^2}{8} \right) = 34.1111 \quad (s_x = 5.84)$</p> <p>$t_8(0.025) \text{ cv} = 2.306$</p> <p>95% CI for μ is $= 20.111 \pm 2.306 \times \frac{5.84}{\sqrt{9}}$</p> <p>$= (15.6, 24.6)$ awrt (15.6, 24.6)</p> <p>$\chi_8^2(0.025) = 2.18(0), \quad \chi_8^2(0.975) = 17.535$</p> <p>95% CI for σ^2 is given by $2.180 < \frac{8s_x^2}{\sigma^2} < 17.535$</p> <p>So 95% CI for σ^2 is $=$ awrt (15.6, 125)</p> <p>Require $P(X < 16) = P\left(Z < \frac{16 - \mu}{\sigma}\right)$ to be as small as possible OR</p> <p>$\frac{16 - \mu}{\sigma}$ to be as large as possible but negative; imply lowest σ and largest μ.</p> <p>$P\left(Z < \frac{16 - 24.6}{\sqrt{15.6}}\right); = 1 - 0.9854 =$ 0.0146 or 0.0147</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1, A1</p> <p>B1B1</p> <p>M1</p> <p>A1</p> <p>(10)</p> <p>M1</p> <p>M1A1ft;A1</p> <p>(4)</p>
Notes		
<p>(a)(i)</p> <p>(ii)</p> <p>(b)</p>	<p>1st M1 ‘ their \bar{x} ’ $\pm t \text{ value} \times \frac{\text{‘their } s\text{’}}{\sqrt{9}}$</p> <p>1st A1 awrt 15.6</p> <p>2nd A1 awrt 24.6</p> <p>2nd M1 $\chi^2 < \frac{8s^2}{\sigma^2} < \chi^2$</p> <p>A1 awrt 15.6 and 125</p> <p>M1 Identify must use lowest σ and largest μ</p> <p>M1 standardising and finding correct area use either limit for μ and σ</p> <p>A1 ft their lowest σ and largest μ</p> <p>A1 awrt 0.0146 or 0.0147</p>	

Question Number	Scheme	Marks
4(a)	The <u>differences</u> in the mean number of hours sleep are <u>normally</u> distributed	B1 (1)
(b)	Differences are 0.8, 0.7, -0.3, 1.2, 0.7, 2.9, 1.3, 0.8 $\bar{d} = \frac{8.1}{8} = 1.0125$ $s_d = \sqrt{\frac{13.89 - 8 \times 1.0125^2}{7}} = 0.901\dots$ both \bar{d} and s $H_0: \mu_D = 1/6 \quad H_1: \mu_D > 1/6$ $t = \frac{1.0125 - 1/6}{0.901/\sqrt{8}} = \text{awrt } 2.65$ or $\frac{c - 1/6}{0.901/\sqrt{8}} = 2.988 \therefore \text{CR } c > \text{awrt } 1.12$ $t_7(1\%) = 2.998$ (or prob. = awrt 0.0164) There is insufficient evidence to suggest the <u>drug increases</u> the mean number of hours slept by <u>more than 10</u> minutes.	M1 M1 M1 B1 M1A1 B1 A1ft (8)

Notes

(a)	B1 for a comment that mentions “differences” and “normal” distribution
(b)	1 st M1 for attempting the d s 2 nd M1 for attempting \bar{d} 1 st M1 for s_d or s_d^2 1 st B1 for both hypotheses correct in terms of μ or μ_d . (allow a defined symbol) Do not allow 10 instead of 1/6 (awrt 0.167) unless working in minutes throughout 3 rd M1 for attempting the correct test statistic $\frac{\bar{d} - 1/6}{s_d/\sqrt{8}}$ or $p = \text{awrt } 0.016$ or $\frac{c - 1/6}{0.901/\sqrt{8}} = t$ value 2 nd A1 awrt 2.65 / 2.655 or awrt 1.12 or awrt 0.016 2 nd B1 2.988 or 0.0164 3 rd A1ft for a correct comment in context based on their test statistic and their cv. Do not allow contradictions.

Question Number	Scheme	Marks
<p>5 (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>$X \sim B(10, 0.5)$ Size = $P(\text{reject } H_0 p = 0.5)$ = $P(X < 3 p = 0.5)$ = 0.0547</p> <p>Power = $P(X = 2) + P(X = 1) + P(X = 0)$ = $45p^2(1 - p)^8 + 10p(1 - p)^9 + (1 - p)^{10}$ = $(1 - p)^8(45p^2 + 10p(1 - p) + (1 - p)^2)$ = $(1 - p)^8(36p^2 + 8p + 1)$</p> <p>$r = 0.68$ $s = 0.17$</p>  <p>$P(\text{Type II error}) \leq 0.4$ $1 - \text{power} \leq 0.4$ Power ≥ 0.6 $p < 0.23$</p>	<p>B1 (1)</p> <p>M1 A1 A1cso (3)</p> <p>B1 B1 (2)</p> <p>B1 points B1 curve (2)</p> <p>M1 A1 A1 (3)</p>
Notes		
<p>(b)</p> <p>(c)</p> <p>(e)</p>	<p>M1 for a correct expression/selection of probabilities A1 for a fully correct expression SC B1 B0 both correct but not given to 2 dp M1 may be implied by Power ≥ 0.6 or correct value or by correct answer A1 may be implied by correct answer A1 allow number between 0.22 and 0.23 inclusive and either $<$ or \leq</p>	

Question Number	Scheme	Marks
6(a)	It is the probability distribution of T .	B1 (1)
(b)	An estimator is biased if $E(T) \neq \theta$	B1 (1)
(c)	$E(\hat{\mu}_1) = \frac{E(X_3)+E(X_5)+E(X_7)}{3} = \frac{\mu+\mu+\mu}{3} = \mu \therefore \text{Bias} = 0$ $E(\hat{\mu}_2) = \frac{5E(X_1)+2E(X_2)+E(X_9)}{6} = \frac{5\mu+2\mu+\mu}{6} = \frac{4\mu}{3} \therefore \text{Bias} = \frac{\mu}{3}$ $E(\hat{\mu}_3) = \frac{3E(X_{10}) - E(X_1)}{3} = \frac{3\mu - \mu}{3} = \frac{2\mu}{3} \therefore \text{Bias} = -\frac{\mu}{3}$	M1A1 A1 A1 (4)
(d)	$\text{Var}(\hat{\mu}_1) = \frac{1}{9}(\text{Var}(X_3) + \text{Var}(X_5) + \text{Var}(X_7))$ $= \frac{1}{9}(\sigma^2 + \sigma^2 + \sigma^2)$ $= \frac{\sigma^2}{3}$ $\text{Var}(\hat{\mu}_2) = \frac{1}{36}(25\text{Var}(X_1) + 4\text{Var}(X_2) + \text{Var}(X_9))$ $= \frac{1}{36}(25\sigma^2 + 4\sigma^2 + \sigma^2)$ $= \frac{5}{6}\sigma^2$ $\text{Var}(\hat{\mu}_3) = \frac{1}{9}(9\text{Var}(X_{10}) + \text{Var}(X_1))$ $= \frac{1}{9}(9\sigma^2 + \sigma^2)$ $= \frac{10\sigma^2}{9}$	M1 A1 M1 A1 M1 A1 (6)
(e)(i)	$\hat{\mu}_1$ is the best estimator. It has no bias	B1
(ii)	It has <u>same magnitude of bias</u> as $\hat{\mu}_2$ but it has the <u>largest variance</u> $\hat{\mu}_3$ is the worst estimator.	B1ft B1dcao (3)
Notes		
(c)	M1 finding $E(\hat{\mu})$ A1 bias 0 A1 $\pm \frac{\mu}{3}$ A1 $\pm \frac{\mu}{3}$	
(d)	For method marks allow an incorrect variance, M1 squaring 9, M1 Squaring 5 and 2, M1 adding variances. Do not penalise same mistake twice.	
(e)(ii)	Must have idea that its bias is the same as another ($\hat{\mu}_2$) and state it has largest variance for first B1 . ft their values of Var. Second B1 dependent on first B1cao SC $\hat{\mu}_3$ because <u>largest variance</u> B1 B0	

Question Number	Scheme	Marks
7(a)	<p>The variance of the two group's marks must be the same.</p> $H_0 : \sigma_1^2 = \sigma_2^2 \quad H_1 : \sigma_1^2 \neq \sigma_2^2$ $s_1^2 = 16.25$ $\left(F_{8,6} = \right) \frac{16.25}{12.9} = (1.2597...) \quad \left(\frac{1}{F_{8,6}} = \frac{12.9}{16.25} = 0.7938... \right)$ <p>$F_{8,6}$ 5% (two-tail) cv = 4.15 (0.241) (or prob. = awrt 0.39)</p> <p>Not significant so can accept the assumption that variances are equal.</p>	<p>B1 B1 B1 M1A1 B1 A1 (7)</p>
(b)	$H_0 : \mu_1 = \mu_2 \quad H_1 : \mu_1 \neq \mu_2$ $s_p^2 = \frac{8 \times 16.25 + 6 \times 12.9}{14} = 14.814... \quad \text{or} \quad s_p = 3.8489...$ $(t_{14} =) (\pm) \frac{30.33 - 31.29}{s_p \sqrt{\frac{1}{9} + \frac{1}{7}}} = (\pm) 0.494927.. \quad = \text{awrt } \underline{\underline{0.49}}$ <p>$t_{14}(0.025)$ two-tail cv = 2.145</p> <p>There is insufficient evidence to reject H_0. There is no evidence of a significant difference between the mean marks of the two groups</p>	<p>B1 M1 B1 M1A1 B1 A1 (7)</p>
Notes		
(a)	<p>2nd B1allow σ or σ^2 3rd B1 allow awrt 16.3 or $s_1 =$ awrt 4.03 M1 for use of the correct test statistic 5th B1 allow "assumption is correct"</p>	
(b)	<p>1st M1 for attempting s_p or s_p^2 1st B1 for 30.33 2nd M1 for use of a correct test statistic 2nd A1 for awrt 0.49 (accept \pm) or 0.495 2nd B1 for 2.145 (allow \pm 1.761 for one-tailed H_1)</p>	

