

Please check the examination details below before entering your candidate information

Candidate surname	Other names
-------------------	-------------

**Pearson Edexcel  
International  
Advanced Level**

Centre Number	Candidate Number
<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>

**Wednesday 23 October 2019**

Morning (Time: 1 hour 30 minutes)	Paper Reference <b>WME01/01</b>
-----------------------------------	---------------------------------

**Mathematics**  
**International Advanced Subsidiary/Advanced Level**  
**Mechanics M1**

<b>You must have:</b> Mathematical Formulae and Statistical Tables (Blue), calculator	Total Marks
--	-------------

**Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

**Instructions**

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of  $g$  is required, take  $g = 9.8 \text{ m s}^{-2}$ , and give your answer to either 2 significant figures or 3 significant figures.

**Information**

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets – *use this as a guide as to how much time to spend on each question.*

**Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ►

P58496A





















4.




Figure 1

A particle,  $P$ , of mass  $km$  lies on a fixed rough plane. The plane is inclined to the horizontal at an acute angle  $\theta$ . A horizontal force of magnitude  $X$  acts on  $P$ , as shown in Figure 1. The line of action of the force lies in the vertical plane which contains the line of greatest slope of the inclined plane that passes through  $P$ . The coefficient of friction between  $P$  and the inclined plane is  $\mu$ .

When  $X = mg$ , the particle  $P$  is in equilibrium and on the point of sliding down the plane.

(a) Show that  $\mu = \frac{k \tan \theta - 1}{k + \tan \theta}$  (10)

(b) Deduce that, when  $k = 1$ ,  $\theta$  must be greater than  $45^\circ$  (2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q4

(Total 12 marks)

5.




Figure 2

A non-uniform beam,  $AB$ , has length  $5\text{ m}$  and mass  $12\text{ kg}$ . The beam is suspended in a horizontal position by two vertical ropes. One rope is attached to the beam at  $A$ . The other rope is attached to the beam at  $C$ , where  $CB = 1.5\text{ m}$ , as shown in Figure 2. The distance of the centre of mass of the beam from  $A$  is  $1.75\text{ m}$ . The beam is modelled as a non-uniform rod and the ropes are modelled as light inextensible strings.

A particle of mass  $M\text{ kg}$  is now placed on the beam at  $B$  and the beam remains in equilibrium in a horizontal position.

- (a) Find the largest possible value of  $M$ . (3)

The particle at  $B$  is now removed and a particle of mass  $15\text{ kg}$  is now placed on the beam at the point  $D$ , where  $AD = x$  metres. The beam remains in equilibrium in a horizontal position.

Given that the tension in the rope attached to the beam at  $C$  is now twice the tension in the rope attached to the beam at  $A$ ,

- (b) find the value of  $x$ . (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA









Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q6

(Total 12 marks)

7. Two forces, **F** and **G**, act on a particle. The force **F** has magnitude 4N and acts in a direction with a bearing of  $120^\circ$  and the force **G** has magnitude 6N and acts due north.

Given that  $\mathbf{P} = 2\mathbf{F} + \mathbf{G}$ , find

(i) the magnitude of **P**

(ii) the direction of **P**, giving your answer as a bearing to the nearest degree.

(7)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA  
DO NOT WRITE IN THIS AREA  
DO NOT WRITE IN THIS AREA

**Question 7 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**(Total 7 marks)**

**Q7**



8. [In this question, the horizontal unit vectors  $\mathbf{i}$  and  $\mathbf{j}$  are directed due east and due north respectively and position vectors are given relative to a fixed origin  $O$ .]

Two speedboats,  $A$  and  $B$ , are each moving with constant velocity. The velocity of  $A$  is  $20\text{ km h}^{-1}$  due west and the velocity of  $B$  is  $40\text{ km h}^{-1}$  on a bearing of  $150^\circ$ . The boats are modelled as particles.

At noon, the position vector of  $A$  is  $60\mathbf{i}$  km and  $B$  is at the origin  $O$ . At time  $t$  hours after noon, the position vector of  $A$  is  $\mathbf{r}$  km and the position vector of  $B$  is  $\mathbf{s}$  km.

(a) Find the velocity of  $B$  in the form  $(p\mathbf{i} + q\mathbf{j})\text{ km h}^{-1}$  (3)

(b) Find expressions for  $\mathbf{r}$  and  $\mathbf{s}$  in terms of  $t$ ,  $\mathbf{i}$  and  $\mathbf{j}$ . (3)

(c) Find the time, to the nearest minute, at which the distance between the boats is the same as it was at noon. (8)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

