

WST01/01: Statistics S1

Question Number	Scheme	Marks
<p>Q1 (a)</p> <p>(b)</p> <p>(c)</p>	<p>$r = \frac{8825}{\sqrt{1022500 \times 130.9}}$, = awrt 0.763</p> <p>Teams with high attendance scored more goals (oe, statement in context)</p> <p>0.76(3)</p>	<p>M1 A1 (2)</p> <p>B1 (1)</p> <p>B1ft (1)</p> <p>Total 4</p>
	<p>(a) M1 for a correct expression, square root required Correct answer award 2/2</p> <p>(b) Context required (attendance and goals). Condone causality. B0 for 'strong positive correlation between attendance and goals' on its own oe</p> <p>(c) Value required. Must be a correlation coefficient between -1 and +1 inclusive. B1ft for 0.76 or better or same answer as their value from part (a) to at least 2 d.p.</p>	

WST01/01: Statistics S1

Question Number	Scheme	Marks
Q2 (a)	<p style="text-align: center;">P(R) and P(B) 2nd set of probabilities</p>	B1 B1 (2)
(b)	$P(H) = \frac{5}{12} \times \frac{2}{3} + \frac{7}{12} \times \frac{1}{2} = \frac{41}{72}$ or awrt 0.569	M1 A1 (2)
(c)	$P(R H) = \frac{\frac{5}{12} \times \frac{2}{3}}{\frac{41}{72}} = \frac{20}{41}$ or awrt 0.488	M1 A1ft A1 (3)
(d)	$\left(\frac{5}{12}\right)^2 + \left(\frac{7}{12}\right)^2$ $= \frac{25}{144} + \frac{49}{144} = \frac{74}{144}$ or $\frac{37}{72}$ or awrt 0.514	M1 A1ft A1 (3)
	Total 10	
(a)	<p>1st B1 for the probabilities on the first 2 branches. Accept 4/6 and 0.5 3/6 8/10</p> <p>2nd B1 for probabilities on the second set of branches. Accept 0.6, 0.3, 0.5 and 1.5/3</p> <p>Allow exact decimal equivalents using clear recurring notation if required.</p>	
(b)	M1 for an expression for P(H) that follows through their sum of two products of probabilities from their tree diagram	
(c)	M1 for $\frac{P(R \cap H)}{P(H)}$ with denominator their (b) substituted e.g. $\frac{P(R \cap H)}{P(H)} = \frac{5}{12}$ award M1.	
Formula seen		
Formula not seen	M1 for $\frac{\text{probability} \times \text{probability}}{\text{their } b}$ but M0 if fraction repeated e.g. $\frac{5}{12} \times \frac{2}{3}$.	
	1 st A1ft for a fully correct expression or correct follow through	
	2 nd A1 for $\frac{20}{41}$ o.e.	
(d)	M1 for $\left(\frac{5}{12}\right)^2$ or $\left(\frac{7}{12}\right)^2$ can follow through their equivalent values from tree diagram	
	1 st A1 for both values correct or follow through from their original tree and +	
	2 nd A1 for a correct answer	
	Special Case $\frac{5}{12} \times \frac{4}{11}$ or $\frac{7}{12} \times \frac{6}{11}$ seen award M1A0A0	

WST01/01: Statistics S1

Question Number	Scheme	Marks
Q3	(a) $2a + \frac{2}{5} + \frac{1}{10} = 1$ (or equivalent)	M1
	$a = \frac{1}{4}$ or 0.25	A1
		(2)
	(b) $E(X) = \underline{1}$	B1
		(1)
	(c) $E(X^2) = 1 \times \frac{1}{5} + 1 \times \frac{1}{10} + 4 \times \frac{1}{4} + 9 \times \frac{1}{5}$ (= 3.1)	M1
$\text{Var}(X) = 3.1 - 1^2,$	$= \underline{2.1}$ or $\frac{21}{10}$ oe	M1 A1
		(3)
(d) $\text{Var}(Y) = (-2)^2 \text{Var}(X),$	$= \underline{8.4}$ or $\frac{42}{5}$ oe	M1 A1
		(2)
(e) $X \geq Y$ when $X = 3$ or $2,$ so probability = " $\frac{1}{4}$ " + $\frac{1}{5}$	$= \underline{\frac{9}{20}}$ oe	M1 A1ft
		A1
		(3)
Total 11		
(a)	M1 for a clear attempt to use $\sum P(X = x) = 1$ Correct answer only 2/2. NB Division by 5 in parts (b), (c) and (d) seen scores 0. Do not apply ISW.	
(b)	B1 for 1	
(c)	1 st M1 for attempting $\sum x^2 P(X = x)$ at least two terms correct. Can follow through. 2 nd M1 for attempting $E(X^2) - [E(X)]^2$ or allow subtracting 1 from their attempt at $E(X^2)$ provided no incorrect formula seen. Correct answer only 3/3.	
(d)	M1 for $(-2)^2 \text{Var}(X)$ or $4\text{Var}(X)$ Condone missing brackets provided final answer correct for their $\text{Var}(X)$. Correct answer only 2/2.	
(e)	Allow M1 for distribution of Y and correct attempt at $-E(Y^2) - [E(Y)]^2$ M1 for identifying $X = 2, 3$ 1 st A1ft for attempting to find their $P(X=2) + P(X=3)$ 2 nd A1 for $\frac{9}{20}$ or 0.45	


WST01/01: Statistics S1

Question Number	Scheme	Marks
Q4 (a)	$\frac{2+3}{\text{their total}} = \frac{5}{\text{their total}} = \frac{1}{6}$ (** given answer**)	M1 A1cso (2)
(b)	$\frac{4+2+5+3}{\text{total}}, = \frac{14}{30}$ or $\frac{7}{15}$ or $0.4\bar{6}$	M1 A1 (2)
(c)	$P(A \cap C) = 0$	B1 (1)
(d)	$P(C \text{reads at least one magazine}) = \frac{6+3}{20} = \frac{9}{20}$	M1 A1 (2)
(e)	$P(B) = \frac{10}{30} = \frac{1}{3}, P(C) = \frac{9}{30} = \frac{3}{10}, P(B \cap C) = \frac{3}{30} = \frac{1}{10}$ or $P(B C) = \frac{3}{9}$	M1
	$P(B) \times P(C) = \frac{1}{3} \times \frac{3}{10} = \frac{1}{10} = P(B \cap C)$ or $P(B C) = \frac{3}{9} = \frac{1}{3} = P(B)$	M1
	So yes they are statistically independent	A1cso (3)
		Total 10
(a)	M1 for $\frac{2+3}{\text{their total}}$ or $\frac{5}{30}$	
(b)	M1 for adding at least 3 of "4, 2, 5, 3" and dividing by their total to give a probability Can be written as separate fractions substituted into the completely correct Addition Rule	
(c)	B1 for 0 or 0/30	
(d)	M1 for a denominator of 20 or $\frac{20}{30}$ leading to an answer with denominator of 20 $\frac{9}{20}$ only, 2/2	
(e)	1 st M1 for attempting all the required probabilities for a suitable test 2 nd M1 for use of a correct test - must have attempted all the correct probabilities. Equality can be implied in line 2. A1 for fully correct test carried out with a comment	

WST01/01: Statistics S1

Question Number	Scheme	Marks
Q5	(a) 23, 35.5 (may be in the table)	B1 B1 (2)
	(b) Width of 10 units is 4 cm so width of 5 units is 2 cm Height = $2.6 \times 4 = \mathbf{10.4 \text{ cm}}$	B1 M1 A1 (3)
	(c) $\sum fx = 1316.5$ $\bar{x} = \frac{1316.5}{56} =$ awrt 23.5 $\sum fx^2 = 37378.25$ can be implied So $\sigma = \sqrt{\frac{37378.25}{56} - \bar{x}^2} =$ awrt 10.7 allow $s = 10.8$	M1 A1 B1 M1 A1 (5)
	(d) $Q_2 = (20.5) + \frac{(28-21)}{11} \times 5 = 23.68\dots$ awrt 23.7 or 23.9	M1 A1 (2)
	(e) $Q_3 - Q_2 = 5.6$, $Q_2 - Q_1 = 7.9$ (or $\bar{x} < Q_2$) [7.9 > 5.6 so] negative skew	M1 A1 (2)
Total 14		
(b)	M1 for their width x their height=20.8. Without labels assume width first, height second and award marks accordingly.	
(c)	1 st M1 for reasonable attempt at $\sum x$ and /56 2 nd M1 for a method for σ or s , $\sqrt{\quad}$ is required Typical errors $\sum (fx)^2 = 354806.3$ M0, $\sum f^2 x = 13922.5$ M0 and $\sum (x)^2 f = 1733172$ M0 Correct answers only, award full marks.	
(d)	Use of $\sum f(x - \bar{x})^2 =$ awrt 6428.75 for B1 lcb can be 20, 20.5 or 21, width can be 4 or 5 and the fraction part of the formula correct for M1 - Allow 28.5 in fraction that gives awrt 23.9 for M1A1	
(e)	M1 for attempting a test for skewness using quartiles or mean and median. Provided median greater than 22.55 and less than 29.3 award for M1 for $Q_2 \quad Q_3 \quad - \quad < \quad -$ without as a valid reason. SC Accept mean close to median and no skew oe for M1A1	

WST01/01: Statistics S1

Question Number	Scheme	Marks
<p>Q6 (a)</p>		<p>B1 B1</p> <p>(2)</p>
(b)	<p>The points lie reasonably close to a straight line (o.e.)</p>	<p>B1</p> <p>(1)</p>
(c)	<p>$\sum d = 27.7, \quad \sum f = 146$ (both, may be implied)</p> <p>$S_{dd} = 152.09 - \frac{(27.7)^2}{6} = 24.208\dots$ awrt <u>24.2</u></p> <p>$S_{fd} = 723.1 - \frac{27.7 \times 146}{6} = 49.06\dots$ awrt <u>49.1</u></p>	<p>B1</p> <p>M1 A1</p> <p>A1</p> <p>(4)</p>
(d)	<p>$b = \frac{S_{fd}}{S_{dd}} = 2.026\dots$ awrt <u>2.03</u></p> <p>$a = \frac{146}{6} - b \times \frac{27.7}{6} = 14.97\dots$ so <u>$f = 15.0 + 2.03d$</u></p>	<p>M1 A1</p> <p>M1 A1</p> <p>(4)</p>
(e)	<p>A flight costs £2.03 (or about £2) for every extra 100km or about 2p per km.</p>	<p>B1ft</p> <p>(1)</p>
(f)	<p>$15.0 + 2.03d < 5d$ so $d > \frac{15.0}{(5 - 2.03)} = 5.00 \sim 5.05$</p> <p>So $t > 500 \sim 505$</p>	<p>M1</p> <p>A1</p> <p>(2)</p> <p>Total 14</p>

WST01/01: Statistics S1

<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>(f)</p>	<p>1st B1 for at least 4 points correct (allow \pm one 2mm square) 2nd B1 for all points correct (allow \pm one 2 mm square)</p> <p>Ignore extra points and lines Require reference to points and line for B1.</p> <p>M1 for a correct method seen for either - a correct expression 1st A1 for S_{dd} awrt 24.2 2nd A1 for S_{fd} awrt 49.1</p> <p>1st M1 for a correct expression for b - can follow through their answers from (c) 2nd M1 for a correct method to find a - follow through their b and their means 2nd A1 for $f=...$ in terms of d and all values awrt given expressions. Accept 15 as rounding from correct answer only.</p> <p>Context of cost and distance required. Follow through their value of b</p> <p>M1 for an attempt to find the intersection of the 2 lines. Value of t in range 500 to 505 seen award M1. Value of d in range 5 to 5.05 award M1. Accept t greater than 500 to 505 inclusive to include graphical solution for M 1A1</p>
---	---

WST01/01: Statistics S1

Question Number	Scheme	Marks
Q7 (a)	$P(D > 20) = P\left(Z > \frac{20-30}{8}\right)$ $= P(Z > -1.25)$ $= \underline{\underline{0.8944}} \qquad \text{awrt } \underline{\underline{0.894}}$	M1 A1 A1 (3)
(b)	$P(D < Q_3) = 0.75 \text{ so } \frac{Q_3 - 30}{8} = 0.67$ $Q_3 = \text{awrt } \underline{\underline{35.4}}$	M1 B1 A1 (3)
(c)	$35.4 - 30 = 5.4 \text{ so } Q_1 = 30 - 5.4 = \text{awrt } \underline{\underline{24.6}}$	B1ft (1)
(d)	$Q_3 - Q_1 = 10.8 \text{ so } 1.5(Q_3 - Q_1) = 16.2 \text{ so } Q_1 - 16.2 = h \text{ or } Q_3 + \quad = k$ $h = \underline{\underline{8.4 \text{ to } 8.6}} \text{ and } k = \underline{\underline{51.4 \text{ to } 51.6}} \qquad \text{both}$	12. M1 A1 (2)
(e)	$2P(D > 51.6) = 2P(Z > 2.7)$ $= 2[1 - 0.9965] = \text{awrt } \underline{\underline{0.007}}$	M1 M1 A1 (3)
		Total 12
(a)	M1 for an attempt to standardise 20 or 40 using 30 and 8. 1 st A1 for $z = \pm 1.25$ 2 nd A1 for awrt 0.894	
(b)	M1 for $\frac{Q_3 - 30}{8} =$ to a z value M0 for 0.7734 on RHS. B1 for (z value) between 0.67~0.675 seen. M1B0A1 for use of $z = 0.68$ in correct expression with awrt 35.4	
(c)	Follow through using their of quartile values.	
(d)	M1 for an attempt to calculate 1.5(IQR) and attempt to add or subtract using one of the formulae given in the question - follow through their quartiles	
(e)	1 st M1 for attempting $2P(D > \text{their } k)$ or $(P(D > \text{their } k) + P(D < \text{their } h))$ 2 nd M1 for standardising their h or k (may have missed the 2) so allow for standardising $P(D > 51.6)$ or $P(D < 8.4)$ Require boths Ms to award A mark.	