

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

Tuesday 18 June 2019

Morning (Time: 1 hour 30 minutes)

Paper Reference **WMA12/01**

Mathematics

International Advanced Subsidiary/Advanced Level
Pure Mathematics P2

You must have:

Mathematical Formulae and Statistical Tables (Lilac), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ►

P62721A

©2019 Pearson Education Ltd.

1/1/1/



Pearson

Answer all questions. Write your answers in the spaces provided.

1. A sequence $a_1, a_2, a_3, \dots$ is defined by

$$a_{n+1} = 4 - a_n$$

$$a_1 = 3$$

starting
point

recursive sequence
(plugging in a term
on the right gives
the next term on
the left)

Find the value of

(a) (i) $a_2 \rightarrow a_2 = a_{1+1} \quad (n=1)$

(ii) a_{107}

(2)

(b) $\sum_{n=1}^{200} (2a_n - 1)$ terms
start value

(2)

(a) This is a recursive sequence :

$$a_{n+1} = 4 - a_n \rightarrow \text{notice how } n+1 \text{ is 1 more than } n$$

Let's find :

(i) a_2 :

$$a_2 = 4 - a_1 = 4 - 3 = 1$$

(ii) a_{107} :

We cannot find a_{107} directly using the given formula. BUT :

$$a_3 = 4 - a_2 = 4 - 1 = 3$$

$$a_4 = 4 - a_3 = 4 - 3 = 1$$

$$a_1 = 3, a_2 = 1, a_3 = 3, a_4 = 1, \dots$$

The series repeats itself every 2 terms;
it is periodic.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 1 continued

$$\underbrace{3 \quad / \quad 1}_1, \underbrace{3 \quad / \quad 1}_2, \underbrace{3 \quad / \quad 1}_3, \dots$$

so : $\frac{107}{2 \text{ terms}} = 53 + 1$ remainder, one more term

$\swarrow$ 53 repeats of 3,1

$$\underbrace{3 \quad / \quad 1}_1, \underbrace{3 \quad / \quad 1}_2, \underbrace{3 \quad / \quad 1}_3, \dots, \underbrace{3 \quad / \quad 1}_{53}, \underbrace{3}_1 \text{ one more!}$$

so : $a_{107} = 3$

(b)

$\sum_{n=1}^{200} (2a_n - 1)$ means start the sequence at $n=1$ and sum all the terms up to $n=200$

$$= (2a_1 - 1) + (2a_2 - 1) + (2a_3 - 1) + \dots + (2a_{200} - 1)$$

Since the series repeats itself :

$$2(3) - 1 = 5$$

$$2(1) - 1 = 1$$

$$\underline{5 + 1} + \underline{5 + 1} + \underline{5 + 1} + \dots$$

we end at $n=200$: $\frac{200}{2 \text{ terms}} = 100 \text{ repeats } (5+1)$

$$\underbrace{5 + 1}_1 + \underbrace{5 + 1}_2 + \dots + \underbrace{5 + 1}_{100}$$

so sum up to $a_{200} = 100(5+1)$
 $= 600$

Q1

(Total 4 marks)



2. A circle C has equation

$$x^2 + y^2 + 4x - 10y - 21 = 0$$

Find

- (a) (i) the coordinates of the centre of C ,
(ii) the exact value of the radius of C .

(3)

The point $P(5, 4)$ lies on C .

- (b) Find the equation of the tangent to C at P , writing your answer in the form $y = mx + c$, where m and c are constants to be found.

(4)

(a) Standard form of equation of a circle:

$$(x - x_{\text{centre}})^2 + (y - y_{\text{centre}})^2 = r^2$$

↳ rewrite given eq. into standard form :

$$x^2 + 4x + y^2 - 10y - 21 = 0$$

(1) complete the square for terms with x :

$$x^2 + 4x + y^2 - 10y = 21$$

↳ add $\left(\frac{b}{2}\right)^2$ to both sides :

$$x^2 + 4x + \left(\frac{4}{2}\right)^2 + y^2 - 10y = 21 + \left(\frac{4}{2}\right)^2$$

$$x^2 + 4x + 4 + y^2 - 10y = 21 + 4$$

$$(x + 2)^2 + y^2 - 10y = 25$$

↳ factorise

$$(x + 2)^2 + y^2 - 10y = 25$$



Question 2 continued

(3) complete the square for terms with y :

$$(x+2)^2 + y^2 - 10y = 25$$

↳ add $\left(\frac{b}{2}\right)^2$ to both sides :

$$(x+2)^2 + y^2 - 10y + \left(\frac{-10}{2}\right)^2 = 25 + \left(\frac{-10}{2}\right)^2$$


$$(x+2)^2 + y^2 - 10y + 25 = 25 + 25$$

↳ factorise

$$(x+2)^2 + (y-5)^2 = 50$$

$$\text{So : } (x - (-2))^2 + (y - 5)^2 = 50$$

$$(x - (-2))^2 + (y - 5)^2 = 50$$

(i) coords of centre = $(-2, 5)$ (ii) radius = $\sqrt{50} = 12$ eq. of tangent $\rightarrow y = mx + c$ radius XP is perpendicular to tangent at P

$$\text{So : } m_{\text{tangent}} = -\frac{1}{m_r}$$

Q2

(Total 7 marks)



Question 2 continued

$$m_r = \frac{y_2 - y_1}{x_2 - x_1}$$

use : (5, 4) and (-2, 5)

$$m_r = \frac{5 - 4}{-2 - 5} = -\frac{1}{7}$$

$$m_{\text{tangent}} = -\frac{1}{-1/7} = 7$$

$$y = mx + c$$

$$y = 7x + c$$

FINDING c :

Method 1 : PLUG (4, -1) IN $y = mx + c$, FIND c

$$y = 7x + c$$

$$-1 = 7(4) + c$$

$$-1 = 28 + c$$

$$c = -29$$

So: $y = 7x - 29$ → eq. of tangent

OR :

★ Method 2 : USE : $y - y_1 = m(x - x_1)$

$$y - 4 = 7(x - 5)$$

$$y - 4 = 7x - 35$$

$$y = 7x - 31$$

note : this
method is faster,
gives equation
directly !

3. (i) Use algebra to prove that for all real values of x

$$(x-4)^2 \geq 2x-9 \quad (3)$$

- (ii) Show that the following statement is untrue.

$2^n + 1$ is a prime number for all values of n , $n \in \mathbb{N}$

(i) $(x-4)^2 \geq 2x-9 \quad (1)$

• expand $(x-4)^2$:

$$x^2 - 8x + 16 \geq 2x - 9$$

• move all terms to one side:

$$x^2 - 8x + 16 \geq 2x - 9$$

$$x^2 - 8x - 2x + 9 + 16 \geq 0$$

$$x^2 - 10x + 25 \geq 0$$

• factorise the inequality:

$$x^2 - 10x + 25 \geq 0$$

$$(x-5)(x-5) \geq 0$$

$$(x-5)^2 \geq 0$$

We know that square numbers are always greater than or equal to zero

so: $(x-5)^2 \geq 0$ is true

hence: $(x-4)^2 \geq 2x-9$ is true

(ii) Here, you can give a counter-example:

ex: $n = 3 \rightarrow 2^3 + 1 = 9$
 $= 3 \times 3$ so NOT PRIME

$n = 5 \rightarrow 2^5 + 1 = 33$
 $= 3 \times 11$ so NOT PRIME



4. (a) Find the first four terms, in ascending powers of x , of the binomial expansion of

$$\left(2 - \frac{1}{4}x\right)^6 \quad (4)$$

- (b) Given that x is small, so terms in x^4 and higher powers of x may be ignored, show

$$\left(2 - \frac{1}{4}x\right)^6 + \left(2 + \frac{1}{4}x\right)^6 = a + bx^2$$

where a and b are constants to be found.

(3)

(a) Binomial expansion formula :

$$(a + bx)^n = {}^nC_0 a^n (bx)^0 + {}^nC_1 a^{n-1} (bx)^1 + {}^nC_2 a^{n-2} (bx)^2 + \dots$$

$${}^nC_r = \frac{n!}{(n-r)!r!}$$

n : power of binomial
 r : (position of term) - 1

OR: USE CHOOSE FUNCTION ON CALCULATOR

for $\left(2 - \frac{1}{4}x\right)^6$:

$$= {}^6C_0 2^6 \left(\frac{-x}{4}\right)^0 + {}^6C_1 2^5 \left(\frac{-x}{4}\right)^1 + {}^6C_2 2^4 \left(\frac{-x}{4}\right)^2 + {}^6C_3 2^3 \left(\frac{-x}{4}\right)^3$$

$${}^6C_0 = 1$$

$${}^6C_1 = 6$$

$${}^6C_2 = \frac{6!}{(6-2)!2!} = 15$$

$${}^6C_3 = \frac{6!}{(6-3)!3!} = 20$$

NOTES:-

• the first and last nC_r are always = 1

• nC_1 is always equal to n

• BINOMIAL COEFFICIENTS ARE SYMMETRIC

for $(a+x)^6$:

$${}^6C_0 = {}^6C_6$$

$${}^6C_2 = {}^6C_4$$

$${}^6C_1 = {}^6C_5$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 4 continued

$$= (1)(64) + (6)(32)\left(\frac{-x}{4}\right) + (15)(16)\left(\frac{x^2}{16}\right) + (20)(8)\left(\frac{-x^3}{64}\right)$$

$$= 64 - 48x + 15x^2 - \frac{5}{2}x^3$$

(b) look at the two fractions :

$$\left(2 - \frac{1}{4}x\right)^6$$

$$\left(2 + \frac{1}{4}x\right)^6$$

only the sign between terms changed from -ve to +ve.

Therefore, the expansion of $\left(2 + \frac{1}{4}x\right)^6$ is the same as the expansion of $\left(2 - \frac{1}{4}x\right)^6$, except all terms become positive :

$$\left(2 - \frac{1}{4}x\right)^6 = 64 - 48x + 15x^2 - \frac{5}{2}x^3$$

$$\left(2 + \frac{1}{4}x\right)^6 = 64 + 48x + 15x^2 + \frac{5}{2}x^3$$

$$\left(2 - \frac{1}{4}x\right)^6 + \left(2 + \frac{1}{4}x\right)^6 \rightarrow \text{add the expansions!}$$

$$64 - 48x + 15x^2 - \frac{5}{2}x^3 + 64 + 48x + 15x^2 + \frac{5}{2}x^3$$

cancel

$$= 128 + 30x^3$$

Q4

(Total 7 marks)



5. A company makes a particular type of watch.

The annual profit made by the company from sales of these watches is modelled by the equation

$$P = 12x - x^{\frac{3}{2}} - 120$$

where P is the annual profit measured in thousands of pounds and $\pounds x$ is the selling price of the watch.

According to this model,

- (a) find, using calculus, the maximum possible annual profit. (6)
- (b) Justify, also using calculus, that the profit you have found is a maximum. (2)

(a) maximum profit = maximum point, which occurs when :

$$\frac{dP}{dx} = 0$$

DIFFERENTIATION:

$$y = ax^n \quad \frac{dy}{dx} = nax^{n-1}$$

$$P = 12x - x^{3/2} - 120$$

$$\frac{dP}{dx} = (1)(12)x^{1-1} - \left(\frac{3}{2}\right)x^{3/2-1} - (1)120x^{1-1}$$

$$\frac{dP}{dx} = 12 - \frac{3}{2}x^{1/2}$$

$$12 - \frac{3}{2}x^{1/2} = 0$$

$$\frac{2}{3} \times (12) = \left(\frac{3}{2}x^{1/2}\right) \times \frac{2}{3}$$

$$8 = x^{1/2}$$



Question 5 continued

$$(8)^2 = (x^{1/2})^2 \rightarrow (x^a)^b = x^{ab}$$

$$x = \text{£ } 64 \text{ per watch}$$

maximum selling price is P at $x = 64$

$$P = 12(64) - 64^{3/2} - 120$$

$$P = 768 - 512 - 120$$

$$= \text{£ } 136 \text{ 000}$$

(b) if $\frac{d^2P}{dx^2} > 0 \rightarrow$ minimum point



$\frac{d^2P}{dx^2} = 0 \rightarrow$ point of inflection



$\frac{d^2P}{dx^2} < 0 \rightarrow$ maximum point



$$\frac{dP}{dx} = 12 - \frac{3}{2}x^{1/2}$$

$$\frac{d^2P}{dx^2} = \frac{1}{2} \left(\frac{3}{2} \right) x^{1/2-1}$$

$$\frac{d^2P}{dx^2} = -\frac{3}{4}x^{-1/2}$$

At $x = 64 : \frac{d^2P}{dx^2} = -\frac{3}{4}(64)^{-1/2} = -0.09375$

$-0.09375 < 0$; it is a maximum

Q5

(Total 8 marks)



6. $f(x) = kx^3 - 15x^2 - 32x - 12$ where k is a constant

Given $(x - 3)$ is a factor of $f(x)$,

- (a) show that $k = 9$

(2)

- (b) Using algebra and showing each step of your working, fully factorise $f(x)$.

(4)

in degrees!

- (c) Solve, for $0 \leq \theta < 360^\circ$, the equation

$$9 \cos^3 \theta - 15 \cos^2 \theta - 32 \cos \theta - 12 = 0$$

giving your answers to one decimal place.

(2)

(a) FACTOR THEOREM :

If $(x - p)$ is a factor of $f(x)$, $f(p) = 0$.

So: $(x - 3)$ is a factor; $f(3) = 0$

• SOLVE $f(3) = 0$:

$$f(x) = kx^3 - 15x^2 - 32x - 12$$

$$f(3) = k(3)^3 - 15(3)^2 - 32(3) - 12 = 0$$

$$27k - 135 - 96 - 12 = 0$$

$$27k - 243 = 0$$

$$27k = 243$$

$$k = 9$$

(b) Use algebraic division to factorise :

$$\begin{array}{r} x-3 \overline{) 9x^3 - 15x^2 - 32x + 12} \end{array}$$

(1) Divide first term of dividend by first term of divisor : $\frac{9x^3}{x} = 9x^2 \rightarrow$ first term of quotient

$$\begin{array}{r} 9x^2 \\ x-3 \overline{) 9x^3 - 15x^2 - 32x + 12} \\ \underline{9x^3 - 27x^2} \\ -8x^2 - 32x + 12 \end{array}$$



Question 6 continued

(2) Multiply out and write under :

$$\begin{array}{r} \overline{9x^3 - 15x^2 - 32x - 12} \\ \underline{9x^3 - 3x^2} \end{array}$$

(3) Subtract then bring down next term :

$$\begin{array}{r} \overline{9x^3 - 15x^2 - 32x - 12} \\ \underline{-(9x^3 - 27x^2)} \\ 12x^2 - 32x \end{array}$$

(4) Divide first term of expression by first term of divisor : $\frac{12x^2}{x} = +12x \rightarrow$ next term of quotient

$$\begin{array}{r} \overline{9x^3 - 15x^2 - 32x - 12} \\ \underline{-(9x^3 - 27x^2)} \\ 12x^2 - 32x \end{array}$$

(5) Multiply out and write under :

$$\begin{array}{r} \overline{9x^3 - 15x^2 - 32x - 12} \\ \underline{-(9x^3 - 27x^2)} \\ 12x^2 - 32x \\ \underline{12x^2 - 36x} \end{array}$$

(6) Subtract then bring down next term :

$$\begin{array}{r} \overline{9x^3 - 15x^2 - 32x - 12} \\ \underline{-(9x^3 - 27x^2)} \\ 12x^2 - 32x \\ \underline{-(12x^2 - 36x)} \\ 4x - 12 \end{array}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 6 continued

(7) Divide first term of expression by first term of divisor : $\frac{4x}{x} = 4 \rightarrow$ next term of quotient

$$\begin{array}{r}
 9x^2 + 12x + 4 \\
 x-3 \overline{) 9x^3 - 15x^2 - 32x - 12} \\
 \underline{-(9x^3 - 27x^2)} \\
 12x^2 - 32x \\
 \underline{-(12x^2 - 36x)} \\
 4x - 12
 \end{array}$$

(8) Multiply out and write under:

$$\begin{array}{r}
 9x^2 + 12x + 4 \\
 x-3 \overline{) 9x^3 - 15x^2 - 32x - 12} \\
 \underline{-(9x^3 - 27x^2)} \\
 12x^2 - 32x \\
 \underline{-(12x^2 - 36x)} \\
 4x - 12 \\
 4x - 12
 \end{array}$$

(9) Subtract:

$$\begin{array}{r}
 9x^2 + 12x + 4 \\
 x-3 \overline{) 9x^3 - 15x^2 - 32x - 12} \\
 \underline{-(9x^3 - 27x^2)} \\
 12x^2 - 32x \\
 \underline{-(12x^2 - 36x)} \\
 4x - 12 \\
 \underline{4x - 12} \\
 0
 \end{array}$$

So: $f(x) = (x-3)(9x^2 + 12x + 4) \rightarrow$ factorise further!
 $= (x-3)(3x+2)^2$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 6 continued

$$(c) \quad 9 \cos^3 \theta - 15 \cos^2 \theta - 32 \cos \theta - 12 = 0$$

$$9x^3 - 15x^2 - 32x - 12 = 0$$

$$\text{so } x = \cos \theta$$

therefore, we can rewrite:

$$9 \cos^3 \theta - 15 \cos^2 \theta - 32 \cos \theta - 12 = 0$$

$$(\cos \theta - 3)(3 \cos \theta + 2)^2 = 0$$

$$\cos \theta - 3 = 0 \quad \text{REJECT!}$$

$$\cos \theta = 3 \rightarrow \text{range of } y = \cos \theta \text{ is } -1 < \theta < 1$$

OR

$$3 \cos \theta + 2 = 0$$


$$\cos \theta = -\frac{2}{3}$$

$$\text{SOLVE: } \cos \theta = -\frac{2}{3}, \quad 0 \leq \theta < 360^\circ$$

ON CALCULATOR:

$$\theta = \cos^{-1}\left(-\frac{2}{3}\right) = 131.8^\circ \quad (1 \text{ d.p.})$$

ON UNIT CIRCLE:



$$\text{where } \cos \theta = -\frac{2}{3},$$

$$\theta = 131.8^\circ$$

$$\theta = 360 - 131.8^\circ$$

$$= 228.8^\circ$$

So:

$$\theta = 131.8^\circ, 228.8^\circ$$

FINAL ANSWER!

(Total 8 marks)

Q6



7. Kim starts working for a company.

- In year 1 her annual salary will be £16200 a_1
- In year 10 her annual salary is predicted to be £31500 a_{10}

Model A assumes that her annual salary will increase by the same amount each year.

arithmetic series

(a) According to model A, determine Kim's annual salary in year 2.

geometric series (3)

Model B assumes that her annual salary will increase by the same percentage each year.

(b) According to model B, determine Kim's annual salary in year 2. Give your answer to the nearest £10

partial sum (3)

(c) Calculate, according to the two models, the difference between the total amounts that Kim is predicted to earn from year 1 to year 10 inclusive. Give your answer to the nearest £10

(3)

(a) In model A, we can find d , then use it to find salary in year 2:

year : 1 , 2 , ... , 10
 salary : £16200 , a_2 , ... , £31500

$+d$ $+d$ 8 more times

$$31500 = 16200 + 9d$$

$$d = \frac{31500 - 16200}{9}$$

$$d = 1700$$

now : $a_n = a + d(n-1)$

$$a_2 = 16200 + (1700)(2-1)$$

$$a_2 = £17900$$

(b) In model B, we can find r then use it to find salary in year 2:



Question 7 continued

year : 1 / 2 / ... / 10
 salary : £16 200 / a_2 / ... / £31 500

$\times r$ $\times r$ 8 more times

So : 31 500 = 16 200 $\times r^9$ $\rightarrow$ multiplying by r 9 times.

$$r^9 = \frac{31\,500}{16\,200}$$

$$r = \sqrt[9]{\frac{31\,500}{16\,200}} = 1.07668\dots$$

$$\begin{aligned} a_n &= ar^{n-1} \\ a_2 &= 16\,200(1.07668\dots)^{2-1} \\ &= 17\,442 = \text{£}17\,440 \text{ (nearest £10)} \end{aligned}$$

(c) ① total for A :

$$S_n = \frac{n}{2}(a_1 + a_n)$$

$$\begin{aligned} S_{10} &= \frac{10}{2}(16\,200 + 31\,500) \\ &= \text{£}238\,500 \end{aligned}$$

② total for B :

$$S_n = \frac{a(1-r^n)}{1-r}$$

$$S_{10} = \frac{16\,200(1 - [1.07668\dots]^{10})}{(1 - [1.07668\dots])}$$

$$= \text{£}231\,019$$



Question 7 continued

$$\text{difference} = 238\,500 - 231\,019$$

$$= 7481 = \text{£}7480 \text{ (nearest } \text{£}10)$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q7

(Total 9 marks)



8. (i) Find the exact solution of the equation

$$8^{2x+1} = 6$$

giving your answer in the form $a + b \log_2 3$, where a and b are constants to be found.

(4)

- (ii) Using the laws of logarithms, solve

$$\log_5(7 - 2y) = 2 \log_5(y + 1) - 1$$

(5)

Recall rules of logarithms :

$$\left[\begin{array}{l} \log_a(b) + \log_a(c) = \log_a(b \times c) \\ \log_a(b) - \log_a(c) = \log_a\left(\frac{b}{c}\right) \\ \log_a b = b \log_a a \end{array} \right]$$

(i) note!

Q wants final answer to include $\log_2 3$

take logs of both sides, (base 2)

$$\log_2 8^{2x+1} = \log_2 6$$

$$\log_2 8^{2x+1} = \log_2 6$$

$$(2x+1) \log_2 8 = \log_2 6$$

We know $8 = 2^3$

$$2x+1 (\log_2 8) = \log_2 6$$

$$(2x+1) (\log_2 (2^3)) = \log_2 6$$

$$(2x+1) (3 \log_2 2) = \log_2 6 \rightarrow \log_2 2 = 1$$

$$(2x+1)(3) = \log_2 6$$

$$\frac{(2x+1)(3)}{3} = \frac{\log_2 6}{3}$$



Question 8 continued

$$2x + 1 = \frac{\log_2 6}{3}$$

$$2x + 1 = \frac{\log_2 (2 \times 3)}{3}$$

$$2x + 1 = \frac{\log_2 (2 \times 3)}{3}$$

$$2x + 1 = \frac{\log_2 2 + \log_2 3}{3}$$

$$2x + 1 = \frac{1 + \log_2 3}{3}$$

$$2x = \frac{1}{3} + \frac{\log_2 3}{3} - 1$$

$$2x = \frac{\log_2 3}{3} - \frac{2}{3}$$

divide both sides by 2 :

$$x = \frac{\log_2 3}{6} - \frac{1}{6}$$

$$x = \frac{\log_2 3}{6} - \frac{1}{6}$$

ALTERNATIVE SOLUTION :

$$8^{2x+1} = 6$$

we know $8 = 2^3$

$$(2^3)^{2x+1} = 6$$

use rule : $(a^b)^c = a^{b \times c}$

$$2^{3(2x+1)} = 6$$

$$2^{6x+3} = 6$$



Question 8 continued

take $\log_2$ of both sides :

$$\log_2 2^{6x+3} = \log_2 6$$

$$\log_2 2^{6x+3} = \log_2 6$$

$$(6x+3) \log_2 2 = \log_2 6$$

$$(6x+3)(1) = \log_2 6$$

$$6x+3 = \log_2 (2 \times 3)$$

$$6x+3 = \log_2 (2 \times 3)$$

$$6x+3 = \log_2 2 + \log_2 3$$

$$6x+3 = 1 + \log_2 3$$

make x subject :

$$6x+3 = 1 + \log_2 3$$

$$6x = 1 - 3 + \log_2 3$$

$$6x = -2 + \log_2 3$$

$$x = \frac{-2}{6} + \frac{\log_2 3}{6}$$

$$x = -\frac{1}{3} + \frac{\log_2 3}{6}$$

$$(ii) \log_5(7-2y) = 2\log_5(y+1) - 1$$

$$\log_5(7-2y) - 2\log_5(y+1) = -1$$

$$\log_5(7-2y) - \log_5(y+1)^2 = -1$$

$$\log_5(7-2y) - \log_5(y+1)^2 = -1$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 8 continued

$$\log_5 \left(\frac{7-2y}{(y+1)^2} \right) = -1$$

use rule : $\log_a b = c \rightarrow a^c = b$

$$\log_5 \left(\frac{7-2y}{(y+1)^2} \right) = -1$$

$$\frac{7-2y}{(y+1)^2} = 5^{-1} \rightarrow a^{-b} = \frac{1}{a^b}$$

$$\frac{7-2y}{(y+1)^2} = \frac{1}{5}$$

multiply both sides by $(y+1)^2$

$$(y+1)^2 \left(\frac{7-2y}{(y+1)^2} \right) = \left(\frac{1}{5} \right) (y+1)^2$$

$$7-2y = \frac{1}{5} (y+1)^2$$

expand $(y+1)^2$:

$$7-2y = \frac{1}{5} (y^2 + 2y + 1)$$

$$7-2y = \frac{1}{5} (y^2 + 2y + 1)$$

$$7-2y = \frac{1}{5} y^2 + \frac{2}{5} y + \frac{1}{5}$$

multiply both sides by 5 :

$$5 \times (7-2y) = \left(\frac{1}{5} y^2 + \frac{2}{5} y + \frac{1}{5} \right) \times 5$$

$$35 - 10y = y^2 + 2y + 1$$

(Total 9 marks)

Q8



Question 8 continued

$$y^2 + 2y + 1 + 10y - 35 = 0$$

$$y^2 + 12y - 34 = 0 \rightarrow \text{factorise!}$$

$$1y^2 + 12y - 34 = 0$$

$$y = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-12 \pm \sqrt{12^2 - 4(1)(-34)}}{2(1)}$$

$$= 6 \pm \sqrt{70}$$

$$y = 6 + \sqrt{70}$$

$$y = 6 - \sqrt{70} \quad \times \text{reject:}$$

$\log_2(6 - \sqrt{70} + 1)$ is
-ve, log of a -ve
value is not possible

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



9. (a) Show that the equation

$$\cos \theta - 1 = 4 \sin \theta \tan \theta$$

can be written in the form

$$5 \cos^2 \theta - \cos \theta - 4 = 0 \quad (4)$$

- (b) Hence solve, for
- $0 \leq x < \frac{\pi}{2}$
- IN RADIANS !!**

$$\cos 2x - 1 = 4 \sin 2x \tan 2x$$

giving your answers, where appropriate, to 2 decimal places.

(4)

$$(a) \cos \theta - 1 = 4 \sin \theta \tan \theta \rightarrow \tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\cos \theta - 1 = 4 \sin \theta \left(\frac{\sin \theta}{\cos \theta} \right)$$

$$\cos \theta \times (\cos \theta - 1) = \frac{4 \sin^2 \theta}{\cos \theta} \times \cos \theta$$

$$\cos^2 \theta - \cos \theta = 4 \sin^2 \theta$$

Use identity : $\sin^2 \theta + \cos^2 \theta = 1$
 $\sin^2 \theta = 1 - \cos^2 \theta$

$$\cos^2 \theta - \cos \theta = 4(1 - \cos^2 \theta)$$

$$\cos^2 \theta - \cos \theta = 4 - 4 \cos^2 \theta$$

$$5 \cos^2 \theta - \cos \theta - 4 = 0$$

$$(b) \begin{array}{l} \cos 2x - 1 = 4 \sin 2x \tan 2x \\ \cos \theta - 1 = 4 \sin \theta \tan \theta \end{array} \left. \begin{array}{l} \\ \end{array} \right\} \begin{array}{l} \text{same} \\ \text{structure} \end{array}$$

$$\text{SO: } 5 \cos^2 2x - \cos 2x - 4 = 0 \rightarrow \text{FACTORISE!}$$

$$(5 \cos 2x + 4)(\cos 2x - 1) = 0$$

$$5 \cos 2x + 4 = 0 \quad \text{or} \quad \cos 2x - 1 = 0$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 9 continued


$$\textcircled{1} \quad 5\cos 2x + 4 = 0 \quad , \quad 2(0) < 2x < 2(\pi/2)$$

$$\cos 2x = \frac{-4}{5}$$

ON CALCULATOR :

$$2x = \cos^{-1}\left(\frac{-4}{5}\right) = 2.50$$


• note : there are no other angles on the unit circle within our given range where $\cos 2x = -4/5$

ALLOWED
DOMAIN

So $2x = 2.50$

$$\textcircled{2} \quad \cos 2x - 1 = 0$$

$$\cos 2x = 1$$



So : $2x = 0$ when $\cos 2x = 1$

$$2x = 0, 2.50$$

$$x = 0, 1.25$$



Question 9 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q9

(Total 8 marks)



10.




Figure 1

Figure 1 shows a sketch of part of the curve C with equation $y = f(x)$ where

$$f(x) = \frac{36}{x^2} + 2x - 13 \quad x > 0$$

Using calculus,

(a) find the range of values of x for which $f(x)$ is increasing, (4)

(b) show that $\int_2^9 \left(\frac{36}{x^2} + 2x - 13 \right) dx = 0$ (4)

The point $P(2, 0)$ and the point $Q(6, 0)$ lie on C .

Given $\int_2^6 \left(\frac{36}{x^2} + 2x - 13 \right) dx = -8$

(c) (i) state the value of $\int_6^9 \left(\frac{36}{x^2} + 2x - 13 \right) dx$
 (ii) find the value of the constant k such that $\int_2^6 \left(\frac{36}{x^2} + 2x + k \right) dx = 0$ (3)

(a) Where $f(x)$ is increasing, there is a positive gradient: $f'(x) > 0$

• Find $f'(x)$ of $f(x)$:

$$f(x) = \frac{36}{x^2} + 2x - 13$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 10 continued

$$f(x) = 36x^{-2} + 2x - 13$$

$$f(x) = ax^n \rightarrow f'(x) = nax^{n-1} \quad \text{differentiation rule :)$$

$$f'(x) = (-2)36x^{-2-1} + (1)2x^{1-1} - (0)13x^{0-1}$$

$$f'(x) = -72x^{-3} + 2$$

$$= -\frac{72}{x^3} + 2$$

$$f'(x) > 0$$

$$\left[-\frac{72}{x^3} \right] + 2 > 0$$

$$2 > \frac{72}{x^3}$$

• multiply both sides by x^3 :

$$x^3 \times (2) > \left(\frac{72}{x^3} \right) \times x^3$$

$$2x^3 > 72$$

• divide by 2 :

$$\frac{2x^3}{2} > \frac{72}{2}$$

$$x^3 > 36 \rightarrow x > \sqrt[3]{36}$$

interval
where $f(x)$
is increasing!

$$(b) \int_2^9 \frac{36}{x^2} + 2x - 13 \, dx$$

$$= \int_2^9 36x^{-2} + 2x - 13 \, dx$$

REMEMBER!

$$\int x^n \, dx = \frac{x^{n+1}}{n+1}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 10 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

$$\left[\frac{36x^{-2+1}}{-2+1} + \frac{2x^{1+1}}{1+1} - \frac{13x^{0+1}}{0+1} \right]_2^9$$

$$\left[\frac{36x^{-2+1}}{-2+1} + \frac{2x^{1+1}}{1+1} - \frac{13x^{0+1}}{0+1} \right]_2^9$$


$$\left[-36x^{-1} + x^2 - 13x \right]_2^9 \quad (\text{RULE: } x^{-a} = 1/x^a)$$

$$\left[\frac{-36}{x} + x^2 - 13x \right]_2^9$$

$$= \left[\frac{-36}{9} + 9^2 - 13(9) \right] - \left[\frac{-36}{2} + 2^2 - 13(2) \right]$$

$$= (-4 + 81 - 117) - (-18 + 4 - 26)$$

$$= -40 - (-40) = 0$$

(c)(i) If : 

$$\int_a^c f(x) dx = \int_a^b f(x) dx + \int_b^c f(x) dx$$

So :

$$\int_2^9 \frac{36}{x^2} + 2x - 13 dx = \int_2^6 \frac{36}{x^2} + 2x - 13 dx + \int_6^9 \frac{36}{x^2} + 2x - 13 dx$$

$$0 = -8 + \int_6^9 \frac{36}{x^2} + 2x - 13 dx$$

$$\int_6^9 \frac{36}{x^2} + 2x - 13 dx = 8$$



Question 10 continued

(ii)

$$\int_2^6 \left(\frac{36}{x^2} + 2x + k \right) dx$$

INTEGRATE : (see steps in (b))

$$\left[-\frac{36}{x} + x^2 + kx \right]_2^6$$

$$= \left[-\frac{36}{6} + 6^2 + k(6) \right] - \left[-\frac{36}{2} + 2^2 + k(2) \right]$$

$$= -6 + 36 + 6k - (-18 + 4 + 2k)$$

$$= 30 + 6k + 14 - 2k$$

$$= 44 + 4k$$

$$\int_2^6 \left(\frac{36}{x^2} + 2x + k \right) dx = 0$$

$$44 + 4k = 0$$

$$4k = -44$$

$$k = -11$$



Leave
blank

Question 10 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q10

(Total 11 marks)

TOTAL FOR PAPER IS 75 MARKS

END

