

Please check the examination details below before entering your candidate information

Candidate surname		Other names	
Centre Number		Candidate Number	
Pearson Edexcel International Advanced Level			
Tuesday 15 October 2024			
Morning (Time: 1 hour 30 minutes)		Paper reference	WMA12/01
Mathematics International Advanced Subsidiary/Advanced Level Pure Mathematics P2			
You must have: Mathematical Formulae and Statistical Tables (Yellow), calculator			Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 11 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ►

P76187RA

©2024 Pearson Education Ltd.
V:1/1/1/1/1/



1. A continuous curve has equation $y = f(x)$.

A table of values of x and y for $y = f(x)$ is shown below.

Interval btwn two x -coords = h of trapezium

x	0.5	1.75	3	4.25	5.5
y	3.479	6.101	7.448	6.823	5.182

y_1

y_6

multiplied by 2 as all are shared by two trapezia

Using the trapezium rule with all the values of y in the given table,

- (a) find an estimate for

$$\int_{0.5}^{5.5} f(x) dx$$


giving your answer to one decimal place.

(3)

- (b) Using your answer to part (a) and making your method clear, estimate

$$\int_{0.5}^{5.5} (f(x) + 4x) dx$$

(2)



(Formula is given in formula booklet. The graph is only a visual representation. DO NOT DRAW A GRAPH IN THE EXAM)



Question 1 continued

$$A = \frac{1}{2} h \left[y_1 + 2(y_2 + y_3 + y_4) + y_6 \right]$$

$$A = \frac{1}{2} (1.25) \left[3.479 + 2(6.101 + 7.448 + 6.823) + 5.182 \right] = 30.8781...$$

$$\text{So: } \int_{0.5}^{5.5} f(x) dx \approx 30.8781... \\ = 30.9 \text{ (1 d.p.)}$$

(b)

$$\text{use rule: } \int f(x) + g(x) dx = \int f(x) dx + \int g(x) dx$$

$$\int_{0.5}^{5.5} f(x) + 4x dx = \int_{0.5}^{5.5} f(x) dx + \int_{0.5}^{5.5} 4x dx$$

$$\int_{0.5}^{5.5} f(x) dx + \int_{0.5}^{5.5} 4x dx$$

$$= 30.9 + \left[\frac{4x^{1+1}}{1+1} \right]_{0.5}^{5.5}$$

$$= 30.9 + [2x^2]_{0.5}^{5.5}$$

$$= 30.9 + (2(5.5)^2 - 2(0.5)^2)$$

$$= 30.9 + 60 = 90.9$$

REMEMBER!

$$\int x^n dx = \frac{x^{n+1}}{n+1}$$

(Total for Question 1 is 5 marks)



2. A sequence of numbers $u_1, u_2, u_3, \dots$ is defined by

$$u_{n+1} = (-1)^n u_n + k$$

recursive sequence
(plugging in a term
on the right gives
the next term on
the left)

where k is a constant.

(a) Show that $u_5 = 7$

end → terms: $u_1, u_2, u_3, \dots$

(3)

Given that $\sum_{r=1}^4 u_r = 30$

sum

starting value

(b) find the value of k .

(2)

(c) Hence find the value of

$$\sum_{r=1}^{150} u_r$$

(2)

(a) This is a recursive sequence :

$$u_{n+1} = (-1)^n u_n + k \rightarrow \text{notice how } n+1 \text{ is 1 more than } n$$

$$\text{If } u_1 = 7 :$$

$$\begin{aligned} u_{1+1} &= (-1)^1 (7) + k \\ &= (-1)(7) + k \\ \rightarrow u_2 &= -7 + k \end{aligned}$$

$$\begin{aligned} u_{2+1} &= (-1)^2 (-7 + k) + k \\ &= (1)(-7 + k) + k \\ \rightarrow u_3 &= -7 + 2k \end{aligned}$$

$$\begin{aligned} u_{3+1} &= (-1)^3 (-7 + 2k) + k \\ &= 7 - 2k + k \\ \rightarrow u_4 &= 7 - k \end{aligned}$$

$$\begin{aligned} u_{4+1} &= (-1)^4 (7 - k) + k \\ &= 1(7 - k) + k \\ \rightarrow u_5 &= 7 \end{aligned}$$

Question 2 continued

(b) $\sum_{r=1}^4 u_r = 30$ means start the sequence at $r=1$ and sum all the terms up to $r=4$

$$u_1 + u_2 + u_3 + u_4 = 30$$

$$7 + -7 + k + -7 + 2k + 7 - k = 30$$

$$2k = 30$$

$$k = 15$$

(c) $\sum_{r=1}^{150} u_r = u_1 + u_2 + u_3 + u_4 + \dots + u_{150}$

We have seen in (a) that the sequence repeats itself every 4 terms ($u_1 = u_5$).

$$7 + -7 + k + -7 + 2k + 7 - k + 7 + -7 + k + -7 + 2k + 7 - k + \dots$$

So: $\frac{150}{4 \text{ terms}} = 37 + 2$ remainder, 2 more terms
 $\rightarrow 37 \text{ repeats } (a_1 \text{ to } a_4)$

We know that $u_1 + u_2 + u_3 + u_4 = 30$.

$$\begin{aligned} \text{So: } \sum_{r=1}^{150} u_r &= 37(30) + 7 + (-7 + 15) \\ &= 1125 \end{aligned}$$

(Total for Question 2 is 7 marks)

3.

$$f(x) = 2x^3 - x^2 + Ax + B$$

where A and B are integers.

Given that when $f(x)$ is divided by $(x + 3)$ the remainder is 55

(a) show that

$$3A - B = -118$$

(2)

Given also that $(2x - 5)$ is a factor of $f(x)$, $\rightarrow$ no remainder when dividing

(b) find the value of A and the value of B .

(3)

(c) Hence find the quotient when $f(x)$ is divided by $(x - 7)$

(2)

(a) USE :

REMAINDER THEOREM :

if $\frac{f(x)}{(x-k)}$ = quotient + remainder :

then $f(k)$ = remainder

$$\text{if } \frac{f(x)}{(x - (-3))} = \text{quotient} + 55$$

$$\text{then } f(-3) = 55 :$$

$$2(-3)^3 - (-3)^2 + A(-3) + B = 55$$

$$-54 - 9 - 3A + B = 55$$

$$3A - B = -54 - 9 - 55$$

$$3A - B = -118$$

(b)

FACTOR THEOREM :

(no remainder)

if $\frac{f(x)}{(ax - k)}$ = quotient + 0, then $f\left(\frac{k}{a}\right) = 0$

$$\text{if } \frac{f(x)}{(2x - 5)} = 0, \text{ then } f\left(\frac{5}{2}\right) = 0 :$$



Question 3 continued

$$2\left(\frac{5}{2}\right)^3 - \left(\frac{5}{2}\right)^2 + A\left(\frac{5}{2}\right) + B = 0$$

$$\frac{125}{4} - \frac{25}{4} + \frac{5}{2}A + B = 0$$

$$2 \times \left(\frac{5}{2}A + B\right) = (-25) \times 2$$

$$5A + 2B = -50$$

NOW WE HAVE TWO EQ. :

$$\textcircled{1} 3A - B = -118$$

$$\textcircled{2} 5A + 2B = -50$$

SOLVE SIMULTANEOUSLY

 $\textcircled{1}$ elimination :

$$\begin{aligned} [3A - B = -118] \times 2 \\ 5A + 2B = -50 \end{aligned}$$

$$\begin{aligned} \oplus \quad 6A - 2B &= -236 \\ 5A + 2B &= -50 \\ \hline 11A &= -286 \end{aligned}$$

$$A = -26$$

$$\begin{aligned} 3(-26) - B &= -118 \\ B &= 40 \end{aligned}$$

$$\text{So : } \begin{aligned} A &= -26 \\ B &= 40 \end{aligned}$$

 $\textcircled{2}$ Substitution :

$$\begin{aligned} 3A - B &= -118 \\ B &= 3A + 118 \end{aligned}$$

$$\begin{aligned} 5A + 2B &= -50 \\ 5A + 2(3A + 118) &= -50 \\ 5A + 6A + 236 &= -50 \\ 11A &= -286 \\ A &= -26 \end{aligned}$$

$$\begin{aligned} B &= 3(-26) + 118 \\ B &= 40 \end{aligned}$$

Question 3 continued

$$(c) f(x) = 2x^3 - x^2 - 26x + 40$$

Use algebraic division to find quotient :

$$\begin{array}{r} x-7 \overline{) 2x^3 - x^2 - 26x + 40} \end{array}$$

(1) Divide first term of dividend by first term of divisor : $\frac{2x^3}{x} = 2x^2 \rightarrow$ first term of quotient

$$\begin{array}{r} 2x^2 \\ x-7 \overline{) 2x^3 - x^2 - 26x + 40} \end{array}$$

(2) Multiply out and write under :

$$\begin{array}{r} 2x^2 \\ x-7 \overline{) 2x^3 - x^2 - 26x + 40} \\ \underline{2x^3 - 14x^2} \end{array}$$

(3) Subtract then bring down next term :

$$\begin{array}{r} 2x^2 \\ x-7 \overline{) 2x^3 - x^2 - 26x + 40} \\ \underline{-(2x^3 - 14x^2)} \quad \downarrow \\ 13x^2 - 26x \end{array}$$

(4) Divide first term of expression by first term of divisor : $\frac{13x^2}{x} = 13x \rightarrow$ next term of quotient

$$\begin{array}{r} 2x^2 + 13x \\ x-7 \overline{) 2x^3 - x^2 - 26x + 40} \\ \underline{-(2x^3 - 14x^2)} \\ 13x^2 - 26x \end{array}$$



Question 3 continued

(8) Multiply out and write under:

$$\begin{array}{r}
 \text{ } \overline{) 2x^3 - x^2 - 26x + 40} \\
 \underline{-(2x^3 - 14x^2)} \\
 13x^2 - 26x \\
 \underline{-(13x^2 - 91x)} \\
 65x + 40
 \end{array}$$

(6) Subtract then bring down next term:

$$\begin{array}{r}
 \text{ } \overline{) 2x^3 - x^2 - 26x + 40} \\
 \underline{-(2x^3 - 14x^2)} \\
 13x^2 - 26x \\
 \underline{-(13x^2 - 91x)} \\
 65x + 40
 \end{array}$$

(7) Divide first term of expression by first term of divisor: $\frac{65x}{x} = 65 \rightarrow$ next term of quotient

$$\begin{array}{r}
 \text{ } \overline{) 2x^3 - x^2 - 26x + 40} \\
 \underline{-(2x^3 - 14x^2)} \\
 13x^2 - 26x \\
 \underline{-(13x^2 - 91x)} \\
 65x + 40
 \end{array}$$

(8) Multiply out and write under:

$$\begin{array}{r}
 \text{ } \overline{) 2x^3 - x^2 - 26x + 40} \\
 \underline{-(2x^3 - 14x^2)} \\
 13x^2 - 26x \\
 \underline{-(13x^2 - 91x)} \\
 65x + 40 \\
 \underline{-(65x - 455)} \\
 495
 \end{array}$$

(Total for Question 3 is 7 marks)

Question 3 continued

(9) Subtract

$$\begin{array}{r}
 \overline{2x^2 + 13x + 65} \\
 x-7 \overline{) 2x^3 - x^2 - 26x + 40} \\
 \underline{-(2x^3 - 14x^2)} \\
 13x^2 - 26x \\
 \underline{-(13x^2 - 91x)} \\
 65x + 40 \\
 \underline{-(65x - 455)} \\
 495
 \end{array}$$

so : $\frac{f(x)}{x-7} = \frac{2x^2 + 13x + 65}{x-7} + \frac{495}{x-7}$

$\swarrow$ quotient.
 $\searrow$ remainder!

quotient = $2x^2 + 13x + 65$

TOTAL FOR PAPER IS 75 MARKS



4.

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

The curve C has equation

$$y = 4x^{\frac{1}{2}} + 9x^{-\frac{1}{2}} + 3$$

$$x > 0$$

don't ignore this!

- (a) Find $\frac{dy}{dx}$ giving each term in simplest form. $\frac{dy}{dx} = 0$ (2)

- (b) Hence find the x coordinate of the stationary point of C . (2)

- (c) (i) Find $\frac{d^2y}{dx^2}$ giving each term in simplest form. (2)

- (ii) Hence determine the nature of the stationary point of C , giving a reason for your answer. (2)

- (d) State the range of values of x for which y is decreasing. (1)

(a)

DIFFERENTIATION

$$y = ax^n \rightarrow \frac{dy}{dx} = nax^{n-1}$$

$$y = 4x^{\frac{1}{2}} + 9x^{-\frac{1}{2}} + 3$$

$$\frac{dy}{dx} = \left(\frac{1}{2}\right) 4x^{\frac{1}{2}-1} + \left(-\frac{1}{2}\right) 9x^{-\frac{1}{2}-1}$$

$$= 2x^{-\frac{1}{2}} - \frac{9}{2}x^{-\frac{3}{2}}$$

(b)

$$\frac{dy}{dx} = 0 \text{ at stationary point}$$

$$2x^{-\frac{1}{2}} - \frac{9}{2}x^{-\frac{3}{2}} = 0$$

$$x^{\frac{3}{2}} \times \left(2x^{-\frac{1}{2}} - \frac{9}{2}x^{-\frac{3}{2}}\right) = (0) \times x^{\frac{3}{2}}$$

$$2x^{-\frac{1}{2}+\frac{3}{2}} - \frac{9}{2}x^{-\frac{3}{2}+\frac{3}{2}} = 0 \rightarrow (a^b \times a^c = a^{b+c})$$

$$2x - \frac{9}{2} = 0 \rightarrow x = \frac{9}{4}$$



Question 4 continued

$$(c)(i) \quad \frac{dy}{dx} = 2x^{-1/2} - \frac{9}{2}x^{-3/2}$$

$$\frac{d^2y}{dx^2} = \left(-\frac{1}{2}\right) 2x^{-1/2-1} - \left(-\frac{3}{2}\right) \frac{9}{2}x^{-3/2-1}$$

$$\frac{d^2y}{dx^2} = -x^{-3/2} + \frac{27}{4}x^{-5/2}$$

$$(ii) \quad \text{at } x = 9/4 :$$

$$\frac{d^2y}{dx^2} = -\left(\frac{9}{4}\right)^{-3/2} + \frac{27}{4}\left(\frac{9}{4}\right)^{-5/2} = \frac{16}{27}$$

RECALL:

if $\frac{d^2y}{dx^2} > 0 \rightarrow$ minimum point



$\frac{d^2y}{dx^2} = 0 \rightarrow$ point of inflection



$\frac{d^2y}{dx^2} < 0 \rightarrow$ maximum point



$\frac{16}{27} > 0$ so point is a minimum ☺

(d) $\frac{9}{4}$ is a minimum :

y is decreasing BEFORE $9/4$:

$$0 < x < 9/4$$

because range in question is for $x > 0$!!

(Total for Question 4 is 7 marks)

5. (a) Find, in terms of a , the first 3 terms, in ascending powers of x , of the binomial expansion of

$$(2 + ax)^6$$

where a is a non-zero constant. Give each term in simplest form.

(3)

$$f(x) = \left(3 + \frac{1}{x}\right)^2 (2 + ax)^6$$

Given that the constant term in the expansion of $f(x)$ is 576

- (b) find the value of a .

(4)

(a) Binomial expansion formula :

$$(a + bx)^n = {}^nC_0 a^n (bx)^0 + {}^nC_1 a^{n-1} (bx)^1 + {}^nC_2 a^{n-2} (bx)^2 + \dots$$

$${}^nC_r = \frac{n!}{(n-r)! r!}$$

n : power of binomial
 r : (position of term) - 1

OR USE CHOOSE FUNCTION ON CALCULATOR!

$$(2 + ax)^6 = {}^6C_0 2^6 (ax)^0 + {}^6C_1 2^5 (ax)^1 + {}^6C_2 2^4 (ax)^2 + \dots$$

$${}^6C_0 = 1$$

$${}^6C_1 = 6$$

$${}^6C_2 = \frac{6!}{(6-2)! 2!} = 15$$

NOTES:-

• the first and last nC_r are always = 1

• nC_1 is always equal to n

$$(2 + ax)^6 = 1(64)(1) + 6(32)(ax) + 15(16)(a^2 x^2) + \dots$$

$$= 64 + 192ax + 240a^2 x^2$$



Question 5 continued

expand using quick method

$$f(x) = \left(3 + \frac{1}{x}\right)^2 (2 + ax)^6$$

expansion estimated in (a).

$$f(x) = \left(9 + \frac{6}{x} + \frac{1}{x^2}\right) (64 + 192ax + 240a^2x^2)$$

After multiplying, the constant terms include all terms where :

- no x is present in either multiplied value :

$$9 \times 64 = 576$$

- the x value cancels out :

$$\frac{6}{x} \times 192ax = 1152a$$

- the x^2 value cancels out :

$$\frac{1}{x^2} \times 240a^2x^2 = 240a^2$$

$$\text{so : } \cancel{576} + 1152a + 240a^2 = \cancel{576}$$

$$a(240a + 1152) = 0$$

$[a = 0]$ or
reject, a is
non-zero!

$$240a + 1152 = 0$$

$$a = \frac{-1152}{240} = \frac{-24}{5}$$

FINAL ANSWER!

(Total for Question 5 is 7 marks)



6.

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

Using the laws of logarithms, solve

$$\log_4(12 - 2x) = 2 + 2\log_4(x + 1)$$

(5)

Recall rules of logarithms :

$$\log_a(b) + \log_a(c) = \log_a(b \times c)$$

$$\log_a(b) - \log_a(c) = \log_a\left(\frac{b}{c}\right)$$

$$\log_a b^c = c \log_a b$$

$$\log_a b = c \rightarrow a^c = b$$

$$\log_4(12 - 2x) = 2 + 2\log_4(x + 1)$$

$$\log_4(12 - 2x) - 2\log_4(x + 1) = 2$$

$$\log_4(12 - 2x) - \log_4(x + 1)^2 = 2$$

$$\log_4\left(\frac{12 - 2x}{(x + 1)^2}\right) = 2$$

$$(x + 1)^2 \times \left(\frac{12 - 2x}{(x + 1)^2}\right) = (4^2) \times (x + 1)^2$$

$$12 - 2x = 16(x^2 + 2x + 1)$$

$$12 - 2x = 16x^2 + 32x + 16$$

$$\frac{16x^2}{2} + \frac{34x}{2} + \frac{4}{2} = 0$$

$$8x^2 + 17x + 2 = 0$$

$$(8x + 1)(x + 2) = 0$$

FINAL
ANSWER

$$x = -1/8 \text{ or } x = -2$$

REJECT ;
 $\log_4(-2 + 1)$
 $= \log_4(-1)$, log
of -ve value has
no real solution!

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 6 continued

ALTERNATIVE SOLUTION :

$$\log_4(12-2x) = 2 + 2\log_4(x+1)$$

$$2\log_4(x+1) - \log_4(12-2x) = -2$$

$$\log_4(x+1)^2 - \log_4(12-2x) = -2$$

$$\log_4\left(\frac{(x+1)^2}{12-2x}\right) = -2$$

$$(12-2x) \times \left(\frac{(x+1)^2}{12-2x}\right) = (4)^{-2} \times (12-2x)$$

$$x^2 + 2x + 1 = \frac{1}{16}(12-2x)$$

$$x^2 + 2x + 1 = \frac{12}{16} - \frac{2}{16}x$$

$$\left(x^2 + \frac{17}{8}x + \frac{1}{4} = 0\right) \times 8$$

$$8x^2 + 17x + 2 = 0$$

$$(8x + 1)(x + 2) = 0$$

$$x = -1/8 \text{ or } x = -2$$

**FINAL
ANSWER**

REJECT ;
 $\log_4(-2+1)$
 $= \log_4(-1)$, log
of -ve value has
no real solution!

(Total for Question 6 is 5 marks)



7. Jem pays money into a savings scheme, A , over a period of 300 months.

Jem pays £20 into scheme A in month 1, £20.50 in month 2, £21 in month 3 and so on, so that the amounts Jem pays each month form an arithmetic sequence.

- (a) Show that Jem pays £69.50 into scheme A in month 100

partial sum S_n

$n=300$

(1)

- (b) Find the total amount that Jem pays into scheme A over the period of 300 months.

(2)

Kim pays money into a different savings scheme, B , over the same period of 300 months.

In a model, the amounts Kim pays into scheme B increase by the same percentage each month, so that the amounts Kim pays each month form a geometric sequence.

Given that Kim pays

- £20 into scheme B in month 1
 - £250 into scheme B in month 300
- (c) use the model to calculate, to the nearest £10, the difference between the total amount paid into scheme A and the total amount paid into scheme B over the period of 300 months.

(3)

(a) $a_1, a_2, a_3, \dots, a_{100}$

£20, £20.50, £21, ..., £??

+0.50 +0.5

For arithmetic series :

$$a_n = a + d(n-1)$$

$$a_{100} = 20 + 0.5(100-1) = £69.50$$

(b) use partial sum formula :

$$S_n = \frac{n}{2} [2a + d(n-1)]$$

$$= \frac{300}{2} [2(20) + 0.5(300-1)] = £28\,425$$



Question 7 continued

OR: use partial sum formula:

$$S_n = \frac{n}{2} [a_1 + a_n]$$

$$a_{300} = 20 + 0.5(300-1) = 169.5$$

$$S_{300} = \frac{300}{2} [20 + 169.5] = £28\,425$$

(c) ① Find r

$$a_n = ar^{n-1}$$

$$250 = 20r^{300-1}$$

use the exact value for calculations!

$$299 \sqrt[299]{r^{299}} = 299 \sqrt[299]{\frac{25}{2}} \rightarrow r = 299 \sqrt[299]{\frac{25}{2}}$$

② use partial sum formula:

$$S_n = \frac{a_1(1-r^n)}{(1-r)}$$

$$S_{300} = \frac{20 \left(1 - \left(299 \sqrt[299]{\frac{25}{2}} \right)^{300} \right)}{\left(1 - 299 \sqrt[299]{\frac{25}{2}} \right)}$$

$$S_{300} = 27\,362.948...$$

$$= £27\,362$$

$$\text{difference} = £28\,425 - £27\,362 = 1063$$

$$= £1060$$

(nearest ten)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 7 is 6 marks)



8.

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.




Figure 1

Figure 1 shows a sketch of part of the curve C_1 with equation

$$y = x^2 + 3 \quad x > 0$$

and part of the curve C_2 with equation

$$y = 13 - \frac{9}{x^2} \quad x > 0$$

The curves C_1 and C_2 intersect at the points P and Q as shown in Figure 1.(a) Use algebra to find the x coordinate of P and the x coordinate of Q .

(4)

The finite region R , shown shaded in Figure 1, is bounded by C_1 and C_2 (b) Use algebraic integration to find the exact area of R .

(4)

(a) P and Q lie on C_1 and C_2 , so:

$$\text{at } x_P \text{ and } x_Q, \quad y_{C_1} = y_{C_2}$$

$$x^2 \times (x^2 + 3) = \left(13 - \frac{9}{x^2}\right) \times x^2$$

$$x^4 + 3x^2 = 13x^2 - 9$$



Question 8 continued

$$x^4 - 10x^2 + 9 = 0$$

$$(x^2)^2 - 10(x^2)^1 + 9 = 0 \rightarrow \text{can be factorised}$$

$$(x^2 - 1)(x^2 - 9) = 0$$

like a quadratic equation!

$$x^2 - 1 = 0 \quad \text{or} \quad x^2 - 9 = 0$$

$$x = 1 \quad \text{or} \quad x = 3$$

$$x_P = 1$$

$$x_Q = 3$$

$$(b) \quad A_R = A_{C2} - A_{C1}$$

$$A_R = \int_1^3 13 - \frac{9}{x^2} dx - \int_1^3 x^2 + 3 dx$$

COMBINE USING RULE:

$$\int f(x) dx \pm \int g(x) dx = \int f(x) \pm g(x) dx$$

$$A_R = \int_1^3 13 - \frac{9}{x^2} - (x^2 + 3) dx$$

$$A_R = \int_1^3 13 - 9x^{-2} - x^2 - 3 dx$$

$$A_R = \int_1^3 10 - 9x^{-2} - x^2 dx$$

REMEMBER!

$$\int x^n dx = \frac{x^{n+1}}{n+1}$$

Question 8 continued

$$A_R = \left[\frac{10x^{0+1}}{0+1} - \frac{9x^{-2+1}}{-2+1} - \frac{x^{2+1}}{2+1} \right]_1^3$$

$$A_R = \left[10x + \frac{9}{x} - \frac{x^3}{3} \right]_1^3$$

$$A_R = \left[10(3) + \frac{9}{3} - \frac{3^3}{3} \right] - \left[10(1) + \frac{9}{1} - \frac{1^3}{3} \right]$$

$$A_R = (30 + 3 - 9) - (10 + 9 - 1/3)$$

$$A_R = 24 - \frac{56}{3}$$

$$A_R = \frac{16}{3}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



9.

In this question you must show detailed reasoning.

Solutions relying entirely on calculator technology are not acceptable.

(a) Show that the equation

$$2 \tan \theta = 3 \cos \theta$$

can be written as

$$3 \sin^2 \theta + 2 \sin \theta - 3 = 0$$

(3)

(b) Hence solve, for $-\pi < x < \pi$, the equation

$$2 \tan \left(2x + \frac{\pi}{3} \right) = 3 \cos \left(2x + \frac{\pi}{3} \right)$$

giving your answers to 3 significant figures.

(4)

$$(a) \quad 2 \tan \theta = 3 \cos \theta \quad \tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\cos \theta \times \left(2 \frac{\sin \theta}{\cos \theta} \right) = (3 \cos \theta) \times \cos \theta$$

$$2 \sin \theta = 3 \cos^2 \theta$$

Use identity: $\sin^2 \theta + \cos^2 \theta = 1$
 $\cos^2 \theta = 1 - \sin^2 \theta$

$$2 \sin \theta = 3(1 - \sin^2 \theta)$$

$$2 \sin \theta = 3 - 3 \sin^2 \theta$$

$$\therefore 3 \sin^2 \theta + 2 \sin \theta - 3 = 0$$

$$(b) \quad 3 \sin^2 \left(2x + \frac{\pi}{3} \right) + 2 \sin \left(2x + \frac{\pi}{3} \right) - 3 = 0$$

Let $\sin^2 \left(2x + \frac{\pi}{3} \right) = x$

$$3x^2 + 2x - 3 = 0 \rightarrow \text{FACTORISE!}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 9 continued

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-2 \pm \sqrt{2^2 - 4(3)(-3)}}{2(3)} \rightarrow x = \frac{-1 \pm \sqrt{10}}{3}$$

$$\sin\left(2x + \frac{\pi}{3}\right) = \frac{-1 - \sqrt{10}}{3} \quad \times \quad \text{REJECT; range of } y = \sin \theta \text{ is } -1 < \theta < 1$$

$$\sin\left(2x + \frac{\pi}{3}\right) = \frac{-1 + \sqrt{10}}{3} \rightarrow = 0.72... \text{ so accept!}$$


NOW, SOLVE:

$$\sin\left(2x + \frac{\pi}{3}\right) = \frac{-1 + \sqrt{10}}{3}, \quad \left[2(-\pi) + \frac{\pi}{3} < 2x + \frac{\pi}{3} < 2(\pi) + \frac{\pi}{3} \right] \quad \text{domain!}$$

ON CALCULATOR:

$$\left(2x + \frac{\pi}{3}\right) = \sin^{-1}\left(\frac{-1 + \sqrt{10}}{3}\right) = 0.8049$$

ON UNIT CIRCLE:



$$\text{SOL 1: } 2x + \frac{\pi}{3} = 0.8049$$

$$\text{SOL 2: } 2x + \frac{\pi}{3} = \pi - 0.8049 = 2.337$$

these two solutions are only within the domain of $0 < 2x + \frac{\pi}{3} < 2\pi$.

(Total for Question 9 is 7 marks)

Question 9 continued

Since our domain is actually :

$$-\frac{5\pi}{3} < 2x + \frac{\pi}{3} < \frac{7\pi}{3}$$

$$\text{SOL 3: } 2x + \frac{\pi}{3} = 2.337 - 2\pi \\ = -3.946$$

$$\text{SOL 4: } 2x + \frac{\pi}{3} = 0.8049 + 2\pi \\ = 7.088$$

$$2x + \frac{\pi}{3} = 0.8049, -3.949, 2.337, 7.088$$

Subtract $\frac{\pi}{3}$, divide by 2;

$$x = -0.121, -2.50, 0.645, 3.02 \quad (3 \text{ s.f.})$$

10. The circle C has equation

$$x^2 + y^2 + 4x - 30y + 209 = 0$$

(a) Find

- (i) the coordinates of the centre of C ,
- (ii) the exact value of the radius of C .

(3)

The line L has equation $y = mx + 1$, where m is a constant.

Given that L is the tangent to C at the point P ,

(b) show that

$$2m^2 - 7m - 22 = 0$$

(3)

(c) Hence find the possible pairs of coordinates of P .

(4)

(a) Standard form of equation of a circle:

$$(x - x_{\text{centre}})^2 + (y - y_{\text{centre}})^2 = r^2$$

↳ rewrite given eq. into standard form :

$$x^2 + 4x + y^2 - 30y + 209 = 0$$

(1) complete the square for terms with x :

$$x^2 + 4x + y^2 - 30y = -209$$

↳ add $\left(\frac{b}{2}\right)^2$ to both sides :

$$x^2 + 4x + \left(\frac{(-4)}{2}\right)^2 + y^2 - 30y = -209 + \left(\frac{(-4)}{2}\right)^2$$

$$x^2 + 4x + 4 + y^2 - 30y = -209 + 4$$

↳ factorise



Question 10 continued

$$(x+2)^2 + y^2 - 30y = -205$$

(3) complete the square for terms with y :

$$(x+2)^2 + y^2 - 30y = -205$$

↳ add $\left(\frac{b}{2}\right)^2$ to both sides :

$$(x+2)^2 + y^2 - 30y + \left(\frac{-30}{2}\right)^2 = -205 + \left(\frac{-30}{2}\right)^2$$

$$(x+2)^2 + y^2 - 30y + 225 = -205 + 225$$

↳ factorise

$$(x+2)^2 + (y-15)^2 = 20$$

$$\text{so: } (x+2)^2 + (y-15)^2 = 20$$

(i) coords of centre : $(-2, 15)$ (ii) radius $\rightarrow r^2 = 20$, $r = \sqrt{20}$ (b) At point P, $y_c = y_L$ ↳ substitute $y = mx+1$ into y_c :

$$(x+2)^2 + (mx+1-15)^2 = 20$$

$$(x+2)^2 + (mx-14)^2 = 20$$

$$x^2 + 4x + 4 + m^2x^2 - 28mx + 196 = 20$$

$$x^2 + m^2x^2 + 4x - 28mx + 180 = 0$$

Question 10 continued

$$(1 + m^2)x^2 + (4 - 28m)x + 180 = 0$$

a
 b
 c

Since line L and circle C only touch at **ONE POINT**, we can use the discriminant to form an equation with m alone.

REMEMBER :

discriminant

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

If discriminant > 0 , then eq. has two solutions
 If discriminant $= 0$, then eq. has **one solution**
 If discriminant < 0 , then eq. has no solutions

$$b^2 - 4ac = 0$$

$$(4 - 28m)^2 - 4(1 + m^2)(180) = 0$$

$$16 - 224m + 784m^2 - 4(180 + 180m^2) = 0$$

$$16 - 224m + 784m^2 - 720 - 720m^2 = 0$$

$$\frac{64m^2}{32} - \frac{224m}{32} - \frac{704}{32} = 0$$

$$\therefore 2m^2 - 7m - 22 = 0$$

(c) $2m^2 - 7m - 22 = 0 \rightarrow$ **FACTORISE!**

$$(2m - 11)(m + 2) = 0$$

$$m = 11/2 \quad \text{or} \quad m = -2$$



Question 10 continued

① WHEN $m = \frac{11}{2}$:

$$x^2 + \left(\frac{11}{2}\right)^2 x^2 + 4x - 28\left(\frac{11}{2}\right)x + 180 = 0$$

$$x^2 + \frac{121}{4}x^2 + 4x - 154x + 180 = 0$$

$$4x \left(\frac{125}{4}x^2 - 150x + 180 \right) = (0) \times 4$$

$$125x^2 - 600x + 720 = 0 \rightarrow \text{FACTORISE}$$

$$(5x - 12)^2 = 0$$

$$x = \frac{12}{5}$$

$$y = \frac{11}{2} \left(\frac{12}{5} \right) + 1 = \frac{71}{5}$$

② WHEN $m = -2$:

$$x^2 + (-2)^2 x^2 + 4x - 28(-2)x + 180 = 0$$

$$x^2 + 4x^2 + 4x + 56x + 180 = 0$$

$$\frac{5x^2}{5} + \frac{60x}{5} + \frac{180}{5} = 0$$

$$x^2 + 12x + 36 = 0 \rightarrow \text{FACTORISE}$$

$$(x + 6)^2 = 0$$

$$x = -6$$

$$y = \frac{11}{2}(-6) + 1 = 13$$

possible coordinates: $\left(\frac{12}{5}, \frac{71}{5} \right)$ or $(-6, 13)$

(Total for Question 10 is 10 marks)

11. (i) Prove by counter example that the statement

"If n is a prime number then $3^n + 2$ is also a prime number."

is false.

(2)

(ii) Use proof by exhaustion to prove that if m is an integer that is **not** divisible by 3, then

$$m^2 - 1$$

is divisible by 3

(4)

(i) Give any example that disproves the statement :

$$\text{ex: } n = 5 : 3^5 + 2 = 245 \\ = 5 \times 7 \times 7 \quad \text{NOT PRIME}$$

$$n = 7 : 3^7 + 2 = 2189 \\ = 11 \times 199 \quad \text{NOT PRIME}$$

$$n = 11 : 3^{11} + 2 = 177149 \\ = 7 \times 25307 \quad \text{NOT PRIME}$$

(ii) If m is divisible by 3, $m = 3k$

If m is NOT divisible by 3 :

$$m = 3k + 1$$

$$m = 3k + 2$$

WHEN $m = 3k + 1$:

$$\begin{aligned} & (3k+1)^2 - 1 \\ &= 9k^2 + 6k + 1 - 1 \\ &= 9k^2 + 6k \\ &= 3(3k^2 + 2k) \end{aligned}$$

WHEN $m = 3k + 2$

$$\begin{aligned} & (3k+2)^2 - 1 \\ &= 9k^2 + 12k + 4 - 1 \\ &= 9k^2 + 12k + 3 \\ &= 3(3k^2 + 4k + 1) \end{aligned}$$



Question 11 continued

$3(3k^2 + 2k)$ and $3(3k^2 + 4 + 1)$ are both multiples of 3, so :

$m^2 - 1$ must be divisible by 3 when m is not divisible by 3.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

