



Pearson  
Edexcel

Mark Scheme (FINAL)

October 2022

Pearson Edexcel International Advanced Level

In Pure Mathematics 4 (WMA14) Paper 01

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

### **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

October 2022

Question Paper Log Number P72068A

Publications Code WMA14\_01\_2210\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2022

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

## General Principles for Core Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

### **Method mark for solving 3 term quadratic:**

#### **1. Factorisation**

$(x^2 + bx + c) = (x + p)(x + q)$ , where  $|pq| = |c|$ , leading to  $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$ , where  $|pq| = |c|$  and  $|mn| = |a|$ , leading to  $x = \dots$

#### **2. Formula**

Attempt to use the correct formula (with values for a, b and c).

#### **3. Completing the square**

Solving  $x^2 + bx + c = 0$ :  $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0$ ,  $q \neq 0$ , leading to  $x = \dots$

### **Method marks for differentiation and integration:**

#### **1. Differentiation**

Power of at least one term decreased by 1. ( $x^n \rightarrow x^{n-1}$ )

#### **2. Integration**

Power of at least one term increased by 1. ( $x^n \rightarrow x^{n+1}$ )

### **Use of a formula**

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small errors in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

### **Exact answers**

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Question Number	Scheme	Marks
1	<p>Attempts to get <math>t</math> in terms of <math>x</math> or <math>y</math>. E.g. <math>x = \frac{t}{t-3} \Rightarrow t = \frac{3x}{x-1}</math></p> <p>Forms a Cartesian equation and makes progress to making <math>y</math> the subject</p> <p>E.g. <math>y = \frac{1}{t} + 2 \Rightarrow y = \frac{x-1}{3x} + 2</math></p> <p><math>\Rightarrow y = \frac{7x-1}{3x}</math></p>	<p>M1</p> <p>dM1</p> <p>A1</p> <p>(3)</p> <p>(3 marks)</p>

M1: For an attempt to get  $t$  or  $\frac{1}{t}$  in terms of  $x$  or  $y$ .

Condone poor attempts here. It is the intention that is important

dM1: For a complete attempt to form an equation of the form  $y = f(x)$

Look for

- getting  $t$  or  $\frac{1}{t}$  in terms of  $x$  or  $y$
- using this to form a Cartesian equation and makes some progress towards making  $y$  the subject.

E.g.I.  $x = \frac{t}{t-3} \Rightarrow t = \frac{3x}{x-1}$  followed by  $y = \frac{1}{t} + 2 \Rightarrow y = \frac{1}{\frac{3x}{x-1}} + 2$

E.g.II.  $y = \frac{1}{t} + 2 \Rightarrow t = \frac{1}{y-2}$  followed by  $x = \frac{1}{\frac{1}{y-2}-3} \Rightarrow x = \frac{1}{1-3(y-2)} \Rightarrow (7-3y)x = 1$

Condone slips, for example in the signs and miscopies/misreads, but the overall mechanics of the attempt should be sound.

In E.g.II. an attempt must be made to make progress towards  $y = f(x)$ , so allow for example when an intermediate line such as  $g(y) \times h(x) = b$  or  $g(y) = h(x)$  where  $g(y)$  is a linear expression in  $y$

A1: Correct equation  $y = \frac{7x-1}{3x}$  but allow  $y = \frac{-1+7x}{3x}$

Alt Method:

M1: Substitutes both  $y = \frac{1}{t} + 2$  and  $x = \frac{t}{t-3}$  into  $y = \frac{ax-1}{bx}$  and makes progress to a point in which sides can be compared

$$\frac{1}{t} + 2 = \frac{a \times \frac{t}{t-3} - 1}{b \times \frac{t}{t-3}} = \left( \frac{a-1}{b} \right) + \frac{3}{bt} \quad \text{or} \quad b + 2bt = (a-1)t + 3$$

dM1: Compares terms to set up and solve simultaneous equations in both  $a$  and  $b$ .

FYI  $\frac{3}{b} = 1, \frac{a-1}{b} = 2$  or  $b = 3, 2b = a-1$

A1:  $y = \frac{7x-1}{3x}$  NOT just for correct values of  $a$  and  $b$

Question Number	Scheme	Marks
2 (a)	$\frac{3x}{(2x-1)(x-2)} \equiv \frac{A}{2x-1} + \frac{B}{x-2} \Rightarrow$ Values for $A$ and $B$	M1
	One correct value, either $A = -1$ or $B = 2$ Correct PF form $\frac{-1}{2x-1} + \frac{2}{x-2}$	A1 A1
(b)	$\int \frac{-1}{2x-1} + \frac{2}{x-2} dx = -\frac{1}{2} \ln(2x-1) + 2 \ln(x-2) + (c)$	M1, A1ft
	$\int_5^{25} \frac{3x}{(2x-1)(x-2)} dx = \left[ -\frac{1}{2} \ln(2x-1) + 2 \ln(x-2) \right]_5^{25}$ $= \left( -\frac{1}{2} \ln 49 + 2 \ln 23 \right) - \left( -\frac{1}{2} \ln 9 + 2 \ln 3 \right)$ $\ln \frac{529}{21}$	dM1 A1
		(3) (4) (7 marks)

(a)

M1: Attempts a correct PF form leading to values for  $A$  and  $B$ .

Condone  $\frac{3x}{(2x-1)(x-2)} \equiv C + \frac{A}{2x-1} + \frac{B}{x-2} \Rightarrow$  Values for at least  $A$  and  $B$

A1: One correct value for  $A$  or  $B$  or one correct term

A1: Correct PF form  $\frac{-1}{2x-1} + \frac{2}{x-2}$ .

This is not just for correct values for  $A$  and  $B$  but it may be awarded from work seen in (b)

(b)

M1:  $\int \frac{A}{2x-1} + \frac{B}{x-2} dx = C \ln(2x-1) + D \ln(x-2)$  o.e. Condone missing brackets here

A1ft: Correct ft on their values for  $A$  and  $B$   $\int \frac{A}{2x-1} + \frac{B}{x-2} dx = \frac{A}{2} \ln(2x-1) + B \ln(x-2) + (c)$

Do not condone missing brackets but the brackets can be implied by subsequent working, E.g.  $\ln 49$   
dM1: Attempts to substitute the limits 5 and 25 (no misreads on the limits but condone slips, so don't accept a consistent lower limit of 0 for instance) into an expression of the correct form (with brackets which may be implied) and subtracts, either way around. Condone attempts where the substitutions were made into incorrectly combined  $\ln$  terms.

A1: cso  $\ln \frac{529}{21}$  Also allow in mixed number form  $\ln 25 \frac{4}{21}$

Question Number	Scheme	Marks
3 (a)	Attempts $(2\mathbf{i} - 3\mathbf{j} + 4\mathbf{k}) - (8\mathbf{i} - 5\mathbf{j} + 3\mathbf{k})$	M1
	$\overrightarrow{RQ} = -6\mathbf{i} + 2\mathbf{j} + \mathbf{k}$	A1
(b)	Attempts $\overrightarrow{PQ} \cdot \overrightarrow{RQ} = 2 \times -6 + -3 \times 2 + 4 \times 1$	M1
	Full attempt to find $\cos PQR$ E.g. $2 \times -6 + -3 \times 2 + 4 \times 1 = \sqrt{29} \sqrt{41} \cos PQR$	dM1
	Angle $PQR = 114^\circ$	A1
		(2)
		(3)
		(5 marks)

(a)

M1: Attempts to subtract vectors  $\overrightarrow{PQ}$  and  $\overrightarrow{PR}$  either way around. Look for  $\overrightarrow{PQ} - \overrightarrow{PR}$  or  $\overrightarrow{PR} - \overrightarrow{PQ}$ .  
If a method is not shown it can be implied by two correct components of  $\pm 6\mathbf{i} \pm 2\mathbf{j} \pm \mathbf{k}$   
Note that an attempt such as  $\overrightarrow{PR} - \overrightarrow{QP}$  is M0

A1:  $\overrightarrow{RQ} = -6\mathbf{i} + 2\mathbf{j} + \mathbf{k}$  o.e. such as  $\begin{pmatrix} -6 \\ 2 \\ 1 \end{pmatrix}$  Do not accept coordinates or indeed  $\begin{pmatrix} -6\mathbf{i} \\ 2\mathbf{j} \\ \mathbf{k} \end{pmatrix}$

(b)

M1: Attempts scalar product of  $\overrightarrow{PQ}$  and their  $\overrightarrow{RQ}$ . Look for an attempt at multiplying together the components and adding. There will be some confusion over direction so allow for sight of  $(\pm 2 \times \pm 6) + (\pm 3 \times \pm 2) + (\pm 4 \times \pm 1)$

This cannot be scored if they attempt a scalar product of  $\overrightarrow{PQ}$  and  $\overrightarrow{PR}$  for instance.

dM1: Full attempt to find  $\cos PQR$  using  $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta$  using vectors  $\pm \overrightarrow{PQ}$  and their  $\pm \overrightarrow{RQ}$ .

There must be an attempt at both moduli with at least one correct (which may be unsimplified) but you should fit on their  $\overrightarrow{RQ}$ . Don't be concerned whether the angle is acute or obtuse.

A1: Angle  $PQR = \text{awrt } 114^\circ$  ISW after sight of this, e.g followed by  $66^\circ$

Alt (b)

M1: Attempts all three lengths or all three lengths<sup>2</sup> using Pythagoras' Theorem. Look for an attempt to square and add with at least one modulus or modulus<sup>2</sup> correct.

dM1: Attempts to use the cosine rule with the lengths in the correct positions in order to find angle  $PQR$

$$\text{Look for } \cos PQR = \frac{PQ^2 + QR^2 - PR^2}{2 \times PQ \times QR}$$

There are more round about methods including finding other angles first and then using the sine rule but this method mark can only be awarded when an attempt is made at angle  $PQR$

A1: Angle  $PQR = \text{awrt } 114^\circ$

It must be found using "correct" vectors, e.g.  $\pm(-6\mathbf{i} + 2\mathbf{j} + \mathbf{k})$  and  $\pm(2\mathbf{i} - 3\mathbf{j} + 4\mathbf{k})$

If, for example  $\overrightarrow{RQ}$  is incorrect, e.g.  $6\mathbf{i} - 2\mathbf{j} + \mathbf{k}$ , A0 will be awarded even if  $114^\circ$  is stated

Question Number	Scheme	Marks
<b>4(a)</b>	$\frac{1}{\sqrt{4-x^2}} = (4-x^2)^{-\frac{1}{2}} = 4^{-\frac{1}{2}}(1-\dots)$	B1
	$\left(1-\frac{1}{4}x^2\right)^{-\frac{1}{2}} = 1 + \left(-\frac{1}{2}\right)\left(-\frac{1}{4}x^2\right) + \frac{\left(-\frac{1}{2}\right)\times\left(-\frac{3}{2}\right)}{2}\left(-\frac{1}{4}x^2\right)^2 + \frac{\left(-\frac{1}{2}\right)\times\left(-\frac{3}{2}\right)\times\left(-\frac{5}{2}\right)}{3!}\left(-\frac{1}{4}x^2\right)^3$	M1, A1
	$\frac{1}{\sqrt{4-x^2}} = \frac{1}{2} + \frac{1}{16}x^2 + \frac{3}{256}x^4 + \frac{5}{2048}x^6$	A1, A1
<b>(b)</b>	$ x  < 2$	B1
<b>(c)</b>	Substitutes an appropriate value of $x$ in both sides with LHS in terms of $\sqrt{3}$ E.g. with $x = 1$ $\sqrt{3} = \frac{2048}{1181}$ or $\frac{3543}{2048}$	M1 A1
		(5) (1) (2) <b>(8 marks)</b>

(a) Note that the first mark in (a) is M1 on open. We are now marking it as B1

B1: Correct constant term/ factor.

Accept as  $4^{-\frac{1}{2}}$  or  $\frac{1}{2}$  as the constant term or as the factor with the bracket starting (1...

M1: Correct attempt at the third or the fourth term of binomial expansion of form  $\left(1+ax^2\right)^{-\frac{1}{2}}$

Look for a correct binomial coefficient with a correct power of  $x$

E.g.  $\frac{\left(-\frac{1}{2}\right)\times\left(-\frac{1}{2}-1\right)}{2}\left(\pm ax^2\right)^2$  or  $\frac{\left(-\frac{1}{2}\right)\times\left(-\frac{1}{2}-1\right)\times\left(-\frac{1}{2}-2\right)}{3!}\left(\pm ax^2\right)^3$  where  $a$  could be 1

A1: Correct unsimplified

For example  $\left(1-\frac{1}{4}x^2\right)^{-\frac{1}{2}} = 1 + \left(-\frac{1}{2}\right)\left(-\frac{1}{4}x^2\right) + \frac{\left(-\frac{1}{2}\right)\times\left(-\frac{1}{2}-1\right)}{2}\left(-\frac{1}{4}x^2\right)^2 + \frac{\left(-\frac{1}{2}\right)\times\left(-\frac{1}{2}-1\right)\times\left(-\frac{1}{2}-2\right)}{3!}\left(-\frac{1}{4}x^2\right)^3$

A1: Two correct and simplified terms of  $\frac{1}{2} + \frac{1}{16}x^2 + \frac{3}{256}x^4 + \frac{5}{2048}x^6$

A1:  $\frac{1}{2} + \frac{1}{16}x^2 + \frac{3}{256}x^4 + \frac{5}{2048}x^6$ . Accept as a list, ignore any terms with indices greater than these.

**(b)**

**B1:** States a correct range. E.g  $|x| < 2$  or  $-2 < x < 2$  o.e.

**(c) For example: See \*\***

**M1:** Substitutes an exact value of  $x$  into both sides of their expansion that enables  $\sqrt{3}$  to be found .



$$\text{E.g. } x=1 \quad \frac{1}{\sqrt{4-1^2}} = \frac{1}{2} + \frac{1}{16} \times 1^2 + \frac{3}{256} \times 1^4 + \frac{5}{2048} \times 1^6$$

Note that  $x = -1$  and  $x^2 = 1$  all work with the same results

Alternatively correctly solves  $\frac{1}{\sqrt{4-x^2}} = \sqrt{3}$  o.e. and substitutes this into their expansion

**A1:** With  $x=1 \quad \sqrt{3} = \frac{2048}{1181}$  or  $\frac{3543}{2048}$ . This can only be scored from a correct expansion

\*\*There are lots of values of  $x$  that work, all more difficult than the one above.

$$\text{E.g. } x^2 = \frac{8}{3} \Rightarrow \frac{1}{\sqrt{4-x^2}} = \frac{\sqrt{3}}{2} \quad \text{and} \quad x^2 = \frac{11}{3} \Rightarrow \frac{1}{\sqrt{4-x^2}} = \sqrt{3}$$

In both these cases the fractions become much more difficult to find.

$$\text{For } x^2 = \frac{8}{3} \Rightarrow \sqrt{3} = \frac{43}{27} \quad \text{and for } x^2 = \frac{11}{3} \Rightarrow \sqrt{3} = \frac{55687}{55296}$$

Alt I (a) By direct expansion

$$\frac{1}{\sqrt{4-x^2}} = \left(4-x^2\right)^{-\frac{1}{2}} = 4^{-\frac{1}{2}} + \left(-\frac{1}{2}\right) \times 4^{-\frac{3}{2}} \left(-x^2\right)^1 + \frac{\left(-\frac{1}{2}\right) \times \left(-\frac{3}{2}\right)}{2} \times 4^{-\frac{5}{2}} \left(-x^2\right)^2 + \frac{\left(-\frac{1}{2}\right) \times \left(-\frac{3}{2}\right) \times \left(-\frac{5}{2}\right)}{3!} \times 4^{-\frac{7}{2}} \left(-x^2\right)^3$$

B1: For  $4^{-\frac{1}{2}} +$

M1: A correct attempt at the third or fourth terms condoning sign slips on the  $-x^2$

A1: Correct and unsimplified expansion. See above

A1: Two correct and simplified terms of  $\frac{1}{2} + \frac{1}{16}x^2 + \frac{3}{256}x^4 + \frac{5}{2048}x^6$

A1:  $\frac{1}{2} + \frac{1}{16}x^2 + \frac{3}{256}x^4 + \frac{5}{2048}x^6$ . Accept as a list

Alt II (a) By difference of two squares

$$\frac{1}{\sqrt{4-x^2}} = (2-x)^{-\frac{1}{2}} \times (2+x)^{-\frac{1}{2}} = 4^{-\frac{1}{2}} + \dots$$

B1: For  $4^{-\frac{1}{2}} +$

M1: For correctly writing  $\frac{1}{\sqrt{4-x^2}} = (2-x)^{-\frac{1}{2}} \times (2+x)^{-\frac{1}{2}} = \dots \left(1-\frac{x}{2}\right)^{-\frac{1}{2}} \times \left(1+\frac{x}{2}\right)^{-\frac{1}{2}} =$

with a correct attempt at the third or fourth term in either expansion, followed by an attempt to combine both the expansions.

A1: One correct term in  $x^2$ ,  $x^4$  or  $x^6$

A1: Two correct and simplified terms of  $\frac{1}{2} + \frac{1}{16}x^2 + \frac{3}{256}x^4 + \frac{5}{2048}x^6$  with no terms in  $x$ ,  $x^3$  or  $x^5$

A1: Fully correct

Question Number	Scheme	Marks
5.	<p>States or implies Volume = <math>\int_{\frac{1}{\sqrt{2}}}^k \pi \left( \frac{12\sqrt{x}}{(2x^2+3)^{1.5}} \right)^2 dx</math></p> <p>Attempts <math>\int \left( \frac{12\sqrt{x}}{(2x^2+3)^{1.5}} \right)^2 dx = \int \frac{144x}{(2x^2+3)^3} dx = -18(2x^2+3)^{-2}</math></p> <p>Sets <math>-\frac{18\pi}{(2k^2+3)^2} + \frac{18\pi}{(2 \times \frac{1}{2} + 3)^2} = \frac{713}{648}\pi</math></p> <p><math>\Rightarrow -\frac{18}{(2k^2+3)^2} + \frac{9}{8} = \frac{713}{648} \Rightarrow (2k^2+3)^2 = 729 \Rightarrow k^2 = \dots</math></p> <p><math>k = 2\sqrt{3}</math></p>	<p>B1</p> <p>M1A1</p> <p>M1</p> <p>ddM1</p> <p>A1</p> <p><b>(6 marks)</b></p>

B1: States or implies Volume =  $\int_{\frac{1}{\sqrt{2}}}^k \pi \left( \frac{12\sqrt{x}}{(2x^2+3)^{1.5}} \right)^2 dx$  o.e. including limits and  $\pi$ .

You can condone a missing dx and the limits the wrong way around.

You may not see the limits until the integration has been attempted which is fine.

M1: Attempts  $\int \left( \frac{\sqrt{x}}{(2x^2+3)^{1.5}} \right)^2 dx$  to achieve  $A(2x^2+3)^{-2}$  o.e.

With substitution of  $u = 2x^2 + 3$  look for  $Au^{-2}$

A1: Correct integration of  $\int \left( \frac{12\sqrt{x}}{(2x^2+3)^{1.5}} \right)^2 dx$  to  $-18(2x^2+3)^{-2}$  or  $-18u^{-2}$  which may be left unsimplified.

Note that this can be scored with or without the  $\pi$  and may even have an incorrect coefficient of  $2\pi$

M1: Uses the limits (subtracts either way around) within their attempted integration of  $y^2$  NOT  $y$  and sets

equal to  $\frac{713}{648}\pi$ . Condone poor attempts at integrating the  $y^2$  here.

Note that if they change limits to  $u$  where  $u = 2x^2 + 3$  look for limits of  $2k^2 + 3$  and 4

ddM1: Dependent upon both previous M's. It is for proceeding to

either  $(2k^2+3)^{\pm 2} = B$  where  $B > 0$  leading to a value for  $k^2$  or  $k$

or setting up a quadratic equation in  $k^2$  leading to a value for  $k^2$  or  $k$

$$\text{FYI } k^4 + 3k^2 - 180 = 0 \Rightarrow (k^2 - 12)(k^2 + 15) = 0 \Rightarrow k^2 = 12$$

A1:  $k = 2\sqrt{3}$  or exact equivalent such as  $\sqrt{12}$  Condone if  $k = \pm 2\sqrt{3}$  o.e.

Question Number	Scheme	Marks
<b>6 (a)</b>	For correct parameter $t = \frac{\pi}{4}$ or $x$ coordinate at $P$ $x = 4$  $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{-4 \sin 2t}{3 \sec^2 t}$ Equation of tangent $y - 0 = -\frac{2}{3}(x - 4) \Rightarrow y = -\frac{2}{3}x + \frac{8}{3}$	B1  M1 A1  dM1 A1  (5)
<b>(b)</b>	$k = 1 - \sqrt{3}$	B1  (1)
<b>(c)</b>	$-1 \leq f \leq 2$	M1 A1  (2)
		<b>(8 marks)</b>

(a)

B1: Correct parameter or  $x$  coordinate at  $P$ .

The correct  $x$  coordinate  $x = 4$  (can even be scored following  $x = 45^\circ$ )

M1: Attempts to find  $\frac{dy}{dx}$  using  $\frac{dy/dt}{dx/dt}$ . Condone poor differentiation.

A1: Correct  $\frac{dy}{dx} = \frac{-4 \sin 2t}{3 \sec^2 t}$ . You may see other forms for this following use of identities.

If you see  $\frac{dy}{dx} = \frac{-4 \sin 2t}{3 \sec^2 x}$  then only allow if they subsequently use this as  $\frac{dy}{dx} = \frac{-4 \sin 2t}{3 \sec^2 t}$

Other possible correct gradients are  $\frac{-8 \sin t \cos t}{3 \sec^2 t}$  and  $\frac{-8 \sin t \cos^3 t}{3}$

ISW after you see a correct answer. Quite often we see candidates incorrectly adapting a correct answer.

It is also possible for candidates to find  $\frac{dy}{dt}$  at  $t = \frac{\pi}{4}$ ,  $\frac{dx}{dt}$  at  $t = \frac{\pi}{4}$  and divide the two values.

dM1: Attempts to find the equation of the tangent at  $t = \frac{\pi}{4}$

It is dependent upon the previous M and use of  $(4, 0)$  and  $t = \frac{\pi}{4}$

Allow substitution into their adapted  $\frac{dy}{dx}$

A1:  $y = -\frac{2}{3}x + \frac{8}{3}$

(b)

B1:  $k = 1 - \sqrt{3}$

(c)

M1: Either end correct. Allow strict inequalities here. E.g  $f < 2$  scores M1 but  $f > 2$  is M0

A1:  $-1 \leq f \leq 2$  o.e such as  $[-1, 2]$  and  $-1 \leq y \leq 2$

Question Number	Scheme	Marks
7 (i)	$u = e^x - 3 \Rightarrow \frac{du}{dx} = e^x$ $\int \frac{4e^{3x}}{e^x - 3} dx = \int \frac{4(u+3)^3}{u(u+3)} du = \int \frac{4(u+3)^2}{u} du$ $= \int \frac{4u^2 + 24u + 36}{u} du$ $= \int 4u + 24 + \frac{36}{u} du = 2u^2 + 24u + 36 \ln u$ $= \left[ 2u^2 + 24u + 36 \ln u \right]_{u=2}^{u=4}$ $= 72 + 36 \ln 2$	B1  M1 A1  dM1 A1  M1 A1  <b>(7)</b>
(ii)	$\int 3e^x \cos 2x dx = \frac{3}{2} e^x \sin 2x - \int \frac{3}{2} e^x \sin 2x dx$ $= \frac{3}{2} e^x \sin 2x - \left( -\frac{3}{4} e^x \cos 2x + \int \frac{3}{4} e^x \cos 2x dx \right)$ <p>Collect terms <math>\frac{15}{4} \int e^x \cos 2x dx = \frac{3}{2} e^x \sin 2x + \frac{3}{4} e^x \cos 2x</math></p> $\int 3e^x \cos 2x dx = \frac{6}{5} e^x \sin 2x + \frac{3}{5} e^x \cos 2x (+c)$	M1 A1  dM1  ddM1  A1  <b>(5)</b> <b>(12 marks)</b>

(i) The 4 must be present to score the A marks. No misreads

B1: States or uses  $\frac{du}{dx} = e^x$  o.e. such as  $\frac{dx}{du} = \frac{1}{u+3}$  This may be seen within the integrand.

M1: A "correct" attempt to write the integrand in terms of  $u$ .

Look for  $\int \frac{4e^{3x}}{e^x - 3} dx = \int \frac{P(u \pm 3)^3}{u(u \pm 3)} du$  which does not need to be simplified.

Condone a missing  $du$  but  $dx$  is M0

A1: Correct  $\int \frac{4(u+3)^2}{u} du$ . Condone a missing  $du$  (cannot be  $dx$ ).

dM1: Integrates by attempting to multiply out and integrating each term.

Look for two "correct" terms, one of which must be  $\int \frac{P}{u} du = P \ln u$ .

It is dependent upon having achieved  $\int \frac{P(u \pm 3)^2}{u} du$

A1: Correct  $2u^2 + 24u + 36\ln u$  o.e.

M1: Uses limits 2 and 4 within their attempted integral and subtracts. Condone poor attempts at the integration  
Alternatively converts their answer in  $u$  back to  $x$ 's using the correct substitution and uses the given limits.

There must have been some attempt to use the substitution  $u = e^x - 3$  at some point.

A1:  $72 + 36\ln 2$  or  $36\ln 2 + 72$

(ii) Condone missing  $dx$ 's here

M1: Attempts to integrate by parts  $\int 3e^x \cos 2x \, dx = Ae^x \sin 2x \pm \int Be^x \sin 2x \, dx$

A1:  $\int 3e^x \cos 2x \, dx = \frac{3}{2}e^x \sin 2x - \int \frac{3}{2}e^x \sin 2x \, dx$  Condone missing  $dx$ 's

The 3 may have been "removed"/factorised out and replaced at the final answer.

If so you can this would be awarded at the sight of a correct answer.

dM1: Attempts to integrate by parts again. See below of what to look for

Look for  $\int 3e^x \cos 2x \, dx = Ae^x \sin 2x \pm Be^x \cos 2x \pm \int Ce^x \cos 2x \, dx$

ddM1: Dependent upon both previous M's

It is awarded for attempting to collect both terms in  $\int e^x \cos 2x \, dx$

A1:  $\int 3e^x \cos 2x \, dx = \frac{6}{5}e^x \sin 2x + \frac{3}{5}e^x \cos 2x (+c)$  condone the omission of the  $+c$ .

Alternative method for (ii). This can be scored in a similar way to the above.

(ii)	$\int 3e^x \cos 2x \, dx = 3e^x \cos 2x + \int 6e^x \sin 2x \, dx$	M1 A1
	$= 3e^x \cos 2x + 6e^x \sin 2x - \int 12e^x \cos 2x \, dx$	dM1
	$15 \int e^x \cos 2x \, dx = 6e^x \sin 2x + 3e^x \cos 2x$	ddM1
	$\int 3e^x \cos 2x \, dx = \frac{6}{5}e^x \sin 2x + \frac{3}{5}e^x \cos 2x (+c)$	A1
<b>(5)</b>		

1stM1:  $\int 3e^x \cos 2x \, dx = Pe^x \cos 2x \pm Q \int e^x \sin 2x \, dx$

2ndM1:  $\int 3e^x \cos 2x \, dx = Pe^x \cos 2x \pm Qe^x \sin 2x \pm R \int e^x \cos 2x \, dx$

Question Number	Scheme	Marks
8	<p>Solves one of <math>(3x - y) = 25</math> and <math>(x + y) = 1</math>  or <math>(3x - y) = 5</math> and <math>(x + y) = 5</math></p> <p>Correct solution of one.</p> <p>Either <math>\left. \begin{array}{l} 3x - y = 25 \\ x + y = 1 \end{array} \right\} \Rightarrow 4x = 26 \Rightarrow x = 6.5, (y = -5.5)</math></p> <p>Or <math>\left. \begin{array}{l} 3x - y = 5 \\ x + y = 5 \end{array} \right\} \Rightarrow 4x = 10 \Rightarrow x = 2.5, (y = 2.5)</math></p> <p>Solves both equations</p> <p>Both solved correctly with a minimal reason given for the contradiction e.g "not integers" with conclusion "hence there are no integers <math>x</math> and <math>y</math> such that <math>3x^2 + 2xy - y^2 = 25</math>"</p>	<p>M1</p> <p>A1</p> <p>dM1</p> <p>A1</p> <p>(4)</p>

#### Notes

M1: Attempts to solve one of the two possible cases.

Take as a minimum, one correct pair of equations followed by a value for  $x$  or a value for  $y$

A1: Correctly solves one of the two possible cases.

To solve you need only find a value for  $x$  or  $y$ . Once a correct value is found you can ISW

dM1: Attempts to solve both possible cases

A1: Correctly solves the two possible cases and makes a concluding argument.

To score this mark (i) **all** calculations must be correct.

(ii) reason(s) for the contradiction must be written down. Allow "not integers", "x" and (iii) gives a concluding statement must be given. Allow for example "hence proven"

Ignore any possible cases which would give rise to negative numbers but satisfy  $(3x - y)(x + y) = 25$

E.g  $(3x - y) = -5, (x + y) = -5$

Withhold the final mark only if they include cases which do not satisfy  $(3x - y)(x + y) = 25$

E.g  $(3x - y) = 20, (x + y) = 5$

Question Number	Scheme	Marks
<p><b>9</b></p>	$\begin{pmatrix} 2-\lambda \\ 8+2\lambda \\ 10+3\lambda \end{pmatrix} = \begin{pmatrix} -4+5\mu \\ -1+4\mu \\ 2+8\mu \end{pmatrix}$ <p>Attempts to solve any two of the three equations</p> <p>Either (1) and (2) <math>\begin{cases} 2-\lambda = -4+5\mu \\ 8+2\lambda = -1+4\mu \end{cases} \Rightarrow \lambda = -\frac{3}{2}, \mu = \frac{3}{2}</math></p> <p>(1) and (3) <math>\begin{cases} 2-\lambda = -4+5\mu \\ 10+3\lambda = 2+8\mu \end{cases} \Rightarrow \lambda = \frac{8}{23}, \mu = \frac{26}{23}</math></p> <p>(2) and (3) <math>\begin{cases} 8+2\lambda = -1+4\mu \\ 10+3\lambda = 2+8\mu \end{cases} \Rightarrow \lambda = -10, \mu = -\frac{11}{4}</math></p> <p>Substitutes their values of <math>\lambda</math> and <math>\mu</math> into both sides of the "third" equation</p> <p>E.g. <math>\lambda = -\frac{3}{2}</math> into <math>10+3\lambda = \frac{11}{2}</math> and <math>\mu = \frac{3}{2}</math> into <math>2+8\mu = 14</math></p> <p>Concludes that lines don't intersect with correct calculations and minimal reason</p> <p>Additionally states that <math>\begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix}</math> is not parallel to <math>\begin{pmatrix} 5 \\ 4 \\ 8 \end{pmatrix}</math> with a minimal reason</p> <p>So lines are skew CSO *</p>	<p>M1, A1</p> <p>dM1</p> <p>A1</p> <p>A1*</p> <p><b>(5)</b> <b>(5 marks)</b></p>

Notes:

Main method seen

M1: Attempts to solve two of the three equations.

Accept as an attempt, writing down two of the three equations (condoning slips) followed by values for both  $\lambda$  and  $\mu$

A1: Solves two of the three equations to find correct values for both  $\lambda$  and  $\mu$ , Allow equivalent fractions

dM1: Either: Substitutes their values of  $\lambda$  and  $\mu$  into both sides of the third equation....or into the equations of both lines to find both coordinates

A1: Having achieved correct values for  $\lambda$  and  $\mu$ , the values for the third equation are found to enable a

comparison to be made. E.g. solving equations (1) and (2) and using equation (3) stating  $10 + 3 \times -\frac{3}{2} \neq 2 + 8 \times \frac{3}{2}$

is sufficient. If the values are found they must be correct.

**Important: Additionally, to score this mark, a minimal statement must be made that states that the lines do not intersect /cross. Condone statements such as  $l_1 \neq l_2$**

**Stating that the lines are skew at this point is not sufficient to score this mark**

In the alternative stating that "as the values are not the same, the lines cannot intersect" is sufficient.

A1\*: CSO . Hence all previous marks must have been scored.

In addition to not intersecting there must be a statement, with a minimal reason, that the lines are not parallel and hence skew. Accept statements like, not intersecting, not parallel (with reason), hence proven.

Reasons could be  $\begin{pmatrix} 5 \\ 4 \\ 8 \end{pmatrix} \neq k \begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix}$  o.e such as  $5 = -5 \times -1$  but  $4 = 2 \times 2$  so they are not parallel.

Accept an argument based around the scalar product of the direction vectors. If parallel  $\cos \theta = 1$

A reason for the lines not being parallel cannot be  $\begin{pmatrix} 5 \\ 4 \\ 8 \end{pmatrix} \neq \begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix}$

Note: Other methods are possible and it is important that you look at their complete attempt at proving that they don't intersect.

Alternative 1

For example it is possible to solve equations (1) and (2) to find just  $\lambda$

then solve equations (1) and (3) to find just  $\lambda$

and then conclude that "as the two values are not the same, the lines don't intersect"

M1 dM1 marks are scored together. Both aspects have to be attempted

Attempts to solve two of the three equations to find  $\lambda$  (or  $\mu$ )


Attempts to solve a different pair of equations to find  $\lambda$  (or  $\mu$ )

A1: Correct values for  $\lambda$  (or  $\mu$ ).

A1: conclude that "as the two values are not the same, the lines don't intersect"

If you see something that you feel deserves credit AND that you cannot mark, then please send to review



Question Number	Scheme	Marks
10 (a)	<p>States or uses <math>\frac{dr}{dt} \propto \pm \frac{1}{r^2}</math> or <math>\frac{dr}{dt} = \pm \frac{k}{r^2}</math></p> $\frac{dr}{dt} = \pm \frac{k}{r^2} \Rightarrow \int r^2 dr = \int \pm k dt$ $\frac{1}{3} r^3 = \pm kt + c$ <p>Substitutes <math>t = 0, r = 12 \Rightarrow c = (576)</math></p> <p>AND <math>t = 15, r = 6</math> and their <math>c = (576) \Rightarrow k = (\pm 33.6)</math></p> $\frac{1}{3} r^3 = -33.6t + 576 \text{ or } r^3 = -100.8t + 1728 \text{ o.e.}$	<p>B1</p> <p>M1 A1</p> <p>M1</p> <p>A1</p> <p>(5)</p> <p>M1</p> <p>A1</p> <p>(2)</p> <p>B1</p> <p>(1)</p> <p><b>(8 marks)</b></p>
(b)	<p>Sets <math>r = 0 \Rightarrow -100.8t + 1728 = 0</math></p> $\frac{120}{7} \text{ minutes or awrt } 17.1 \text{ minutes}$	<p>M1</p> <p>A1</p>
(c)		<p>B1</p>

(a)

B1: States or uses  $\frac{dr}{dt} \propto \pm \frac{1}{r^2}$ ,  $\frac{dr}{dt} = \pm \frac{k}{r^2}$  or  $\frac{dr}{dt} = \pm \frac{1}{kr^2}$  Accept  $k \leftrightarrow \lambda$  or other constant. Condone  $\frac{dr}{dt} = \pm \frac{1}{r^2}$

M1: Integrates to achieve an expression of the correct form  $\dots r^3 = \dots t (+c)$  o.e

There is no requirement for  $+c$

A1: Correct integration with two different unknown constants. E.g. look for  $\frac{1}{3} r^3 = \pm kt + c$  o.e

So if for example  $\frac{dr}{dt} = -\frac{1}{r^2} \Rightarrow \frac{1}{3} r^3 = -t + c$  the score would be B1, M1, A0 then dM0, A0

M1: Uses both boundary conditions to form an equation involving  $r$  and  $t$ . For this to be awarded there must be two different unknown constants that are initially correctly placed.

It is dependent upon having achieved  $\dots r^n = \pm k "t + "c"$  which may have been achieved from an incorrect assumption. E.g.  $\frac{dr}{dt} = \pm kr^2$

A1: Correct equation. E.g.  $\frac{1}{3}r^3 = -33.6t + 576$  or  $r^3 = -100.8t + 1728$  o.e. such as  $t = -\frac{5}{504}r^3 + \frac{120}{7}$

(b)

M1: Sets  $r = 0 \Rightarrow -100.8t + 1728 = 0 \Rightarrow t = \dots$  Follow through on their equation. Condone if this produces a negative value for  $t$ . Alt sets  $V = 0$  to find  $t = \dots$

A1: Awrt 17.1 minutes. Must include the units. Also allow 17 minutes 8 seconds or 17 minutes 9 seconds


(c)

B1: For the correct shaped curve only in quadrant 1 starting at (0,12), ignoring value of  $t$ .

As  $\frac{dr}{dt} = -\frac{k}{r^2}$  the gradient should appear to get increasingly steeper.

If there are two curves given on the axes, both need to be correct for this mark to be awarded

Do not be concerned if it is not infinite at the  $t$  axis.



The curve on the left would be at the limit of what we would allow as the gradient on the left hand side does appear to increase.

We condone its appearance as it approaches the  $t$ -axis.

It does not get less steep

Note that it is possible to use  $V = \frac{4}{3}\pi r^3 \Rightarrow \frac{dV}{dr} = 4\pi r^2$  with  $\frac{dr}{dt} = \pm \frac{k}{r^2}$  to set up and solve  $\frac{dV}{dt} = \pm 4\pi k$  or

$$\frac{dV}{dt} = \pm \beta \quad \text{In this case the equation becomes } V = 2304\pi - \left(\frac{672\pi}{5}\right)t$$

(a)

B1: States or uses either  $\frac{dV}{dt} = \pm \beta$  or its exact value which is  $\frac{dV}{dt} = -\frac{2016\pi}{15}$  o.e. such as  $\frac{dV}{dt} = -\frac{672\pi}{5}$ .

M1: For  $\frac{dV}{dt} = \pm \beta \Rightarrow V = \pm \beta t + c$  and attempts to find the values of  $\beta$  and  $c$  using the given conditions.

It is dependent upon knowing that  $V = \alpha r^3$  and using this to find  $V$  at  $r = 6$  and  $12$  with  $t = 15$  and  $t = 0$

$$\text{A1: } V = 2304\pi - \left(\frac{672\pi}{5}\right)t$$

M1: Substitutes  $V = \alpha r^3$  to form an equation linking  $r^3$  with  $t$ .

It is dependent on the previous M1 in this method

$$\text{A1: Achieves } \frac{4}{3}\pi r^3 = 2304\pi - \left(\frac{672\pi}{5}\right)t$$

(b)

M1: Sets  $V = 0$  o.e and finds  $t$

Question Number	Scheme	Marks
<b>11 (a)</b>	$(x + y)^3 + 10y^2 = 108x$	
	Differentiates $10y^2$ to $20y \frac{dy}{dx}$	B1
	Differentiates $(x + y)^3$ to $3(x + y)^2 \left(1 + \frac{dy}{dx}\right)$	M1
	$3(x + y)^2 \left(1 + \frac{dy}{dx}\right) + 20y \frac{dy}{dx} = 108$	A1
	$3(x + y)^2 \frac{dy}{dx} + 20y \frac{dy}{dx} = 108 - 3(x + y)^2$	dM1
	$\frac{dy}{dx} = \frac{108 - 3(x + y)^2}{20y + 3(x + y)^2}$ *	A1*
		<b>(5)</b>
<b>(b)</b>	Deduces that $P$ and $Q$ are where $108 - 3(x + y)^2 = 0 \Rightarrow (x + y)^2 = 36$	M1
	Attempts to substitute $(x + y) = \pm 6$ into $(x + y)^3 + 10y^2 = 108x$ to form an equation in $y$ (or $x$ )	dM1
	Solves $216 + 10y^2 = 108(6 - y)$ and finds the negative root	ddM1
	Awrt 13900 metres	A1
		<b>(4)</b>
		<b>(9 marks)</b>

(a) Allow use of  $y'$  for  $\frac{dy}{dx}$ . This is a proof. Look carefully for evidence of candidates working backwards.

B1: Differentiates  $10y^2$  to  $20y \frac{dy}{dx}$

M1: Attempts to differentiate  $(x + y)^3$  Via the chain rule look for  $k(x + y)^2 \left(1 + \frac{dy}{dx}\right)$  See \*\*

Alternatively you will see attempts where  $(x + y)^3$  is multiplied out. Terms may not be collected but

look for 3 of  $x^3 \rightarrow \dots x^2$ ,  $\dots x^2 y \rightarrow 2xy + x^2 \frac{dy}{dx}$ ,  $\dots xy^2 \rightarrow y^2 + 2xy \frac{dy}{dx}$  and  $y^3 \rightarrow 3y^2 \frac{dy}{dx}$

differentiated to the correct form.

A1: Correct differentiation  $3(x + y)^2 \left(1 + \frac{dy}{dx}\right) + 20y \frac{dy}{dx} = 108$

$$\text{or } 3x^2 + 3\left(2xy + x^2 \frac{dy}{dx}\right) + 3\left(y^2 + 2xy \frac{dy}{dx}\right) + 3y^2 \frac{dy}{dx} + 20y \frac{dy}{dx} = 108$$

dM1: Collects the terms in  $\frac{dy}{dx}$ .

Look for either  $\frac{dy}{dx}$  being factorised out of the relevant terms

or the terms in  $\frac{dy}{dx}$  being set on one side of the equation and the other terms on the other side

It is dependent upon **BOTH** the B and M marks.

A1\*: Proceeds to the given answer.

In the approach  $3x^2 + 3\left(2xy + x^2 \frac{dy}{dx}\right) + 3\left(y^2 + 2xy \frac{dy}{dx}\right) + 3y^2 \frac{dy}{dx} + 20y \frac{dy}{dx} = 108$  it is acceptable to go from  $(3x^2 + 6xy + 3y^2 + 20y) \frac{dy}{dx} = 108 - 3x^2 - 6xy - 3y^2$  straight to the given answer

---

**Watch for candidates who use the given answer and work backwards\*\*.**

$$\frac{dy}{dx} = \frac{108 - 3(x+y)^2}{20y + 3(x+y)^2} \Rightarrow 3(x+y)^2 \frac{dy}{dx} + 20y \frac{dy}{dx} = 108 - 3(x+y)^2$$

Therefore candidates who start with

$$3(x+y)^2 \frac{dy}{dx} + 20y \frac{dy}{dx} = 108 - 3(x+y)^2 \quad \text{or indeed} \quad 3(x+y)^2 \frac{dy}{dx} + 3(x+y)^2 + 20y \frac{dy}{dx} = 108$$

without any other working or evidence can only score B1 M0 A0 M0 A0

---

(b)

M1: Deduces that  $P$  and  $Q$  are where  $108 - 3(x+y)^2 = 0 \Rightarrow (x+y)^2 = 36$

This is implied if there is a statement that  $(x+y) = \pm 6$

dM1: Attempts to solve  $(x+y) = \pm 6$  and  $(x+y)^3 + 10y^2 = 108x$  simultaneously to form a quadratic equation in  $y$  or  $x$ .

ddM1: Solves  $216 + 10y^2 = 108(6-y)$  and finds the negative root. Allow if both roots are found

Note that  $(x+y) = 6$  must have been used.

If an equation was set up in  $x$ , then look for an attempt to find the larger root of

$$216 + 10(6-x)^2 = 108x \quad \text{which should then be used to find a negative } y \text{ value}$$

A1: Awrt 13900 metres or awrt 13.9 km with the correct units. Condone answers like  $-13.9$  km