

# Mathematics (MEI)

Advanced Subsidiary GCE

Unit **4751**: Introduction to Advanced Mathematics

## Mark Scheme for June 2012

---

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2012

Any enquiries about publications should be addressed to:

OCR Publications  
PO Box 5050  
Annesley  
NOTTINGHAM  
NG15 0DL

Telephone: 0870 770 6622  
Facsimile: 01223 552610  
E-mail: [publications@ocr.org.uk](mailto:publications@ocr.org.uk)

## Annotations

Annotation in scoris	Meaning
✓and *	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining
U1	Mark for correct units
G1	Mark for a correct feature on a graph
M1 dep*	Method mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working

### Subject-specific Marking Instructions: GCE Mathematics (MEI) Pure strand

- a. Annotations should be used whenever appropriate during your marking.

**The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded marks.** It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded.

- b. An element of professional judgement is required in the marking of any written paper. Remember that the mark schemes assist in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must be the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the work must be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such methods must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, award marks in the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) discuss the matter with your Team Leader.

- c. The following types of marks are available.

**M**

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood and not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate to state an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of a mark is specified.

**A**

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

**B**

Mark for a correct result or statement independent of Method marks.

## E

A given result is to be established or a result has to be explained. This usually requires more working or a correct argument than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, eg wrong working following a correct answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply if a candidate passes through the correct answer as part of a wrong argument.

- d. When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme indicates otherwise; and similarly where there are several B marks allocated. (The notation 'dep \*' is used to indicate that a part is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate is wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, if two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e. The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect work. Otherwise, A and B marks are given for correct work only — differences in notation are of course permitted. A (accuracy) mark is given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will be given for 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is in a different image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f. Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise. Candidates are expected to give numerical answers to an appropriate degree of accuracy, with 3 significant figures often being the expected accuracy. Variations in the degree of accuracy to which an answer is given (e.g. 2 or 4 significant figures where 3 is expected) should not be penalised, while answers which are grossly over- or under-specified should normally result in the loss of a mark. The mark scheme will specify any particular cases where the accuracy of the answer may be a marking issue should be detailed in the mark scheme rationale. If this is not the case please contact your Team Leader.

g.

Rules for replaced work

If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, the candidate requests.

If there are two or more attempts at a question which have not been crossed out, examiners should mark what (complete) attempt and ignore the others.

NB Follow these maths-specific instructions rather than those in the assessor handbook.

h.

For a *genuine* misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally this may differ for some units. This is achieved by withholding one A mark in the question.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

Question		Answer	Marks	Guida.
1		$y = -2x + 7$ isw  (0, 7) and (3.5, 0) oe or ft their $y = -2x + c$	2  1  <b>[3]</b>	M1 for $y - 1 = -2(x - 3)$ or $1 = -2 \times 3 + c$ oe  condon. $x = 3.5$ or such as (3.5, seen
2		$[b =] \pm \sqrt{\frac{3a}{2c}}$ oe www	3  <b>[3]</b>	M2 for $[b^2 =] \frac{3a}{2c}$ soi  or M1 for other $[b^2 =] \frac{ka}{c}$ or $[b^2 =] \frac{a}{kc}$ oe  and M1 for correctly taking the square root of their $b^2$ , including the $\pm$ sign;  eg M2 for $[b =]$  allow M1 for a quadruple-dec eg $\frac{1.5a}{c}$ , if no  square root m fraction line
3	(i)	25	2  <b>[2]</b>	M1 for $\frac{1}{\frac{1}{25}}$ or $\left(\frac{1}{25}\right)^{-1}$ or $5^2$ or $\frac{25}{1}$
3	(ii)	$\frac{4}{9}$	2  <b>[2]</b>	M1 for 4 or 9 or $\frac{1}{9}$ or $\frac{2}{3}$ or $\left(\frac{2}{3}\right)^2$ or $\sqrt[3]{\frac{64}{729}}$  seen  0 for just $\left(\frac{64}{729}\right)$
4		$\frac{x-3}{x+2}$ or $1 - \frac{5}{x+2}$ as final answer www	3  <b>[3]</b>	B2 for correct answer seen and then spoilt M1 for $(x+3)(x-3)$ and M1 for $(x+2)(x+3)$

Question		Answer	Marks	Guidance
5	(i)	30	3       <b>[3]</b>	M1 for $(\sqrt{6})^3 = 6\sqrt{6}$ soi and M1 for $\sqrt{24} = 2\sqrt{6}$ soi  or allow SC2 for final answer of $5(\sqrt{6})^2$ or $5\sqrt{36}$ or $10\sqrt{9}$ etc
5	(ii)	$\frac{8}{11}$	2       <b>[2]</b>	M1 for common denominator $(4 + \sqrt{5})(4 - \sqrt{5})$ soi - may be in separate fractions or for a final answer with denominator 11, even if worked with only one fraction
6	(i)	10 cao	1 <b>[1]</b>	
6	(ii)	$-720 [x^3]$	4       <b>[4]</b>	condone $-720$  allow equivalent forms as part of a long answer eg M2 for $3^5$ for $10 \times \left(\frac{-2}{3}\right)^3$

Question	Answer	Marks	Guidance
7	$4k^2 - 4 \times 1 \times 5$ or $k^2 - 5$ [ $< 0$ ] oe or $[(x+k)^2 +] 5 - k^2$ [ $> 0$ ] oe  $-\sqrt{5} < k < \sqrt{5}$	M2          A2   <b>[4]</b>	allow =, >, < etc instead of <  or M1 for $b^2 - 4ac$ soi (may be in formula) or for attempt at completing square       may be two separate inequalities or A1 for one 'end' correct or B1 for 'endpoint' = $\sqrt{5}$
8	$16 + 2b + c = 0$ oe  $81 - 3b + c = 85$ oe  $20 + 5b = 0$ oe  $b = -4$ and $c = -8$	M1   B2       M1       A1   <b>[5]</b>	need not be simplified; condone 8 or 32 as first term if $2^4$ not seen       M1 for $f(-3)$ seen or used, condoning one error except $+3b$ – need not be simplified or for long division as far as obtaining $x^3 - 3x^2$ in quotient       for elimination of one variable, ft their equations in $b$ and $c$ , condoning one error in rearrangement of their original equations or in one term in the elimination       allow correct answers to imply last M1 after correct earlier equations

Question	Answer	Marks	Guidance
9	$6n + 9$ isw or $3(2n + 3)$  $6n$ is even [but 9 is odd], even + odd = odd or $2n + 3$ is odd since even + odd = odd and $odd \times odd = odd$  'n is a multiple of 3' or 'n is divisible by 3' without additional incorrect statement(s)	B1  B1 dep  B2  [4]	this mark is dependent on the previous B1  accept equiv. general statements using either $6n + 9$ or $3(2n + 3)$  B2 for 'it is divisible by 9, so n is divisible by 3'  M1 for '6n is divisible by 9' or '2n + 3 is divisible by 3' or for 'n is a multiple of 3' oe with additional incorrect statement(s)

B2 for just 'it  
 but M1 for 'it  
 divisible by 3'  
  
 eg M1 for 'n  
 divisible by 3'  
  
 N.B. 0 for 'n i  
 may be earned



Question		Answer	Marks	Guidance
10	(ii)	[grad. of AC =] $\frac{5 - (-1)}{1 - 3}$ or $\frac{6}{-2}$ oe	M1	award at first step shown even if errors after
		[grad. of BD =] $\frac{5 - 1}{11 - (-1)}$ or $\frac{4}{12}$ oe	M1	if one o... grad BD =... working for u... M1 only. For... that they are o
		showing or stating product of gradients = -1 or that one gradient is the negative reciprocal of the other oe	B1	eg accept $m_1 \times m_2 = -1$ or 'one gradient is negative reciprocal of the other'  B0 for 'opposite' used instead of 'negative' or 'reciprocal'
			[3]	may be earned... correct gradien... be earned the... correct

Question		Answer	Marks	Guidance
10	(iii)	midpoint E of AC = (2, 2) www	B1	condone missing brackets for both B1s
		eqn BD is $y = \frac{1}{3}x + \frac{4}{3}$ oe	M1	accept any correct form isw or correct ft their gradients or their midpt F of BD  this mark will often be gained on the first line of their working for BD
		eqn AC is $y = -3x + 8$ oe	M1	accept any correct form isw or correct ft their gradients or their midpt E of AC  this mark will often be gained on the first line of their working for AC  [see appendix for alternative methods instead showing E is on BD for this M1]
		using both lines and obtaining intersection E is (2, 2) (NB must be independently obtained from midpt of AC)	A1	
		midpoint F of BD = (5,3)	B1	this mark is often earned earlier  see the appendix for some common alternative methods for this question; for all methods, for A1 to be earned, all work for the 5 marks must be correct
			[5]	

Question		Answer	Marks	Guidance
11	(i)	$(2x + 1)(x + 2)(x - 5)$  correct expansion of two linear factors of their product of three linear factors  expansion of their linear and quadratic factors   $[y =] 2x^3 - 5x^2 - 23x - 10$ or $a = -5, b = -23$ and $c = -10$	M1  M1  M1  A1  [4]	or $(x + 1/2)(x + 2)(x - 5)$ ; need not be written as product  dep on first M1; ft one error in previous expansion; condone one error in this expansion  or for direct expansion of all three factors, allow M2 for $2x^3 - 10x^2 + 4x^2 + x^2 - 20x - 5x + 2x - 10$ [or half all these], or M1 if one or two errors,  condone poor 'doubling' to $2x^3...$  $250 + 25a + 5$ $-16 + 4a - 2b$ $-1/4 + 1/4 a - 1/2$  then M2 for correctly eliminating any two variables or M1 for correctly eliminating one variable to get two equations in two unknowns  and then A1 for values.

Question		Answer	Marks	Guidance
11	(ii)	<p>graph of cubic correct way up</p> <p>crossing <math>x</math> axis at <math>-2, -1/2</math> and <math>5</math></p> <p>crossing <math>y</math> axis at <math>-10</math> or ft their cubic in (i)</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>[3]</p>	<p>B0 if stops at <math>x</math>-axis on graph or nearby in this part</p> <p>mark intent for intersections with both axes</p> <p>or <math>x = 0, y = -10</math> or ft in this part if consistent with graph drawn;</p>
11	(iii)	$(0, -18)$ ; accept $-18$ or ft their constant $-8$	<p>1</p> <p>[1]</p>	or ft their intn on $y$ -axis $-8$
11	(iv)	<p>roots at <math>2.5, 1, 8</math></p> <p><math>(2x - 5)(x - 1)(x - 8)</math></p> <p><math>(0, -40)</math>; accept <math>-40</math></p>	<p>M1</p> <p>A1</p> <p>B2</p> <p>[4]</p>	<p>or attempt to substitute <math>(x - 3)</math> in <math>(2x + 1)(x + 2)(x - 5)</math> or in <math>(x + 1/2)(x + 2)(x - 5)</math> or in their unfactorised form of <math>f(x)</math>– attempt need not be simplified</p> <p>accept <math>2(x - 2.5)</math> oe instead of <math>(2x - 5)</math></p> <p>M1 for <math>-5 \times -1 \times -8</math> or ft or for <math>f(-3)</math> attempted or <math>g(0)</math> attempted or for their answer ft from their factorised form</p>

Question		Answer	Marks	Guidance
12	(i)	(-1, 6) (0,1) (1,-2) (2,-3) (3,-2) (4, 1) (5,6) seen plotted	B2	or for a curve within 2 mm of these points; B1 for 3 correct plots or for at least 3 of the pairs of values seen eg in table
		smooth curve through all 7 points	B1 dep	dep on correct points; tolerance 2 mm;
		(0.3 to 0.5, -0.3 to -0.5) and (2.5 to 2.7, -2.5 to -2.7) and (4, 1)	B2	may be given in form $x = \dots, y = \dots$ B1 for two intersections correct or for all the $x$ values given correctly
			[5]	
12	(ii)	$\frac{1}{x-3} = x^2 - 4x + 1$ $1 = (x-3)(x^2 - 4x + 1)$	M1	
		at least one further correct interim step with '=1' or '=0', as appropriate, leading to given answer, which must be stated correctly	M1  A1	condone omission of brackets only if used correctly afterwards, with at most one error;  there may also be a previous step of expansion of terms without an equation, eg in grid  if M0, allow SC1 for correct division of given cubic by quadratic to gain $(x-3)$ with remainder $-1$ , or vice-versa
			[3]	

Question	Answer	Marks	Guidance
12 (iii)	quadratic factor is $x^2 - 3x + 1$  substitution into quadratic formula or for completing the square used as far as $(x - \frac{3}{2})^2 = \frac{5}{4}$ $\frac{3 \pm \sqrt{5}}{2}$ oe	B2  M1  A2  <b>[5]</b>	found by division or inspection; allow M1 for division by $x - 4$ as far as $x^3 - 4x^2$ in the working, or for inspection with two terms correct  condone one error  A1 if one error in final numerical expression, but only if roots are real  no ft from a w  isw factors

Appendix: alternative methods for 10(iii) [details of equations etc are in main scheme]

for a mixture of methods, look for the method which gives most benefit to candidate, but take care not to award the second M1 twice the final A1 is not earned if there is wrong work leading to the required statements

ignore wrong working which has not been used for the required statements

for full marks to be earned in this part, there must be enough to show both the required statements

find midpt E of AC find eqn BD	B1 M1	find midpt E of AC find eqn BD	B1 M1	find midpt E of AC find eqn BD	B1 M1	find midpt E of AC use gradients show E is on BD grad BE = $\frac{2-1}{2-1}$ ED = $\frac{5-2}{11-2} = \frac{1}{3}$ [condone poor notation]
show E on BD find midpt F of BD	M1 B1	show E on BD find midpt F of BD	M1 B1	show E on BD show $BE^2 = 10$ and $DE^2 = 90$ oe	M1 B1	find midpt F of BD
state so not E	A1	find eqn of AC and correctly show F not on AC (the correct eqn for AC earns the second M1 as per the main scheme, if not already earned)	A1	showing $BE^2 = 10$ and $DE^2 = 90$ oe earns this A mark as well as the B1 if there are no errors elsewhere	A1	state so not E show F not on AC
	<b>[5]</b>					