

Mathematics (MEI)

Advanced GCE A2 7895-8

Advanced Subsidiary GCE AS 3895-8

Mark Schemes for the Units

January 2008

3895-8/7895-8/MS/R/08J

OCR (Oxford, Cambridge and RSA Examinations) is a unitary awarding body, established by the University of Cambridge Local Examinations Syndicate and the RSA Examinations Board in January 1998. OCR provides a full range of GCSE, A level, GNVQ, Key Skills and other qualifications for schools and colleges in the United Kingdom, including those previously provided by MEG and OCEAC. It is also responsible for developing new syllabuses to meet national requirements and the needs of students and teachers.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2008

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552310
E-mail: publications@ocr.org.uk

CONTENTS

Advanced GCE Further Mathematics (MEI) (7896)
Advanced GCE Further Mathematics (Additional) (MEI) (7897)
Advanced GCE Mathematics (MEI) (7895)
Advanced GCE Pure Mathematics (MEI) (7898)

Advanced Subsidiary GCE Further Mathematics (MEI) (3896)
Advanced Subsidiary GCE Further Mathematics (Additional) (MEI) (3897)
Advanced Subsidiary GCE Mathematics (MEI) (3895)
Advanced Subsidiary GCE Pure Mathematics (MEI) (3898)

MARK SCHEME FOR THE UNITS

Unit/Content	Page
4751 (C1) Introduction to Advanced Mathematics	1
4752 (C2) Concepts for Advanced Mathematics	5
4753 (C3) Methods for Advanced Mathematics	7
4754 (C4) Applications of Advanced Mathematics	11
4755 (FP1) Further Concepts for Advanced Mathematics	16
4756 (FP2) Further Methods for Advanced Mathematics	20
4758 Differential Equations	27
4761 Mechanics 1	31
4762 Mechanics 2	37
4763 Mechanics 3	41
4766 Statistics 1	46
4767 Statistics 2	50
4768 Statistics 3	54
4771 Decision Mathematics 1	58
4776 Numerical Methods	64
Grade Thresholds	66

4751 (C1) Introduction to Advanced Mathematics

Section A

1	$[v =][\pm] \sqrt{\frac{2E}{m}}$ www	3	M2 for $v^2 = \frac{2E}{m}$ or for $[v =][\pm] \sqrt{\frac{E}{\frac{1}{2}m}}$ or M1 for a correct constructive first step and M1 for $v = [\pm] \sqrt{k}$ ft their $v^2 = k$; if M0 then SC1 for $\sqrt{E/ \frac{1}{2} m}$ or $\sqrt{2E/m}$ etc	3
2	$\frac{3x-4}{x+1}$ or $3 - \frac{7}{x+1}$ www as final answer	3	M1 for $(3x - 4)(x - 1)$ and M1 for $(x + 1)(x - 1)$	3
3	(i) 1 (ii) 1/64 www	1 3	M1 for dealing correctly with each of reciprocal, square root and cubing (allow 3 only for 1/64) eg M2 for 64 or -64 or $1/\sqrt{4096}$ or $\frac{1}{4^3}$ or M1 for $1/16^{3/2}$ or 4^3 or -4^3 or 4^{-3} etc	4
4	$6x + 2(2x - 5) = 7$ $10x = 17$ $x = 1.7$ o.e. isw $y = -1.6$ o.e. isw	M1 M1 A1 A1	for subst or multn of eqns so one pair of coeffs equal (condone one error) simplification (condone one error) or appropriate addn/subtn to eliminate variable allow as separate or coordinates as requested graphical soln: M0	4
5	(i) -4/5 or -0.8 o.e. (ii) (15, 0) or 15 found www	2 3	M1 for 4/5 or 4/-5 or 0.8 or -4.8/6 or correct method using two points on the line (at least one correct) (may be graphical) or for -0.8x o.e. M1 for $y =$ their (i) $x + 12$ o.e. or $4x + 5y = k$ and (0, 12) subst and M1 for using $y = 0$ eg $-12 = -0.8x$ or ft their eqn <u>or</u> M1 for given line goes through (0, 4.8) and (6, 0) and M1 for $6 \times 12/4.8$ graphical soln: allow M1 for correct required line drawn and M1 for answer within 2mm of (15, 0)	5

6	f(2) used $2^3 + 2k + 7 = 3$ $k = -6$	M1 M1 A1	or division by $x - 2$ as far as $x^2 + 2x$ obtained correctly or remainder $3 = 2(4 + k) + 7$ o.e. 2nd M1 dep on first	3
7	(i) 56 (ii) -7 or ft from -their (i)/8	2 2	M1 for $\frac{8 \times 7 \times 6}{3 \times 2 \times 1}$ or more simplified M1 for 7 or ft their (i)/8 or for $56 \times (-1/2)^3$ o.e. or ft; condone x^3 in answer or in M1 expression; 0 in qn for just Pascal's triangle seen	4
8	(i) $5\sqrt{3}$ (ii) common denominator = $(5 - \sqrt{2})(5 + \sqrt{2}) = 23$ numerator = 10	2 M1 A1 B1	M1 for $\sqrt{48} = 4\sqrt{3}$ allow M1A1 for $\frac{5 - \sqrt{2}}{23} + \frac{5 + \sqrt{2}}{23}$ allow 3 only for 10/23	5
9	(i) $n = 2m$ $3n^2 + 6n = 12m^2 + 12m$ or $= 12m(m + 1)$ (ii) showing false when n is odd e.g. $3n^2 + 6n = \text{odd} + \text{even} = \text{odd}$	M1 M2 B2	or any attempt at generalising; M0 for just trying numbers or M1 for $3n^2 + 6n = 3n(n + 2) = 3 \times \text{even} \times \text{even}$ and M1 for explaining that 4 is a factor of even \times even or M1 for 12 is a factor of $6n$ when n is even and M1 for 4 is a factor of n^2 so 12 is a factor of $3n^2$ or $3n(n + 2) = 3 \times \text{odd} \times \text{odd} = \text{odd}$ or counterexample showing not always true; M1 for false with partial explanation or incorrect calculation	5

Section B

10	i	correct graph with clear asymptote $x = 2$ (though need not be marked)	G2	G1 for one branch correct; condone $(0, -\frac{1}{2})$ not shown SC1 for both sections of graph shifted two to left	
	ii	$(0, -\frac{1}{2})$ shown 11/5 or 2.2 o.e. isw	G1	allow seen calculated	3
	iii	$x = \frac{1}{x-2}$ $x(x-2) = 1$ o.e. $x^2 - 2x - 1 [= 0]$; ft their equiv eqn attempt at quadratic formula $1 \pm \sqrt{2}$ cao position of points shown	2 M1 M1 M1 M1 A1 B1	M1 for correct first step or equivs with ys or $(x-1)^2 - 1 = 1$ o.e. or $(x-1) = \pm\sqrt{2}$ (condone one error) on their curve with $y = x$ (line drawn or $y = x$ indicated by both coords); condone intent of diagonal line with gradient approx 1 through origin as $y = x$ if unlabelled	2 6
11	i	$(x-2.5)^2$ o.e. $-2.5^2 + 8$ $(x-2.5)^2 + 7/4$ o.e. min $y = 7/4$ o.e. [so above x axis] or commenting $(x-2.5)^2 \geq 0$	M1 M1 A1 B1	for clear attempt at -2.5^2 allow M2A0 for $(x-2.5) + 7/4$ o.e. with no $(x-2.5)^2$ seen ft, dep on $(x-a)^2 + b$ with b positive; condone starting again, showing $b^2 - 4ac < 0$ or using calculus	4
ii	correct symmetrical quadratic shape 8 marked as intercept on y axis tp $(5/2, 7/4)$ o.e. or ft from (i)	G1 G1 G1	or $(0, 8)$ seen in table	3	
iii	$x^2 - 5x - 6$ seen or used -1 and 6 obtained $x < -1$ and $x > 6$ isw or ft their solns	M1 M1 M1	or $(x-2.5)^2$ [$>$ or $=$] 12.25 or ft $14 - b$ also implies first M1 if M0, allow B1 for one of $x < -1$ and $x > 6$	3	
iv	min = $(2.5, -8.25)$ or ft from (i) so yes, crosses	M1 A1	or M1 for other clear comment re translated 10 down and A1 for referring to min in (i) or graph in (ii); or M1 for correct method for solving $x^2 - 5x - 2 = 0$ or using $b^2 - 4ac$ with this and A1 for showing real solns eg $b^2 - 4ac = 33$; allow M1A0 for valid comment but error in -8.25 ft; allow M1 for showing y can be neg eg $(0, -2)$ found and A1 for correct conclusion	2	


11

12

12	i	$(x - 4)^2 - 16 + (y - 2)^2 - 4 = 9$ o.e. $\text{rad} = \sqrt{29}$	M2	M1 for one completing square or for $(x - 4)^2$ or $(y - 2)^2$ expanded correctly <u>or</u> starting with $(x - 4)^2 + (y - 2)^2 = r^2$: M1 for correct expn of at least one bracket and M1 for $9 + 20 = r^2$ o.e.	3
			B1	<u>or</u> using $x^2 - 2gx + y^2 - 2fy + c = 0$ M1 for using centre is (g, f) [must be quoted] and M1 for $r^2 = g^2 + f^2 - c$	
	ii	$4^2 + 2^2$ o.e. $= 20$ which is less than 29	M1 A1	allow 2 for showing circle crosses x axis at -1 and 9 or equiv for y (or showing one positive; one negative); 0 for graphical solutions (often using A and B from (iii) to draw circle)	2
	iii	showing midpt of AB = $(4, 2)$ and showing AB = $2\sqrt{29}$ or showing AC or BC = $\sqrt{29}$ or that A or B lie on circle <u>or</u> showing both A and B lie on circle (or AC = BC = $\sqrt{29}$), and showing AB = $2\sqrt{29}$ or that C is midpt of AB or that C is on AB or that gradients of AB and AC are the same or equiv. <u>or</u> showing C is on AB and showing both A and B are on circle or AC = BC = $\sqrt{29}$	2 2 2 2	in each method, two things need to be established. Allow M1 for the concept of what should be shown and A1 for correct completion with method shown allow M1A0 for AB just shown as $\sqrt{116}$ not $2\sqrt{29}$ allow M1A0 for stating mid point of AB = $(4, 2)$ without working/method shown NB showing AB = $2\sqrt{29}$ and C lies on AB is not sufficient – earns 2 marks only	4
iv	$\text{grad AC or AB or BC} = -5/2$ o.e. $\text{grad tgt} = -1/\text{their grad AC}$ $\text{tgt is } y - 7 = \text{their } m(x - 2)$ o.e. $y = 2/5x + 31/5$ o.e.	M1 M1 M1 A1	may be seen in (iii) but only allow this M1 if they go on to use in this part allow for $m_1m_2 = -1$ used eg $y = \text{their } mx + c$ then $(2, 7)$ subst; M0 if grad AC used condone $y = 2/5x + c$ and $c = 31/5$ o.e.	4	

4752 (C2) Concepts for Advanced Mathematics

Section A


1	$40x^3$	2	-1 if extra term	2
2	(i) 3 (ii) 141	1 2	M1 for $9 \times (1 + 2 + 3 + 4 + 5) + 1 + 2 + 3$	3
3	right angled triangle with 1 and 2 on correct sides Pythagoras used to obtain hyp = $\sqrt{5}$ $\cos \theta = \frac{a}{h} = \frac{2}{\sqrt{5}}$	M1 M1 A1	or M1 for $\sin \theta = \frac{1}{2} \cos \theta$ and M1 for substituting in $\sin^2 \theta + \cos^2 \theta = 1$ E1 for sufficient working	3
4	(i) line along $y = 6$ with V (1, 6), (2, 2), (3, 6) (ii) line along $y = 3$ with V (-2, 3), (-1, 1), (0, 3)	2 2	1 for two points correct 1 for two points correct	4
5	$2x^6 + \frac{3}{4}x^{\frac{4}{3}} + 7x + c$	5	1 for $2x^6$; 2 for $\frac{3}{4}x^{\frac{4}{3}}$ or 1 for other $kx^{\frac{4}{3}}$; 1 for $7x$; 1 for $+c$	5
6	(i) correct sine shape through O amplitude of 1 and period 2π shown (ii) $7\pi/6$ and $11\pi/6$	1 1 3	B2 for one of these; 1 for $-\pi/6$ found	5
7	(i) 60 (ii) -6 (iii) 	2 1 1 1	M1 for $2^2 + 2^3 + 2^4 + 2^5$ o.e. Correct in both quadrants Through (0, 1) shown dep.	5
8	$r = 1/3$ s.o.i. $a = 54$ or ft $18 \div$ their r $S = \frac{a}{1-r}$ used with $-1 < r < 1$ $S = 81$ c.a.o.	2 M1 M1 A1	1 mark for $ar = 18$ and $ar^3 = 2$ s.o.i.	5
9	(i) 0.23 c.a.o. (ii) 0.1 or $1/10$ (iii) $4(3x + 2)$ or $12x + 8$ (iv) $[y =] 10^{3x+2}$ o.e.	1 1 1 1	10^{-1} not sufficient	4

Section B

10	i	$h = 120/x^2$ $A = 2x^2 + 4xh$ o.e. completion to given answer	B1 M1 A1	at least one interim step shown	3
	ii	$A' = 4x - 480/x^2$ o.e. $A'' = 4 + 960/x^3$	2 2	1 for kx^2 o.e. included ft their A' only if kx^2 seen ; 1 if one error	4
	iii	use of $A' = 0$ $x = \sqrt[3]{120}$ or 4.9(3..) <p>Test using A' or A'' to confirm minimum</p> <p>Substitution of their x in A</p> $A = 145.9$ to 146	M1 A1 T1 M1 A1	Dependent on previous M1	5
11	iA	$BC^2 = 348^2 + 302^2 - 2 \times 348 \times 302 \times \cos 72^\circ$ $BC = 383.86\dots$ $1033.86\dots$ [m] or ft 650 + their BC	M2 A1 1	M1 for recognisable attempt at Cosine Rule to 3 sf or more accept to 3 sf or more	4
	iB	$\frac{\sin B}{302} = \frac{\sin 72}{\text{their } BC}$ $B = 48.4\dots$ $355 - \text{their } B$ o.e. answer in range 306 to 307	M1 A1 M1 A1	Cosine Rule acceptable or Sine Rule to find C or 247 + their C	4
	ii	Arc length PQ = $\frac{224}{360} \times 2\pi \times 120$ o.e. or 469.1... to 3 sf or more $QP = 222.5\dots$ to 3 sf or more answer in range 690 to 692 [m]	M2 B1 A1	M1 for $\frac{136}{360} \times 2\pi \times 120$	4
12	iA	$x^4 = 8x$ (2, 16) c.a.o. $PQ = 16$ and completion to show $\frac{1}{2} \times 2 \times 16 = 16$	M1 A1 A1	NB answer 16 given	3
	iB	$x^5/5$ evaluating their integral at their co-ord of P and zero [or $32/5$ o.e.] 9.6 o.e.	M1 M1 A1	ft only if integral attempted, not for x^4 or differentiation c.a.o.	3
	iiA	$6x^2h^2 + 4xh^3 + h^4$	2	B1 for two terms correct.	2
	iiB	$4x^3 + 6x^2h + 4xh^2 + h^3$	2	B1 for three terms correct	2
	iiC	$4x^3$	1		1
	iiD	gradient of [tangent to] curve	1		1

4753 (C3) Methods for Advanced Mathematics


Section A

1 $y = (1 + 6x^2)^{1/3}$ $\Rightarrow \frac{dy}{dx} = \frac{1}{3}(1 + 6x^2)^{-2/3} \cdot 12x$ $= 4x(1 + 6x^2)^{-2/3}$	M1 B1 A1 A1 [4]	chain rule used $\frac{1}{3}u^{-2/3}$ $\times 12x$ cao (must resolve $1/3 \times 12$) Mark final answer
2 (i) $fg(x) = f(x - 2)$ $= (x - 2)^2$ $gf(x) = g(x^2) = x^2 - 2.$	M1 A1 A1 [3]	forming a composite function mark final answer If fg and gf the wrong way round, M1A0A0
(ii) 	B1ft B1ft [2]	fg – must have (2, 0) labelled (or inferable from scale). Condone no y-intercept, unless wrong gf – must have (0, -2) labelled (or inferable from scale) Condone no x-intercepts, unless wrong Allow ft only if fg and gf are correct but wrong way round.
3 (i) When $n = 1$, $10\,000 = A e^b$ when $n = 2$, $16\,000 = A e^{2b}$ $\Rightarrow \frac{16000}{10000} = \frac{Ae^{2b}}{Ae^b} = e^b$ $\Rightarrow e^b = 1.6$ $\Rightarrow b = \ln 1.6 = 0.470$ $A = 10000/1.6 = 6250.$	B1 B1 M1 E1 B1 B1 [6]	soi soi eliminating A (do not allow verification) SCB2 if initial 'B's are missing, and ratio of years = 1.6 $= e^b$ ln 1.6 or 0.47 or better (mark final answer) cao – allow recovery from inexact b's
(ii) When $n = 20$, $P = 6250 \times e^{0.470 \times 20}$ $= \text{£}75,550,000$	M1 A1 [2]	substituting $n = 20$ into their equation with their A and b Allow answers from £75 000 000 to £76 000 000.
4 (i) $5 = k/100 \Rightarrow k = 500^*$	E1 [1]	NB answer given
(ii) $\frac{dP}{dV} = -500V^{-2} = -\frac{500}{V^2}$	M1 A1 [2]	$(-1)V^{-2}$ o.e. – allow $-k/V^2$
(iii) $\frac{dP}{dt} = \frac{dP}{dV} \cdot \frac{dV}{dt}$ When $V = 100$, $dP/dV = -500/10000 = -0.05$ $dV/dt = 10$ $\Rightarrow dP/dt = -0.05 \times 10 = -0.5$ So P is decreasing at 0.5 Atm/s	M1 B1ft B1 A1 [4]	chain rule (any correct version) (soi) (soi) -0.5 cao

5(i) $p = 2, 2^p - 1 = 3$, prime $p = 3, 2^p - 1 = 7$, prime $p = 5, 2^p - 1 = 31$, prime $p = 7, 2^p - 1 = 127$, prime	M1 E1 [2]	Testing at least one prime testing all 4 primes (correctly) Must comment on answers being prime (allow ticks) Testing $p = 1$ is E0
(ii) $23 \times 89 = 2047 = 2^{11} - 1$ 11 is prime, 2047 is not So statement is false.	M1 E1 [2]	$2^{11} - 1$ must state or imply that 11 is prime ($p = 11$ is sufficient)
6 (i) $e^{2y} = x^2 + y$ $\Rightarrow 2e^{2y} \frac{dy}{dx} = 2x + \frac{dy}{dx}$ $\Rightarrow (2e^{2y} - 1) \frac{dy}{dx} = 2x$ $\Rightarrow \frac{dy}{dx} = \frac{2x}{2e^{2y} - 1} *$	M1 A1 M1 E1 [4]	Implicit differentiation – allow one slip (but with dy/dx both sides) collecting terms
(ii) Gradient is infinite when $2e^{2y} - 1 = 0$ $\Rightarrow e^{2y} = \frac{1}{2}$ $\Rightarrow 2y = \ln \frac{1}{2}$ $\Rightarrow y = \frac{1}{2} \ln \frac{1}{2} = -0.347$ (3 s.f.) $x^2 = e^{2y} - y = \frac{1}{2} - (-0.347)$ $= 0.8465$ $\Rightarrow x = 0.920$	M1 A1 M1 A1 [4]	must be to 3 s.f. substituting their y and solving for x cao – must be to 3 s.f., but penalise accuracy once only.

Section B

<p>7(i) $y = 2x \ln(1+x)$ $\Rightarrow \frac{dy}{dx} = \frac{2x}{1+x} + 2 \ln(1+x)$ When $x = 0$, $dy/dx = 0 + 2 \ln 1 = 0$ \Rightarrow origin is a stationary point.</p>	M1 B1 A1 E1 [4]	product rule $d/dx(\ln(1+x)) = 1/(1+x)$ soi www (i.e. from correct derivative)
<p>(ii) $\frac{d^2y}{dx^2} = \frac{(1+x) \cdot 2 - 2x \cdot 1}{(1+x)^2} + \frac{2}{1+x}$ $= \frac{2}{(1+x)^2} + \frac{2}{1+x}$ When $x = 0$, $d^2y/dx^2 = 2 + 2 = 4 > 0$ $\Rightarrow (0, 0)$ is a min point</p>	M1 A1ft A1 M1 E1 [5]	Quotient or product rule on their $2x/(1+x)$ correctly applied to their $2x/(1+x)$ o.e., e.g. $\frac{4+2x}{(1+x)^2}$ cao substituting $x = 0$ into their d^2y/dx^2 www – dep previous A1
<p>(iii) Let $u = 1+x \Rightarrow du = dx$ $\Rightarrow \int \frac{x^2}{1+x} dx = \int \frac{(u-1)^2}{u} du$ $= \int \frac{(u^2 - 2u + 1)}{u} du$ $= \int (u - 2 + \frac{1}{u}) du$ * $\Rightarrow \int_0^1 \frac{x^2}{1+x} dx = \int_1^2 (u - 2 + \frac{1}{u}) du$ $= \left[\frac{1}{2}u^2 - 2u + \ln u \right]_1^2$ $= 2 - 4 + \ln 2 - (\frac{1}{2} - 2 + \ln 1)$ $= \ln 2 - \frac{1}{2}$</p>	M1 E1 B1 B1 M1 A1 [6]	$\frac{(u-1)^2}{u}$ www (but condone du omitted except in final answer) changing limits (or substituting back for x and using 0 and 1) $\left[\frac{1}{2}u^2 - 2u + \ln u \right]$ substituting limits (consistent with u or x) cao
<p>(iv) $A = \int_0^1 2x \ln(1+x) dx$ Parts: $u = \ln(1+x)$, $du/dx = 1/(1+x)$ $dv/dx = 2x \Rightarrow v = x^2$ $= \left[x^2 \ln(1+x) \right]_0^1 - \int_0^1 \frac{x^2}{1+x} dx$ $= \ln 2 - \ln 2 + \frac{1}{2} = \frac{1}{2}$</p>	M1 A1 M1 A1 [4]	soi substituting their $\ln 2 - \frac{1}{2}$ for $\int_0^1 \frac{x^2}{1+x} dx$ cao

<p>8 (i) Stretch in x-direction s.f. $\frac{1}{2}$ translation in y-direction 1 unit up</p>	<p>M1 A1 M1 A1 [4]</p>	<p>(in either order) – allow ‘contraction’ dep ‘stretch’ allow ‘move’, ‘shift’, etc – direction can be inferred from $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ or $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ dep ‘translation’. $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ alone is M1 A0</p>
<p>(ii) $A = \int_{-\pi/4}^{\pi/4} (1 + \sin 2x) dx$ $= \left[x - \frac{1}{2} \cos 2x \right]_{-\pi/4}^{\pi/4}$ $= \pi/4 - \frac{1}{2} \cos \pi/2 + \pi/4 + \frac{1}{2} \cos (-\pi/2)$ $= \pi/2$</p>	<p>M1 B1 M1 A1 [4]</p>	<p>correct integral and limits. Condone dx missing; limits may be implied from subsequent working. substituting their limits (if zero lower limit used, must show evidence of substitution) or 1.57 or better – cao (www)</p>
<p>(iii) $y = 1 + \sin 2x$ $\Rightarrow dy/dx = 2\cos 2x$ When $x = 0$, $dy/dx = 2$ So gradient at $(0, 1)$ on $f(x)$ is 2 \Rightarrow gradient at $(1, 0)$ on $f^{-1}(x) = \frac{1}{2}$</p>	<p>M1 A1 A1ft B1ft [4]</p>	<p>differentiating – allow 1 error (but not $x + 2\cos 2x$) If 1, then must show evidence of using reciprocal, e.g. 1/1</p>
<p>(iv) Domain is $0 \leq x \leq 2$.</p> 	<p>B1 M1 A1 [3]</p>	<p>Allow 0 to 2, but not $0 < x < 2$ or y instead of x clear attempt to reflect in $y = x$ correct domain indicated (0 to 2), and reasonable shape</p>
<p>(v) $y = 1 + \sin 2x$ $x \leftrightarrow y$ $x = 1 + \sin 2y$ $\Rightarrow \sin 2y = x - 1$ $\Rightarrow 2y = \arcsin(x - 1)$ $\Rightarrow y = \frac{1}{2} \arcsin(x - 1)$</p>	<p>M1 A1 [2]</p>	<p>or $\sin 2x = y - 1$ cao</p>

4754 (C4) Applications of Advanced Mathematics


Section A

<p>1 $3 \cos \theta + 4 \sin \theta = R \cos(\theta - \alpha)$ $= R(\cos \theta \cos \alpha + \sin \theta \sin \alpha)$ $\Rightarrow R \cos \alpha = 3, R \sin \alpha = 4$ $\Rightarrow R^2 = 3^2 + 4^2 = 25 \Rightarrow R = 5$ $\tan \alpha = 4/3 \Rightarrow \alpha = 0.9273$</p> <p>$5 \cos(\theta - 0.9273) = 2$ $\Rightarrow \cos(\theta - 0.9273) = 2/5$ $\theta - 0.9273 = 1.1593, -1.1593$ $\Rightarrow \theta = 2.087, -0.232$</p>	<p>M1 B1 M1A1</p> <p>M1 A1 A1 [7]</p>	<p>$R = 5$ cwo</p> <p>and no others in the range</p>
<p>2(i) $(1-2x)^{-\frac{1}{2}} = 1 - \frac{1}{2}(-2x) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!}(-2x)^2 + \dots$ $= 1 + x + \frac{3}{2}x^2 + \dots$ Valid for $-1 < -2x < 1 \Rightarrow -\frac{1}{2} < x < \frac{1}{2}$</p>	<p>M1 A1</p> <p>A1 M1 A1 [5]</p>	<p>binomial expansion with $p = -\frac{1}{2}$ correct expression</p> <p>cao</p>
<p>(ii) $\frac{1+2x}{\sqrt{1-2x}} = (1+2x)(1+x+\frac{3}{2}x^2+\dots)$ $= 1+x+\frac{3}{2}x^2+2x+2x^2+\dots$ $= 1+3x+\frac{7}{2}x^2+\dots$</p>	<p>M1</p> <p>A1ft</p> <p>A1 [3]</p>	<p>substituting their $1+x+\frac{3}{2}x^2+\dots$ and expanding</p> <p>cao</p>
<p>3 $V = \int_1^2 \pi x^2 dy$ $y = 1 + x^2 \Rightarrow x^2 = y - 1$ $\Rightarrow V = \int_1^2 \pi(y-1) dy$ $= \pi \left[\frac{1}{2}y^2 - y \right]_1^2$ $= \pi(2 - 2 - \frac{1}{2} + 1)$ $= \frac{1}{2} \pi$</p>	<p>B1</p> <p>M1</p> <p>B1</p> <p>M1 A1 [5]</p>	<p>$\left[\frac{1}{2}y^2 - y \right]$</p> <p>substituting limits into integrand</p>

<p>4(i) $\sin(\theta + 45^\circ) = \cos \theta$ $\Rightarrow \sin \theta \cos 45 + \cos \theta \sin 45 = \cos \theta$ $\Rightarrow (1/\sqrt{2}) \sin \theta + (1/\sqrt{2}) \cos \theta = \cos \theta$ $\Rightarrow \sin \theta + \cos \theta = \sqrt{2} \cos \theta$ $\Rightarrow \sin \theta = (\sqrt{2} - 1) \cos \theta$ $\Rightarrow \frac{\sin \theta}{\cos \theta} = \tan \theta = \sqrt{2} - 1 *$</p>	<p>M1 B1 A1 M1 E1 [5]</p>	<p>compound angle formula $\sin 45 = 1/\sqrt{2}$, $\cos 45 = 1/\sqrt{2}$ collecting terms</p>
<p>(ii) $\tan \theta = \sqrt{2} - 1$ $\Rightarrow \theta = 22.5^\circ,$ 202.5°</p>	<p>B1 B1 [2]</p>	<p>and no others in the range</p>
<p>5 $\frac{4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{x^2 + 4}$ $= \frac{A(x^2 + 4) + (Bx + C)x}{x(x^2 + 4)}$ $\Rightarrow 4 = A(x^2 + 4) + (Bx + C)x$ $x = 0 \Rightarrow 4 = 4A \Rightarrow A = 1$ coefft of x^2: $0 = A + B \Rightarrow B = -1$ coeffts of x: $0 = C$ $\Rightarrow \frac{4}{x(x^2 + 4)} = \frac{1}{x} - \frac{x}{x^2 + 4}$</p>	<p>M1 M1 B1 DM1 A1 A1 [6]</p>	<p>correct partial fractions A=1 Substitution or equating coeffts B= -1 C= 0</p>
<p>6 $\operatorname{cosec} \theta = 3$ $\Rightarrow \sin \theta = 1/3$ $\Rightarrow \theta = 19.47^\circ,$ 160.53°</p>	<p>M1 A1 A1 [3]</p>	<p>and no others in the range</p>

Section B


7(i) $\overline{CD} = \begin{pmatrix} -6 \\ 6 \\ 24 \end{pmatrix}$ $\overline{CB} = \begin{pmatrix} 0 \\ 20 \\ 0 \end{pmatrix}$	B1 B1 [2]	
(ii) $\sqrt{(-6)^2 + 6^2 + 24^2}$ = 25.46 cm	M1 A1 [2]	
(iii) $\overline{CD} \cdot \begin{pmatrix} 4 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} -6 \\ 6 \\ 24 \end{pmatrix} \cdot \begin{pmatrix} 4 \\ 0 \\ 1 \end{pmatrix} = -24 + 0 + 24 = 0$ $\overline{CB} \cdot \begin{pmatrix} 4 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 20 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 4 \\ 0 \\ 1 \end{pmatrix} = 0 + 0 + 0 = 0$ \Rightarrow plane BCDE is $4x + z = c$ At C (say) $4 \times 15 + 0 = c \Rightarrow c = 60$ \Rightarrow plane BCDE is $4x + z = 60$	M1 A1 B1 M1 A1 [5]	using scalar product or other equivalent methods
(iv) OG: $\mathbf{r} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 3 \\ 6 \\ 24 \end{pmatrix}$ AF: $\mathbf{r} = \begin{pmatrix} 0 \\ 20 \\ 0 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ -6 \\ 24 \end{pmatrix}$ At (5, 10, 40), $3\lambda = 5 \Rightarrow \lambda = 5/3$ $\Rightarrow 6\lambda = 10, 24\lambda = 40$, so consistent. At (5, 10, 40), $3\mu = 5 \Rightarrow \mu = 5/3$ $\Rightarrow 20 - 6\mu = 10, 24\mu = 40$, so consistent. So lines meet at (5, 10, 40)*	B1 B1 M1 E1 E1 [5]	evaluating parameter and checking consistency. [or other methods, e.g. solving]
(v) $h=40$ POABC: $V = 1/3 \times 20 \times 15 \times 40$ $= 4000 \text{ cm}^3$. PDEFG: $V = 1/3 \times 8 \times 6 \times (40-24)$ $= 256 \text{ cm}^3$ \Rightarrow vol of ornament = $4000 - 256 = 3744 \text{ cm}^3$	B1 M1 A1 A1 [4]	soi $1/3 \times w \times d \times h$ used for either –condone one error both volumes correct cao

<p>8(i) $\cos \theta = \frac{x}{k}, \sin \theta = \frac{2y}{k}$ $\cos^2 \theta + \sin^2 \theta = 1$ $\Rightarrow \left(\frac{x}{k}\right)^2 + \left(\frac{2y}{k}\right)^2 = 1$ $\Rightarrow \frac{x^2}{k^2} + \frac{4y^2}{k^2} = 1$ $\Rightarrow x^2 + 4y^2 = k^2 *$</p>	<p>M1 M1 E1 [3]</p>	<p>Used substitution</p>
<p>(ii) $\frac{dx}{d\theta} = -k \sin \theta, \frac{dy}{d\theta} = \frac{1}{2}k \cos \theta$ $\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = -\frac{\frac{1}{2}k \cos \theta}{k \sin \theta}$ $= -\frac{1}{2} \cot \theta$ $-\frac{x}{4y} = -\frac{2k \cos \theta}{4k \sin \theta} = -\frac{1}{2} \cot \theta = \frac{dy}{dx}$</p>	<p>M1 A1 E1</p>	<p>oe</p>
<p>or, by differentiating implicitly $2x + 8y \, dy/dx = 0$ $\Rightarrow \, dy/dx = -2x/8y = -x/4y*$</p>	<p>M1 A1 E1 [3]</p>	
<p>(iii) $k = 2$</p>	<p>B1 [1]</p>	
<p>(iv)</p> 	<p>B1 B1 B1 [3]</p>	<p>1 correct curve –shape and position 2 or more curves correct shape- in concentric form all 3 curves correct</p>
<p>(v) grad of stream path = $-1/\text{grad of contour}$ $\Rightarrow \frac{dy}{dx} = -\frac{1}{(-x/4y)} = \frac{4y}{x} *$</p>	<p>M1 E1 [2]</p>	
<p>(vi) $\frac{dy}{dx} = \frac{4y}{x} \Rightarrow \int \frac{dy}{y} = \int \frac{4dx}{x}$ $\Rightarrow \ln y = 4 \ln x + c = \ln e^c x^4$ $\Rightarrow y = Ax^4$ where $A = e^c$.</p> <p>When $x = 2, y = 1 \Rightarrow 1 = 16A \Rightarrow A = 1/16$ $\Rightarrow y = x^4/16 *$</p>	<p>M1 A1 M1 M1 A1 E1 [6]</p>	<p>Separating variables $\ln y = 4 \ln x (+c)$ antilogging correctly (at any stage) substituting $x = 2, y = 1$ evaluating a correct constant www</p>

Paper B Comprehension 4754 (C4)




1	4, 1, 5, 6, 11, 17	B1 B1	for 11 and 17 for 1 and 4
2	Even, odd, odd, even, odd, odd recurs 100 th term is therefore even	M1 A1	for reason www
3	$\phi^6 = (3\phi + 2) + (5\phi + 3) = 8\phi + 5$	B1	
4	$1 - EH = 1 - CG = 1 - (\phi - 1)$ $= 2 - \phi = 2 - \left(\frac{1 + \sqrt{5}}{2}\right)$ $= \frac{3 - \sqrt{5}}{2}$	M1 A1 A1	oe
5	(i) Gradients $-\frac{1}{\phi}$ and $\frac{1}{\phi - 1}$ (ii) Product of gradients: $-\frac{1}{\phi} \times \frac{1}{\phi - 1} = -\frac{1}{\phi^2 - \phi}$ $= -\frac{1}{1} = -1$	B1 B1 M1 E1	
6	$\frac{\phi + 1}{2\phi - 1} = \frac{\frac{1 + \sqrt{5}}{2} + 1}{1 + \sqrt{5} - 1}$ $= \frac{3 + \sqrt{5}}{2\sqrt{5}}$ $= \frac{(3 + \sqrt{5})\sqrt{5}}{2\sqrt{5} \times \sqrt{5}} = \frac{3\sqrt{5} + 5}{10}$	M1 A1 E1	
7	$a + (a + d) = a + 2d \Rightarrow a = d$ $(a + d) + (a + 2d) = a + 3d \Rightarrow a = 0$ $a = d = 0$ *	M1 M1 E1 [18]	

4755 (FP1) Further Concepts for Advanced Mathematics

Qu	Answer	Mark	Comment
Section A			
1(i)	$\mathbf{BA} = \begin{pmatrix} 3 & 1 \\ -2 & 4 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ 0 & 3 \end{pmatrix} = \begin{pmatrix} 6 & 0 \\ -4 & 14 \end{pmatrix}$	M1 A1 [2]	Attempt to multiply c.a.o.
1(ii)	$\det \mathbf{BA} = (6 \times 14) - (-4 \times 0) = 84$ $3 \times 84 = 252$ square units	M1 A1 A1(ft) [3]	Attempt to calculate any determinant c.a.o. Correct area
2(i)	$\alpha^2 = (-3 + 4j)(-3 + 4j) = (-7 - 24j)$	M1 A1 [2]	Attempt to multiply with use of $j^2 = -1$ c.a.o.
2(ii)	$ \alpha = 5$ $\arg \alpha = \pi - \arctan \frac{4}{3} = 2.21$ (2d.p.) (or 126.87°) $\alpha = 5(\cos 2.21 + j \sin 2.21)$	B1 B1 B1(ft) [3]	Accept 2.2 or 127° Accept degrees and (r, θ) form s.c. lose 1 mark only if α^2 used throughout (ii)
3(i)	$3^3 + 3^2 - 7 \times 3 - 15 = 0$ $z^3 + z^2 - 7z - 15 = (z - 3)(z^2 + 4z + 5)$ $z = \frac{-4 \pm \sqrt{16 - 20}}{2} = -2 \pm j$ So other roots are $-2 + j$ and $-2 - j$	B1 M1 A1 M1 A1 [5]	Showing 3 satisfies the equation (may be implied) Valid attempt to factorise Correct quadratic factor Use of quadratic formula, or other valid method One mark for both c.a.o.
3(ii)		B2 [2]	Minus 1 for each error ft provided conjugate imaginary roots

4	$\sum_{r=1}^n [(r+1)(r-2)] = \sum_{r=1}^n r^2 - \sum_{r=1}^n r - 2n$ $= \frac{1}{6}n(n+1)(2n+1) - \frac{1}{2}n(n+1) - 2n$ $= \frac{1}{6}n[(n+1)(2n+1) - 3(n+1) - 12]$ $= \frac{1}{6}n(2n^2 + 3n + 1 - 3n - 3 - 12)$ $= \frac{1}{6}n(2n^2 - 14)$ $= \frac{1}{3}n(n^2 - 7)$	M1 A2 M1 M1 A1 [6]	Attempt to split sum up Minus one each error Attempt to factorise Collecting terms All correct
5(i) 5(ii)	$p = -3, r = 7$ $q = \alpha\beta + \alpha\gamma + \beta\gamma$ $\alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \alpha\gamma + \beta\gamma)$ $= (\alpha + \beta + \gamma)^2 - 2q$ $\Rightarrow 13 = 3^2 - 2q$ $\Rightarrow q = -2$	B2 [2] B1 M1 A1 [3]	One mark for each s.c. B1 if b and d used instead of p and r Attempt to find q using $\alpha^2 + \beta^2 + \gamma^2$ and $\alpha + \beta + \gamma$, but not $\alpha\beta\gamma$ c.a.o.
6(i) 6(ii)	$a_2 = 7 \times 7 - 3 = 46$ $a_3 = 7 \times 46 - 3 = 319$ <p>When $n = 1$, $\frac{13 \times 7^0 + 1}{2} = 7$, so true for $n = 1$</p> <p>Assume true for $n = k$</p> $a_k = \frac{13 \times 7^{k-1} + 1}{2}$ $\Rightarrow a_{k+1} = 7 \times \frac{13 \times 7^{k-1} + 1}{2} - 3$ $= \frac{13 \times 7^k + 7}{2} - 3$ $= \frac{13 \times 7^k + 7 - 6}{2}$ $= \frac{13 \times 7^k + 1}{2}$ <p>But this is the given result with $k + 1$ replacing k. Therefore if it is true for k it is true for $k + 1$. Since it is true for $k = 1$, it is true for $k = 1, 2, 3$ and so true for all positive integers.</p>	M1 A1 [2] B1 E1 M1 A1 E1 E1 [6]	Use of inductive definition c.a.o. Correct use of part (i) (may be implied) Assuming true for k Attempt to use $a_{k+1} = 7a_k - 3$ Correct simplification Dependent on A1 and previous E1 Dependent on B1 and previous E1


Section A Total: 36

Section B			
7(i)	$(1, 0)$ and $(0, \frac{1}{18})$	B1 B1 [2]	
7(ii)	$x = 2, x = -3, x = \frac{-3}{2}, y = 0$	B4 [4]	Minus 1 for each error
7(iii)		B1 B1 [2]	Correct approaches to vertical asymptotes Through clearly marked $(1, 0)$ and $(0, \frac{1}{18})$
7(iv)	$x < -3, x > 2$ $\frac{-3}{2} < x \leq 1$	B1 B2 [3]	B1 for $\frac{-3}{2} < x < 1$, or $\frac{-3}{2} \leq x \leq 1$
8(i)		B3 B3 [6]	Circle, B1; radius 2, B1; centre 3j, B1 Half line, B1; from -1, B1; $\frac{\pi}{4}$ to x-axis, B1
8(ii)	 <p>Sketch should clearly show the radius and centre of the circle and the starting point and angle of the half-line.</p>	B2(ft) [2] M1	Correct region between their circle and half line indicated s.c. B1 for interior of circle Tangent from origin to circle
8(iii)	$\arg z = \frac{\pi}{2} - \arcsin \frac{2}{3} = 0.84$ (2d.p.)	A1(ft) M1 A1 [4]	Correct point placed by eye where tangent from origin meets circle Attempt to use right angled triangle c.a.o. Accept 48.20° (2d.p.)

9(i)	$(-3, -3)$	B1 [1]	
9(ii)	(x, x)	B1 B1 [2]	
9(iii)	$\begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix}$	B3 [3]	Minus 1 each error to min of 0
9(iv)	Rotation through $\frac{\pi}{2}$ anticlockwise about the origin	B1 B1 [2]	Rotation and angle (accept 90°) Centre and sense
9(v)	$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \times \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 1 & 0 \end{pmatrix}$	M1 A1 [2]	Attempt to multiply using their T in correct order c.a.o.
9(vi)	$\begin{pmatrix} -1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -x \\ x \end{pmatrix}$ So $(-x, x)$ Line $y = -x$	M1 A1(ft) A1 [3]	May be implied c.a.o. from correct matrix

4756 (FP2) Further Methods for Advanced Mathematics

1(a)	Area is $\int_0^{\pi} \frac{1}{2} a^2 (1 - \cos 2\theta)^2 d\theta$ $= \int_0^{\pi} \frac{1}{2} a^2 (1 - 2 \cos 2\theta + \frac{1}{2} (1 + \cos 4\theta)) d\theta$ $= \frac{1}{2} a^2 \left[\frac{3}{2} \theta - \sin 2\theta + \frac{1}{8} \sin 4\theta \right]_0^{\pi}$ $= \frac{3}{4} \pi a^2$	M1 A1 B1 B1B1B1 ft A1 7	For $\int (1 - \cos 2\theta)^2 d\theta$ Correct integral expression including limits (may be implied by later work) For $\cos^2 2\theta = \frac{1}{2} (1 + \cos 4\theta)$ Integrating $a + b \cos 2\theta + c \cos 4\theta$ <i>[Max B2 if answer incorrect and no mark has previously been lost]</i>
(b)(i)	$f'(x) = \frac{1}{1 + (\sqrt{3} + x)^2}$ $f''(x) = \frac{-2(\sqrt{3} + x)}{(1 + (\sqrt{3} + x)^2)^2}$	M1 A1 M1 A1 4	Applying $\frac{d}{du} \arctan u = \frac{1}{1 + u^2}$ or $\frac{dy}{dx} = \frac{1}{\sec^2 y}$ Applying chain (or quotient) rule
(ii)	$f(0) = \frac{1}{3} \pi$ $f'(0) = \frac{1}{4}, f''(0) = -\frac{1}{8} \sqrt{3}$ $\arctan(\sqrt{3} + x) = \frac{1}{3} \pi + \frac{1}{4} x - \frac{1}{16} \sqrt{3} x^2 + \dots$	B1 M1 A1A1 ft 4	Stated; or appearing in series <i>Accept 1.05</i> Evaluating $f'(0)$ or $f''(0)$ For $\frac{1}{4} x$ and $-\frac{1}{16} \sqrt{3} x^2$ <i>ft provided coefficients are non-zero</i>
(iii)	$\int_{-h}^h (\frac{1}{3} \pi x + \frac{1}{4} x^2 - \frac{1}{16} \sqrt{3} x^3 + \dots) dx$ $= \left[\frac{1}{6} \pi x^2 + \frac{1}{12} x^3 - \frac{1}{64} \sqrt{3} x^4 + \dots \right]_{-h}^h$ $\approx (\frac{1}{6} \pi h^2 + \frac{1}{12} h^3 - \frac{1}{64} \sqrt{3} h^4)$ $\quad - (\frac{1}{6} \pi h^2 - \frac{1}{12} h^3 - \frac{1}{64} \sqrt{3} h^4)$ $= \frac{1}{6} h^3$	M1 A1 ft A1 ag 3	Integrating (award if x is missed) for $\frac{1}{12} x^3$ Allow ft from $a + \frac{1}{4} x + cx^2$ provided that $a \neq 0$ Condone a proof which neglects h^4

2(a)	<p>4th roots of $16j = 16e^{\frac{1}{2}\pi j}$ are $re^{j\theta}$ where</p> $r = 2$ $\theta = \frac{1}{8}\pi$ $\theta = \frac{\pi}{8} + \frac{2k\pi}{4}$ $\theta = -\frac{7}{8}\pi, -\frac{3}{8}\pi, \frac{5}{8}\pi$ 	B1 B1 M1 A1 M1 A1 6	<p>Accept $16^{\frac{1}{4}}$</p> <p>Implied by at least two correct (ft) further values or stating $k = -2, -1, (0), 1$</p> <p>Points at vertices of a square centre O or 3 correct points (ft) or 1 point in each quadrant</p>
(b)(i)	$(1 - 2e^{j\theta})(1 - 2e^{-j\theta}) = 1 - 2e^{j\theta} - 2e^{-j\theta} + 4$ $= 5 - 2(e^{j\theta} + e^{-j\theta})$ $= 5 - 4\cos\theta$ <p>OR</p> $(1 - 2\cos\theta - 2j\sin\theta)(1 - 2\cos\theta + 2j\sin\theta)$ $= (1 - 2\cos\theta)^2 + 4\sin^2\theta$ $= 1 - 4\cos\theta + 4(\cos^2\theta + \sin^2\theta)$ $= 5 - 4\cos\theta$	M1 A1 A1 ag 3	<p>For $e^{j\theta}e^{-j\theta} = 1$</p>
(ii)	$C + jS = 2e^{j\theta} + 4e^{2j\theta} + 8e^{3j\theta} + \dots + 2^n e^{nj\theta}$ $= \frac{2e^{j\theta}(1 - (2e^{j\theta})^n)}{1 - 2e^{j\theta}}$ $= \frac{2e^{j\theta}(1 - 2^n e^{nj\theta})(1 - 2e^{-j\theta})}{(1 - 2e^{j\theta})(1 - 2e^{-j\theta})}$ $= \frac{2e^{j\theta} - 4 - 2^{n+1}e^{(n+1)j\theta} + 2^{n+2}e^{nj\theta}}{5 - 4\cos\theta}$ $C = \frac{2\cos\theta - 4 - 2^{n+1}\cos(n+1)\theta + 2^{n+2}\cos n\theta}{5 - 4\cos\theta}$ $S = \frac{2\sin\theta - 2^{n+1}\sin(n+1)\theta + 2^{n+2}\sin n\theta}{5 - 4\cos\theta}$	M1 M1 A1 M1 A2 M1 A1 ag A1 9	<p>Obtaining a geometric series</p> <p>Summing (M0 for sum to infinity)</p> <p>Give A1 for two correct terms in numerator</p> <p>Equating real (or imaginary) parts</p>


3 (i)	Characteristic equation is $(7 - \lambda)(-1 - \lambda) + 12 = 0$ $\lambda^2 - 6\lambda + 5 = 0$ $\lambda = 1, 5$ When $\lambda = 1$, $\begin{pmatrix} 7 & 3 \\ -4 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix}$ $7x + 3y = x$ $-4x - y = y$ $y = -2x$, eigenvector is $\begin{pmatrix} 1 \\ -2 \end{pmatrix}$ When $\lambda = 5$, $\begin{pmatrix} 7 & 3 \\ -4 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 5 \begin{pmatrix} x \\ y \end{pmatrix}$ $7x + 3y = 5x$ $-4x - y = 5y$ $y = -\frac{2}{3}x$, eigenvector is $\begin{pmatrix} 3 \\ -2 \end{pmatrix}$	M1 A1A1 M1 M1 A1 M1 A1	$\text{or } \begin{pmatrix} 6 & 3 \\ -4 & -2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ <i>can be awarded for either eigenvalue</i> Equation relating x and y or any (non-zero) multiple SR $(\mathbf{M} - \lambda\mathbf{I})\mathbf{x} = \lambda\mathbf{x}$ can earn M1A1A1M0M1A0M1A0
(ii)	$\mathbf{P} = \begin{pmatrix} 1 & 3 \\ -2 & -2 \end{pmatrix}$ $\mathbf{D} = \begin{pmatrix} 1 & 0 \\ 0 & 5 \end{pmatrix}$	B1 ft B1 ft	B0 if \mathbf{P} is singular For B2, the order must be consistent

8

2


(iii) $\mathbf{M} = \mathbf{PDP}^{-1}$ $\mathbf{M}^n = \mathbf{PD}^n \mathbf{P}^{-1}$ $= \mathbf{P} \begin{pmatrix} 1 & 0 \\ 0 & 5^n \end{pmatrix} \mathbf{P}^{-1}$ $= \begin{pmatrix} 1 & 3 \\ -2 & -2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 5^n \end{pmatrix} \frac{1}{4} \begin{pmatrix} -2 & -3 \\ 2 & 1 \end{pmatrix}$ $= \begin{pmatrix} 1 & 3 \times 5^n \\ -2 & -2 \times 5^n \end{pmatrix} \frac{1}{4} \begin{pmatrix} -2 & -3 \\ 2 & 1 \end{pmatrix}$ $= \frac{1}{4} \begin{pmatrix} -2 + 6 \times 5^n & -3 + 3 \times 5^n \\ 4 - 4 \times 5^n & 6 - 2 \times 5^n \end{pmatrix}$ $a = -\frac{1}{2} + \frac{3}{2} \times 5^n$ $b = -\frac{3}{4} + \frac{3}{4} \times 5^n$ $c = 1 - 5^n$ $d = \frac{3}{2} - \frac{1}{2} \times 5^n$	M1	May be implied
	M1	
	A1 ft	Dependent on M1M1
	B1 ft	For \mathbf{P}^{-1}
	M1	Obtaining at least one element in a product of three matrices
	A1 ag	
A2	Give A1 for one of b, c, d correct	
	8	SR If $\mathbf{M}^n = \mathbf{P}^{-1} \mathbf{D}^n \mathbf{P}$ is used, max marks are M0M1A0B1M1A0A1 (d should be correct)
		SR If their \mathbf{P} is singular, max marks are M1M1A1B0M0

<p>OR Gradient $g = 6 \cosh x - 2 \cosh 2x$ B1</p> <p>$g' = 6 \sinh x - 4 \sinh 2x = 2 \sinh x(3 - 4 \cosh x)$</p> <p>$= 0$ when $x = 0$ (only) M1</p> <p>$g'' = 6 \cosh x - 8 \cosh 2x = -2$ when $x = 0$ M1</p> <p>Max value $g = 4$ when $x = 0$</p> <p>So g is never equal to 5 A1</p>		<p>Final A1 requires a complete proof showing this is the only turning point</p>
---	--	--

5 (i)	$\lambda = -1$ $\lambda = 0$ $\lambda = 1$  <div style="display: flex; justify-content: space-around;"> cusps loop </div>	B1B1B1 B1B1	 5 Two different features (cusp, loop, asymptote) correctly identified
(ii)	$x = 1$	B1	1
(iii)	Intersects itself when $y = 0$ $t = (\pm)\sqrt{\lambda}$ $\left(\frac{\lambda}{1+\lambda}, 0\right)$	M1 A1 A1	3
(iv)	$\frac{dy}{dt} = 3t^2 - \lambda = 0$ $t = \pm\sqrt{\frac{\lambda}{3}}$ $x = \frac{\lambda/3}{1+\lambda/3} = \frac{\lambda}{3+\lambda}$ $y = \pm\left(\left(\frac{\lambda}{3}\right)^{3/2} - \lambda\left(\frac{\lambda}{3}\right)^{1/2}\right)$ $= \pm\lambda^{3/2}\left(\frac{1}{3\sqrt{3}} - \frac{1}{\sqrt{3}}\right) = \pm\lambda^{3/2}\left(-\frac{2}{3\sqrt{3}}\right)$ $= \pm\sqrt{\frac{4\lambda^3}{27}}$	M1 A1 ag M1 A1 ag	 One value sufficient 4
(v)	From asymptote, $a = 8$ From intersection point, $\frac{a\lambda}{1+\lambda} = 2$ $\lambda = \frac{1}{3}$ From maximum point, $b\sqrt{\frac{4\lambda^3}{27}} = 2$ $b = 27$	B1 M1 A1 M1 A1	 5



4758

Differential Equations

<p>1(i)</p> $\alpha^2 + 2\alpha + 1 = 0$ <p>)</p> $\alpha = -1 \text{ (repeated)}$ <p>CF $y = (A + Bt)e^{-t}$</p> <p>PI $y = a$</p> <p>in DE $\Rightarrow y = 2$</p> $y = 2 + (A + Bt)e^{-t}$ $t = 0, y = 0 \Rightarrow 0 = 2 + A \Rightarrow A = -2$ $\dot{y} = (B - A - Bt)e^{-t}$ $t = 0, \dot{y} = 0 \Rightarrow 0 = B - A \Rightarrow B = -2$ $y = 2 - (2 + 2t)e^{-t}$	<p>M1 Auxiliary equation</p> <p>A1</p> <p>F1 CF for their roots</p> <p>B1 Constant PI</p> <p>B1 PI correct</p> <p>F1 Their PI + CF (with two arbitrary constants)</p> <p>M1 Condition on y</p> <p>M1 Differentiate (product rule)</p> <p>M1 Condition on \dot{y}</p> <p>A1</p>	10
<p>(ii)</p> <p>Both terms in CF hence will give zero if substituted in LHS</p> <p>PI $y = bt^2 e^{-t}$</p> $\dot{y} = (2bt - bt^2)e^{-t}, \ddot{y} = (2b - 4bt + bt^2)e^{-t}$ <p>in DE $\Rightarrow (2b - 4bt + bt^2 + 2(2bt - bt^2) + bt^2)e^{-t} = e^{-t}$</p> $\Rightarrow b = \frac{1}{2}$ $y = (C + Dt + \frac{1}{2}t^2)e^{-t}$ $t = 0, y = 0 \Rightarrow 0 = C$ $\dot{y} = (D + t - C - Dt - \frac{1}{2}t^2)e^{-t}$ $t = 0, \dot{y} = 0 \Rightarrow 0 = D - C \Rightarrow D = 0$ $y = \frac{1}{2}t^2 e^{-t}$	<p>E1</p> <p>B1</p> <p>M1 Differentiate twice and substitute</p> <p>A1 PI correct</p> <p>F1 Their PI + CF (with two arbitrary constants)</p> <p>M1 Condition on y</p> <p>M1 Condition on \dot{y}</p> <p>A1</p>	8
<p>(iii)</p> $t > 0 \Rightarrow \frac{1}{2}t^2 > 0 \text{ and } e^{-t} > 0 \Rightarrow y > 0$ $\dot{y} = (t - \frac{1}{2}t^2)e^{-t} \text{ so } \dot{y} = 0 \Leftrightarrow t - \frac{1}{2}t^2 = 0 \Leftrightarrow t = 0 \text{ or } 2$ <p>Maximum at $t = 2, y = 2e^{-2}$</p> 	<p>E1</p> <p>M1 Solve $\dot{y} = 0$</p> <p>A1 Maximum value of y</p> <p>B1 Starts at origin</p> <p>B1 Maximum at their value of y</p> <p>B1 $y > 0$</p>	6


2(i)	$\frac{dv}{dt} + \frac{3}{1+t}v = g - \frac{3}{1+t}$ $I = \exp\left(\int \frac{3}{1+t} dt\right) = e^{3\ln(1+t)} = (1+t)^3$ $(1+t)^3 \frac{dv}{dt} + 3(1+t)^2 v = g(1+t)^3 - 3(1+t)^2$ $\frac{d}{dt}\left((1+t)^3 v\right) = g(1+t)^3 - 3(1+t)^2$ $(1+t)^3 v = \int \left(g(1+t)^3 - 3(1+t)^2\right) dx$ $= \frac{1}{4}g(1+t)^4 - (1+t)^3 + A$ $v = \frac{1}{4}g(1+t) - 1 + A(1+t)^{-3}$ $t = 0, v = 0 \Rightarrow 0 = \frac{1}{4}g - 1 + A$ $v = \frac{1}{4}g(1+t) - 1 + \left(1 - \frac{1}{4}g\right)(1+t)^{-3}$	<p>M1 Rearrange</p> <p>M1 Attempt integrating factor</p> <p>A1 Correct</p> <p>A1 Simplified</p> <p>F1 Multiply DE by their I</p> <p>M1 Integrate</p> <p>A1 RHS</p> <p>F1 Divide by their I (must also divide constant)</p> <p>M1 Use condition</p> <p>E1 Convincingly shown</p>	10
(ii)	$(1+t) \frac{dv}{dt} + 5v = (1+t)g$ $\frac{dv}{dt} + \frac{5}{1+t}v = g$ $I = \exp\left(\int \frac{5}{1+t} dt\right) = e^{5\ln(1+t)} = (1+t)^5$ $(1+t)^5 \frac{dv}{dt} + 5(1+t)^4 v = g(1+t)^5$ $\frac{d}{dt}\left((1+t)^5 v\right) = g(1+t)^5$ $(1+t)^5 v = \int g(1+t)^5 dx$ $= \frac{1}{6}g(1+t)^6 + B$ $v = \frac{1}{6}g(1+t) + B(1+t)^{-5}$ $t = 0, v = 0 \Rightarrow 0 = \frac{1}{6}g + B$ $v = \frac{1}{6}g\left(1+t - (1+t)^{-5}\right)$	<p>M1 Rearrange</p> <p>M1 Attempt integrating factor</p> <p>A1 Simplified</p> <p>F1 Multiply DE by their I</p> <p>M1 Integrate</p> <p>A1 RHS</p> <p>F1 Divide by their I (must also divide constant)</p> <p>M1 Use condition</p> <p>F1 Follow a non-trivial GS</p>	9
(iii)	<p>First model: $\frac{dv}{dt} = \frac{1}{4}g - 3\left(1 - \frac{1}{4}g\right)(1+t)^{-4}$</p> <p>As $t \rightarrow \infty, (1+t)^{-4} \rightarrow 0$</p> <p>Hence acceleration tends to $\frac{1}{4}g$</p> <p>Second model $\frac{dv}{dt} = \frac{1}{6}g\left(1 + 5(1+t)^{-6}\right)$</p> <p>Hence acceleration tends to $\frac{1}{6}g$</p>	<p>M1 Find acceleration</p> <p>B1 Identify term(s) $\rightarrow 0$ in their solution for either model</p> <p>A1</p> <p>M1 Find acceleration</p> <p>A1</p>	5

3(i)	$P = Ae^{0.5t}$ $t = 0, P = 2000 \Rightarrow A = 2000$ $P = 2000e^{0.5t}$	M1 Any valid method M1 Use condition A1	3												
(ii)	CF $P = Ae^{0.5t}$ PI $P = a \cos 2t + b \sin 2t$ $\dot{P} = -2a \sin 2t + 2b \cos 2t$ $-2a \sin 2t + 2b \cos 2t = 0.5(a \cos 2t + b \sin 2t) + 170 \sin 2t$ $-2a = 0.5b + 170$ $2b = 0.5a$ solving $\Rightarrow a = -80, b = -20$ GS $P = Ae^{0.5t} - 80 \cos 2t - 20 \sin 2t$	F1 Correct or follows (i) B1 M1 Differentiate M1 Substitute M1 Compare coefficients M1 Solve A1 F1 Their PI + CF (with one arbitrary constant)	8												
(iii)	$t = 0, P = 2000 \Rightarrow A = 2080$ $P = 2080e^{0.5t} - 80 \cos 2t - 20 \sin 2t$	M1 Use condition F1 Follow a non-trivial GS	2												
(iv)	<table border="1" style="border-collapse: collapse; width: 100%;"> <thead> <tr> <th style="text-align: left;">t</th> <th style="text-align: left;">P</th> <th style="text-align: left;">\dot{P}</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>2000</td> <td>1000</td> </tr> <tr> <td>0.1</td> <td>2100</td> <td>1082.58</td> </tr> <tr> <td>0.2</td> <td>2208</td> <td></td> </tr> </tbody> </table>	t	P	\dot{P}	0	2000	1000	0.1	2100	1082.58	0.2	2208		M1 Use of algorithm A1 2100 A1 1082.5... A1 2208	4
t	P	\dot{P}													
0	2000	1000													
0.1	2100	1082.58													
0.2	2208														
(v)	(A) Limiting value $\Rightarrow \dot{P} = 0$ $\Rightarrow P \left(1 - \frac{P}{12000}\right)^{\frac{1}{2}} = 0$ (as limit non-zero) limiting value = 12000	M1 Set $\dot{P} = 0$ M1 Solve A1	3												
	(B) Growth rate max when $f(P) = P \left(1 - \frac{P}{12000}\right)^{\frac{1}{2}}$ max $f'(P) = \left(1 - \frac{P}{12000}\right)^{\frac{1}{2}} - \frac{1}{2 \times 12000} P \left(1 - \frac{P}{12000}\right)^{-\frac{1}{2}}$ $f'(P) = 0 \Leftrightarrow \left(1 - \frac{P}{12000}\right) - \frac{1}{2 \times 12000} P = 0$ $\Leftrightarrow P = 8000$	M1 Recognise expression to maximise M1 Reasonable attempt at derivative M1 Set derivative to zero A1	4												

4(i) $\ddot{x} = -3\dot{x} + \dot{y}$ $= -3\dot{x} + (-5x + y + 15)$ $y = 3x - 9 + \dot{x}$ $\ddot{x} = -3\dot{x} - 5x + (3x - 9 + \dot{x}) + 15$ $\ddot{x} + 2\dot{x} + 2x = 6$	M1 Differentiate first equation M1 Substitute for \dot{y} M1 y in terms of x, \dot{x} M1 Substitute for y E1	5
(ii) $\lambda^2 + 2\lambda + 2 = 0$ $\lambda = -1 \pm j$ CF $x = e^{-t} (A \cos t + B \sin t)$ PI $x = a$ $2a = 6 \Rightarrow a = 3$ GS $x = 3 + e^{-t} (A \cos t + B \sin t)$	M1 Auxiliary equation A1 M1 CF for complex roots F1 CF for their roots B1 Constant PI B1 PI correct F1 Their CF + PI (with two arbitrary constants)	7
(iii) $y = 3x - 9 + \dot{x}$ $= 9 + 3e^{-t} (A \cos t + B \sin t) - 9$ $- e^{-t} (A \cos t + B \sin t) + e^{-t} (-A \sin t + B \cos t)$ $y = e^{-t} ((2A + B) \cos t + (2B - A) \sin t)$	M1 y in terms of x, \dot{x} M1 Differentiate x and substitute A1 Constants must correspond with those in x	3
(iv) $0 = 3 + A \Rightarrow A = -3$ $0 = 2A + B \Rightarrow B = 6$ $x = 3 + 3e^{-t} (2 \sin t - \cos t)$ $y = 15e^{-t} \sin t$	M1 Condition on x M1 Condition on y F1 Follow their GS F1 Follow their GS	4
(v)  	B1 Sketch of x starts at origin B1 Asymptote $x = 3$ B1 Sketch of y starts at origin B1 Decaying oscillations (may decay rapidly) B1 Asymptote $y = 0$	5

4761

Mechanics 1

Q 1		Mark	Comment	Sub
(i)		B1 B1 B1	Acc and dec shown as straight lines Horizontal straight section All correct with v and times marked and at least one axis labelled. Accept (t, v) or (v, t) used.	3
(ii)	<p>Distance is found from the area</p> <p>area is $\frac{1}{2} \times 10 \times 15 + 20 \times 15 + \frac{1}{2} \times 5 \times 15$</p> <p>(or $\frac{1}{2} \times (20 + 35) \times 15$)</p> <p>= 412.5 so distance is 412.5 m</p>	M1 A1 A1	At least one area attempted or equivalent $uvast$ attempted over one appropriate interval. Award for at least two areas (or equivalent) correct Allow if a trapezium used and only 1 substitution error. FT their diagram. cao (Accept 410 or better accuracy)	3
		6		
2 (i)	$\begin{pmatrix} 6 \\ 9 \end{pmatrix} = 1.5\mathbf{a}$ giving $\mathbf{a} = \begin{pmatrix} 4 \\ 6 \end{pmatrix}$ so $\begin{pmatrix} 4 \\ 6 \end{pmatrix} \text{ m s}^{-2}$	M1 A1	Use of N2L with an attempt to find \mathbf{a} . Condone spurious notation. Must be a vector in proper form. Penalise only once in paper.	2
(ii)	<p>Angle is $\arctan\left(\frac{6}{4}\right)$</p> <p>= 56.309... so 56.3° (3 s. f.)</p>	M1 F1	Use of arctan with their $\frac{6}{4}$ or $\frac{4}{6}$ or equiv. May use F . FT their a provided both cpts are +ve and non-zero.	2
(iii)	<p>Using $\mathbf{s} = t\mathbf{u} + 0.5t^2\mathbf{a}$ we have</p> $\mathbf{s} = 2 \begin{pmatrix} -2 \\ 3 \end{pmatrix} + 0.5 \times 4 \begin{pmatrix} 4 \\ 6 \end{pmatrix}$ <p>so $\begin{pmatrix} 4 \\ 18 \end{pmatrix} \text{ m}$</p>	M1 A1 A1	Appropriate single $uvast$ (or equivalent sequence of $uvast$). If integration used twice condone omission of $\mathbf{r}(0)$ but not $\mathbf{v}(0)$. FT their a only cao. isw for magnitude subsequently found. Vector must be in proper form (penalise only once in paper).	3
		7		

Q 3		Mark	Comment	Sub
(i)	$m \times 9.8 = 58.8$ so $m = 6$	M1 A1	$T = mg$. Condone sign error. cao. CWO.	2
(ii)	Resolve \rightarrow $58.8 \cos 40 - F = 0$ $F = 45.043\dots$ so 45.0 N (3 s. f.)	M1 B1 A1	Resolving their tension. Accept $s \leftrightarrow c$. Condone sign errors but not extra forces. (their T) $\times \cos 40$ (or equivalent) seen Accept ± 45 only.	3
(iii)	Resolve \uparrow $R + 58.8 \sin 40 - 15 \times 9.8 = 0$ $R = 109.204\dots$ so 109 N (3 s. f.)	M1 A1 A1	Resolving their tension. All forces present. No extra forces. Accept $s \leftrightarrow c$. Condone errors in sign. All correct cao	3
		8		
Q 4		Mark	Comment	Sub
(i)	Resultant is $\begin{pmatrix} 4 \\ 1 \\ 2 \end{pmatrix} + \begin{pmatrix} -6 \\ 2 \\ 4 \end{pmatrix} = \begin{pmatrix} -2 \\ 3 \\ 6 \end{pmatrix}$ Magnitude is $\sqrt{(-2)^2 + 3^2 + 6^2} = \sqrt{49} = 7$ N	M1 A1 M1 F1	Adding the vectors. Condone spurious notation. Vector must be in proper form (penalise only once in the paper). Accept clear components. Pythagoras on their 3 component vector. Allow e.g. -2^2 for $(-2)^2$ even if evaluated as -4 . FT their resultant.	4
(ii)	$\mathbf{F} + 2\mathbf{G} + \mathbf{H} = \mathbf{0}$ So $\mathbf{H} = -2\mathbf{G} - \mathbf{F} = -\begin{pmatrix} -12 \\ 4 \\ 8 \end{pmatrix} - \begin{pmatrix} 4 \\ 1 \\ 2 \end{pmatrix}$ $= \begin{pmatrix} 8 \\ -5 \\ -10 \end{pmatrix}$	M1 A1 A1	Either $\mathbf{F} + 2\mathbf{G} + \mathbf{H} = \mathbf{0}$ or $\mathbf{F} + 2\mathbf{G} = \mathbf{H}$ Must see attempt at $\mathbf{H} = -2\mathbf{G} - \mathbf{F}$ cao. Vector must be in proper form (penalise only once in the paper).	3
		7		

Q 5	Mark	Comment	Sub
$a = 12 - 6t$ $a = 0$ gives $t = 2$ $x = \int (2 + 12t - 3t^2) dx$ $2t + 6t^2 - t^3 + C$ $x = 3$ when $t = 0$ so $3 = C$ and $x = 2t + 6t^2 - t^3 + 3$ $x(2) = 4 + 24 - 8 + 3 = 23$ m	M1 A1 F1 M1 A1 M1 A1 B1	Differentiation, at least one term correct. Follow their a Integration indefinite or definite, at least one term correct. Correct. Need not be simplified. Allow as definite integral. Ignore C or limits Allow $x = \pm 3$ or argue it is \int_0^2 from A then ± 3 Award if seen WWW or $x = 2t + 6t^2 - t^3$ seen with +3 added later. FT their t and their x if obtained by integration but not if -3 obtained instead of +3. [If 20 m seen WWW for displacement award SC6] [Award SC1 for position if constant acceleration used for displacement and then +3 applied]	8
	8		


Q 6		Mark	Comment	Sub
(i)	$3.5 = 0.5 + 1.5T$ so $T = 2$ so 2 s	M1 A1	Suitable <i>uvast</i> , condone sign errors. cao	4
	$s = \frac{3.5 + 0.5}{2} \times 2$ so $s = 4$ so 4 m	M1 F1	Suitable <i>uvast</i> , condone sign errors. FT their T . [If s found first then it is <i>cao</i> . In this case when finding T , FT their s , if used.]	
(ii)	(A) N2L \downarrow : $80 \times 9.8 - T = 80 \times 1.5$ $T = 664$ so 664 N	M1 B1 A1	Use of N2L. Allow weight omitted and use of $F = mga$ Condone errors in sign but do not allow extra forces. weight correct (seen in (A) or (B)) cao	5
	(B) N2L \downarrow : $80 \times 9.8 - T = 80 \times (-1.5)$ $T = 904$ so 904 N	M1 A1	N2L with all forces and using $F = ma$. Condone errors in sign but do not allow extra forces. cao [Accept 904 N seen for M1 A1]	
(iii)	N2L \uparrow : $2500 - 80 \times 9.8 - 116 = 80a$ $a = 20$ so 20 m s^{-2} upwards.	M1 A1 A1 A1	Use of N2L with $F = ma$. Allow 1 force missing. No extra forces. Condone errors in sign. ± 20 , accept direction wrong or omitted upwards made clear (accept diagram)	4
(iv)	N2L \uparrow on equipment: $80 - 10 \times 9.8 = 10a$ $a = -1.8$ N2L \uparrow either all: $T - (80 + 10) \times 9.8 - 116 = 90 \times (-1.8)$ or on man: $T - (80 \times 9.8) - 116 - 80 = 80 \times (-1.8)$ $T = 836$ so 836 N	M1 A1 M1 A1	Use of N2L on equipment. All forces. $F = ma$. No extra forces. Allow sign errors. Allow ± 1.8 N2L for system or for man alone. Forces correct (with no extras); accept sign errors; their ± 1.8 used cao [NB The answer 836 N is independent of the value taken for g and hence may be obtained if all weights are omitted.]	4
		17		

Q 7		Mark	Comment	Sub
(i)	Horiz $21t = 60$ so $\frac{20}{7}$ s (2.8571...) either $0 = u - 9.8 \times \frac{20}{7}$ or $-u = u - 9.8 \times (\frac{40}{7})$ or $40 = u \times \frac{20}{7} - 4.9(\frac{20}{7})^2$ so $u = 28$ so 28 m s^{-1}	M1 A1 M1 E1	Use of horizontal components and $a = 0$ or $s = vt - 0.5at^2$ with $v = 0$. Any form acceptable. Allow M1 A1 for answer seen WW. [If $s = ut + 0.5at^2$ and $u = 0$ used without justification award M1 A0] [If $u = 28$ assumed to find time then award SC1] Use of $v = u + at$ (or $v^2 = u^2 + 2as$) with $v = 0$. or Use of $v = u + at$ with $v = -u$ and appropriate t . or Use of $s = ut + 0.5at^2$ with $s = 40$ and appropriate t Condone sign errors and, where appropriate, $u \leftrightarrow v$. Accept signs not clear but not errors. Enough working must be given for 28 to be properly shown. [NB $u = 28$ may be found first and used to find time]	4
(ii)	$y = 28t - 0.5 \times 9.8t^2$	E1	<i>Clear & convincing</i> use of $g = -9.8$ in $s = ut + 0.5at^2$ or $s = vt - 0.5at^2$ NB: AG	1
(iii)	Start from same height with same (zero) vertical speed at same time, same acceleration Distance apart is $0.75 \times 21t = 15.75t$	E1 M1 A1	For two of these reasons $0.75 \times 21t$ seen or $21t$ and $5.25t$ both seen with intention to subtract. Need simplification - LHS alone insufficient. CWO.	3
(iv) (A)	either Time is $\frac{20}{7}$ s by symmetry so $15.75 \times \frac{20}{7} = 45$ so 45 m or Hit ground at same time. By symmetry one travels 60 m so the other travels 15 m in this time ($\frac{1}{4}$ speed) so 45 m.	B1 B1 B1 B1	Symmetry or <i>uvast</i> FT their (iii) with $t = \frac{20}{7}$ [SC1 if 90 m seen]	2
(B)	see next page			

Q7	continued			
(B)	<p>either Time to fall is $40 - 10 = 0.5 \times 9.8 \times t^2$</p> <p>$t = 2.47435\dots$ need $15.75 \times 2.47435\dots = 38.971\dots$ so 39.0 (3sf)</p> <p>or Need time so $10 = 28t - 4.9t^2$</p> <p>$4.9t^2 - 28t + 10 = 0$</p> <p>so $t = \frac{28 \pm \sqrt{28^2 - 4 \times 4.9 \times 10}}{9.8}$ so 0.382784... or 5.33150...</p> <p>Time required is 5.33150... $-\frac{20}{7} =$ 2.47435.. need $15.75 \times 2.47435\dots = 38.971\dots$ so 39.0 (3sf)</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>F1</p> <p>M1</p> <p>M1*</p> <p>A1</p> <p>M1</p> <p>F1</p>	<p>[SC1 if either and or methods mixed to give $\pm 30 = 28t - 4.9t^2$ or $\pm 10 = 4.9t^2$]</p> <p>Considering time from explosion with $u = 0$. Condone sign errors.</p> <p>LHS. Allow ± 30</p> <p>All correct</p> <p>cao</p> <p>FT their (iii) only.</p> <p>Equating $28t - 4.9t^2 = \pm 10$ Dep. Attempt to solve quadratic by a method that could give two roots.</p> <p>Larger root correct to at least 2 s. f. Both method marks may be implied from two correct roots alone (to at least 1 s. f.). [SC1 for either root seen WW]</p> <p>M1</p> <p>FT their (iii) only.</p>	5
(v)	<p>Horiz ($x =$) $21t$ Elim t between $x = 21t$ and $y = 28t - 4.9t^2$</p> <p>so $y = 28\left(\frac{x}{21}\right) - 4.9\left(\frac{x}{21}\right)^2$</p> <p>so $y = \frac{4x}{3} - \frac{0.1x^2}{9} = \frac{1}{90}(120x - x^2)$</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>E1</p>	<p>Intention must be clear, with some attempt made.</p> <p>t completely and correctly eliminated from their expression for x and correct y. Only accept wrong notation if subsequently explicitly given correct value e.g. $\frac{x^2}{21}$ seen as $\frac{x^2}{441}$.</p> <p>Some simplification must be shown.</p> <p>[SC2 for 3 points shown to be on the curve. Award more only if it is made clear that (a) trajectory is a parabola (b) 3 points define a parabola]</p>	4
		19		

4762

Mechanics 2

Q1	Mark	Comment	Sub
(a) (i) either In direction of the force $I = Ft = mv$ so $1500 \times 8 = 4000v$ giving $v = 3$ so 3 m s^{-1} or N2L gives $a = \frac{1500}{4000}$ $v = 0 + \frac{1500}{4000} \times 8$ giving $v = 3$ so 3 m s^{-1}	M1 A1 A1 M1 A1 A1	Use of $Ft = mv$ Appropriate use of N2L and $uvast$	3
(ii) <div style="text-align: center; margin: 10px 0;">  </div> PCLM $12000 = 4000V_R + 500V_S$ so $24 = 8V_R + V_S$ NEL $\frac{V_S - V_R}{0 - 3} = -0.2$ so $V_S - V_R = 0.6$ Solving $V_R = 2.6, V_S = 3.2$ so ram 2.6 m s^{-1} and stone 3.2 m s^{-1}	M1 A1 M1 A1 A1 F1	Appropriate use of PCLM Any form Appropriate use of NEL Any form Either value	6
(iii) $0.5 \times 4000 \times 3^2 - 0.5 \times 4000 \times 2.6^2 - 0.5 \times 500 \times 3.2^2$ $= 1920 \text{ J}$	M1 B1 A1	Change in KE. Accept two terms Any relevant KE term correct (FT their speeds) cao	3
(b) see over			

1		Mark	Comment	Sub
(b)				
(i)	$72\mathbf{i} \text{ N s}$ $8(9 \cos 60\mathbf{i} + 9 \sin 60\mathbf{j})$ $= (36\mathbf{i} + 36\sqrt{3}\mathbf{j}) \text{ N s}$	B1 E1	Neglect units but must include direction Evidence of use of 8 kg , 9 m s^{-1} and 60°	2
(ii)	$72\mathbf{i} + (36\mathbf{i} + 36\sqrt{3}\mathbf{j}) = 12(u\mathbf{i} + v\mathbf{j})$ Equating components $72 + 36 = 12u$ so $u = 9$ $36\sqrt{3} = 12v$ so $v = 3\sqrt{3}$	M1 M1 A1	PCLM. Must be momenta both sides Both	3
(iii)	either $4 \times 18 = 8 \times 9$ so equal momenta so $60/2 = 30^\circ$ or $\arctan\left(\frac{3\sqrt{3}}{9}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = 30^\circ$	M1 A1 M1 A1	Must be clear statements cao FT their u and v . cao	2
		19		

Q 2		Mark	Comment	Sub
(i)				
(A)	$0.5 \times 80 \times 3^2 = 360 \text{ J}$	M1 A1	Use of KE	2
(B)	$360 = F \times 12$ so $F = 30$ so 30 N	M1 F1	$W = Fd$ attempted FT their WD	2
(ii)	Using the WE equation $0.5 \times 80 \times 10^2 - 0.5 \times 80 \times 4^2$ $= 80 \times 9.8 \times h - 1600$ $h = 6.32653\dots$ so 6.33 (3 s. f.)	M1 M1 B1 A1 A1	Attempt to use the WE equation. Condone one missing term Δ KE attempted 1600 with correct sign All terms present and correct (neglect signs) cao	5
(iii)				
(A)	We have driving force $F = 40$ so $200 = 40v$ and $v = 5$ so 5 m s^{-1}	B1 M1 A1	May be implied Use of $P = Fv$	3
(B)	From N2L, force required to give accn is $F - 40 = 80 \times 2$ so $F = 200$ $P = 200 \times 0.5 = 100$ so 100 W	M1 A1 A1 M1 A1	Use of N2L with all terms present (neglect signs) All terms correct correct use of $P = Fv$ cao	5
		17		

Q 3		Mark	Comment	Sub
(i)	For \bar{z} $(2 \times 20 \times 100 + 2 \times 50 \times 120) \bar{z}$ $= 2 \times 2000 \times 50 + 2 \times 6000 \times 60$ so $\bar{z} = 57.5$ and $\bar{y} = 0$	M1 B1 B1 A1 B1	Method for c.m. Total mass of 16000 (or equivalent) At least one term correct NB This result is given below. NB This result is given below. Statement (or proof) required. N.B. If incorrect axes specified, award max 4/5	5
(ii)	\bar{y} and \bar{z} are not changed with the folding For \bar{x} $100 \times 120 \times 0 + 2 \times 20 \times 100 \times 10 = 16000 \bar{x}$ so $\bar{x} = \frac{40000}{16000} = 2.5$	E1 M1 B1 E1	A statement, calculation or diagram required. Method for the c.m. with the folding Use of the 10 Clearly shown	4
(iii)	Moments about AH. Normal reaction acts through this line c.w. $P \times 120 - 72 \times (20 - 2.5) = 0$ so $P = 10.5$	M1 B1 B1 A1 A1	May be implied by diagram or statement 20 - 2.5 or equivalent All correct cao	5
(iv)	$F_{\max} = \mu R$ so $F_{\max} = 72\mu$ For slipping before tipping we require $72\mu < 10.5$ so $\mu < 0.1458333... \left(\frac{7}{48}\right)$	M1 A1 M1 A1	Allow $F = \mu R$ Must have clear indication that this is max F Accept \leq . Accept their F_{\max} and R . cao	4
		18		

Q 4	Mark	Comment	Sub
(i) Centre of CE is 0.5 m from D a.c. moment about D $2200 \times 0.5 = 1100$ so 1100 N m c.w moments about D $R \times 2.75 - 1100 = 0$ $R = 400$ so 400 N	B1 M1 E1 M1 B1 A1	Used below correctly Use of their 0.5 0.5 must be clearly established. Use of moments about D in an equation Use of 1100 and 2.75 or equiv	6
(ii) c.w moments about D $W \times 1.5 - 1100 - 440 \times 2.75 = 0$ so $W = 1540$	M1 A1 E1	Moments of all relevant forces attempted All correct Some working shown	3
(iii) (A) c.w. moments about D $1.5 \times 1540 \cos 20 - 1.75T$ $- 1100 \cos 20 - 400 \times 2.75 \cos 20 = 0$ $T = 59.0663\dots$ so 59.1 N (3 s. f.)	M1 M1 A1 B1 A1 A1	Moments equation. Allow one missing term; there must be some attempt at resolution. At least one res attempt with correct length Allow $\sin \leftrightarrow \cos$ Any two of the terms have $\cos 20$ correctly used (or equiv) 1.75 T All correct cao Accept no direction given	6
(iii) (B) either Angle required is at 70° to the normal to CE so $T_1 \cos 70 = 59.0663\dots$ so $T_1 = 172.698\dots$ so 173 N (3 s.f.) or $400 \cos 20 \times 2.75 + 1100 \cos 20$ $= 1540 \cos 20 \times 1.5 - T \sin 20 \times 1.75$ $T = 172.698\dots$ so 173 N (3s.f.)	B1 M1 A1 M1 A1 A1	FT (iii) (A) Moments attempted with all terms present All correct (neglect signs) FT(iii)(A)	3
	18		

4763

Mechanics 3

1(a)(i)	$[\text{Force}] = \text{MLT}^{-2}$ $[\text{Density}] = \text{ML}^{-3}$	B1 B1 2	
(ii)	$[E] = \frac{[F][l_0]}{[A][l - l_0]} = \frac{(\text{MLT}^{-2})(\text{L})}{(\text{L}^2)(\text{L})}$ $= \text{ML}^{-1} \text{T}^{-2}$	B1 M1 A1 3	for $[A] = \text{L}^2$ Obtaining the dimensions of E
(iii)	$T = \text{L}^\alpha (\text{ML}^{-3})^\beta (\text{ML}^{-1} \text{T}^{-2})^\gamma$ $-2\gamma = 1, \quad \beta + \gamma = 0$ $\gamma = -\frac{1}{2}$ $\beta = \frac{1}{2}$ $\alpha - 3\beta - \gamma = 0$ $\alpha = 1$	B1 cao F1 M1 A1 A1 5	Obtaining equation involving α, β, γ
(b)	$AP = 1.7 \text{ m}$ $F = T \cos \theta$ $R + T \sin \theta = 5 \times 9.8$ $T \cos \theta = 0.4(49 - T \sin \theta)$ $\frac{8}{17} T = 0.4(49 - \frac{15}{17} T)$ $T = 23.8$ $T = k(1.7 - 1.5)$ Stiffness is 119 N m^{-1}	B1 M1 M1 M1 A1 A1 M1 A1 8	Resolving in any direction Resolving in another direction <i>(M1 for resolving requires no force omitted, with attempt to resolve all appropriate forces)</i> Using $F = 0.4R$ to obtain an equation involving just one force (or k) Correct equation <i>Allow</i> $T \cos 61.9$ etc or $R = 28$ or $F = 11.2$ <i>May be implied</i> <i>Allow M1 for $T = \frac{\lambda}{1.5} \times 0.2$</i> If $R = 49$ is assumed, max marks are B1M1M0M0A0A0M1A0

2(a)(i)	$0.1 + 0.01 \times 9.8 = 0.01 \times \frac{u^2}{0.55}$ <p>Speed is 3.3 ms^{-1}</p>	M1 A1 A1 3	Using acceleration $u^2 / 0.55$
(ii)	$\frac{1}{2}m(v^2 - u^2) = mg(2 \times 0.55 - 0.15)$ $\frac{1}{2}(v^2 - 3.3^2) = 9.8 \times 0.95$ $v^2 = 29.51$ $R - mg \cos \theta = m \frac{v^2}{a}$ $R - 0.01 \times 9.8 \times \frac{0.4}{0.55} = 0.01 \times \frac{29.51}{0.55}$ <p>Normal reaction is 0.608 N</p>	M1 A1 M1 A1 A1 5	Using conservation of energy <i>(ft is $v^2 = u^2 + 18.62$)</i> Forces and acceleration towards centre <i>(ft is $\frac{u^2 + 22.54}{55}$)</i>
(b)(i)	$T = 0.8r \omega^2$ $T = \frac{160}{2}(r - 2)$ $\omega^2 = \frac{80(r - 2)}{0.8r} = \frac{100(r - 2)}{r}$ $\omega^2 = 100 - \frac{200}{r} < 100, \text{ so } \omega < 10$	B1 B1 E1 E1 4	
(ii)	$EE = \frac{1}{2} \times \frac{160}{2} \times (r - 2)^2 = 40(r - 2)^2$ $KE = \frac{1}{2}m(r\omega)^2$ $= \frac{1}{2} \times 0.8 \times r^2 \times \frac{100(r - 2)}{r}$ $= 40r(r - 2)$ <p>Since $r > r - 2$, $40r(r - 2) > 40(r - 2)^2$ i.e. $KE > EE$</p>	B1 M1 A1 E1 4	Use of $\frac{1}{2}mv^2$ with $v = r\omega$ From fully correct working only
(iii)	<p>When $\omega = 6$, $36 = \frac{100(r - 2)}{r}$ $r = 3.125$</p> $T = 80(r - 2) = 80(3.125 - 2)$ <p>Tension is 90 N</p>	M1 M1 A1 cao 3	Obtaining r

3 (i)	$\frac{dx}{dt} = A\omega \cos \omega t - B\omega \sin \omega t$ $\frac{d^2x}{dt^2} = -A\omega^2 \sin \omega t - B\omega^2 \cos \omega t$ $= -\omega^2(A \sin \omega t + B \cos \omega t) = -\omega^2 x$	B1 B1 ft E1 3	<i>Must follow from their \dot{x}</i> Fully correct completion SR For $\dot{x} = -A\omega \cos \omega t + B\omega \sin \omega t$ $\ddot{x} = -A\omega^2 \sin \omega t - B\omega^2 \cos \omega t$ award B0B1E0
(ii)	$B\omega = -1.44$ $A\omega = -1.44$ $-B\omega^2 = -0.18$ <i>or</i> $-0.18 = -\omega^2(2)$ $\omega = 0.3, \quad A = -4.8$	B1 M1 A1 cao M1 A1 cao A1 cao 6	Using $\frac{dx}{dt} = -1.44$ when $t = 0$ $\frac{d^2x}{dt^2} = -0.18$ when $t = 0$ (or $x = 2$)
(iii)	Period is $\frac{2\pi}{\omega} = \frac{2\pi}{0.3} = 20.94 = 20.9$ s (3 sf) Amplitude is $\sqrt{A^2 + B^2} = \sqrt{4.8^2 + 2^2}$ $= 5.2$ m	E1 M1 A1 3	or $1.44^2 = 0.3^2(a^2 - 2^2)$
(iv)	$x = -4.8 \sin 0.3t + 2 \cos 0.3t$ $v = -1.44 \cos 0.3t - 0.6 \sin 0.3t$ When $t = 12, \quad x = 0.3306 \quad (v = 1.56)$ When $t = 24, \quad x = -2.5929 \quad (v = -1.35)$ Distance travelled is $(5.2 - 0.3306) + 5.2 + 2.5929$ $= 12.7$ m	M1 A1 M1 M1 A1 5	Finding x when $t = 12$ and $t = 24$ Both displacements correct Considering change of direction Correct method for distance ft from their A, B, ω and amplitude: <i>Third M1 requires the method to be comparable to the correct one</i> <i>A1A1 both require</i> $\omega \approx 0.3, \quad A \neq 0, \quad B \neq 0$ Note ft from $A = +4.8$ is $x_{12} = -3.92 \quad (v < 0) \quad x_{24} = 5.03 \quad (v > 0)$ Distance is $(5.2 - 3.92) + 5.2 + 5.03$ $= 11.5$


4 (i)	$V = \int_1^8 \pi (x^{-\frac{1}{3}})^2 dx$ $= \pi \left[3x^{\frac{1}{3}} \right]_1^8 = 3\pi$ $V\bar{x} = \int_1^8 \pi x (x^{-\frac{1}{3}})^2 dx$ $= \pi \left[\frac{3}{4} x^{\frac{4}{3}} \right]_1^8 = \frac{45}{4}\pi$ $\bar{x} = \frac{\frac{45}{4}\pi}{3\pi}$ $= \frac{15}{4} = 3.75$	M1 A1 M1 A1 M1 A1	π may be omitted throughout Dependent on previous M1M1
(ii)	$A = \int_1^8 x^{-\frac{1}{3}} dx$ $= \left[\frac{3}{2} x^{\frac{2}{3}} \right]_1^8 = \frac{9}{2} = 4.5$ $A\bar{x} = \int_1^8 x (x^{-\frac{1}{3}}) dx$ $= \left[\frac{3}{5} x^{\frac{5}{3}} \right]_1^8 = \frac{93}{5} = 18.6$ $\bar{x} = \frac{18.6}{4.5} = \frac{62}{15} (\approx 4.13)$ $A\bar{y} = \int_1^8 \frac{1}{2} (x^{-\frac{1}{3}})^2 dx$ $= \left[\frac{3}{2} x^{\frac{1}{3}} \right]_1^8 = \frac{3}{2} = 1.5$ $\bar{y} = \frac{1.5}{4.5} = \frac{1}{3}$	M1 A1 M1 A1 A1 M1 A1 A1	If $\frac{1}{2}$ omitted, award M1A0A0

(iii)	$(1)\begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} + (3.5)\begin{pmatrix} 4.5 \\ 0.25 \end{pmatrix} = (4.5)\begin{pmatrix} \frac{62}{15} \\ \frac{1}{3} \end{pmatrix} = \begin{pmatrix} 18.6 \\ 1.5 \end{pmatrix}$ $\bar{x} = 2.85$ $\bar{y} = 0.625$	M1 M1 A1 A1	Attempt formula for CM of composite body (one coordinate sufficient) Formulae for both coordinates; signs must now be correct, but areas (1 and 3.5) may be wrong. ft only if $1 < \bar{x} < 8$ ft only if $0.5 < \bar{y} < 1$ <i>Other methods:</i> M1A1 for \bar{x} M1A1 for \bar{y} <i>(In each case, M1 requires a complete and correct method leading to a numerical value)</i>
-------	---	--------------------------	--

4766

Statistics 1

Q1 (i)	Mode = 7 Median = 12.5	B1 cao B1 cao	2
(ii)	Positive or positively skewed	E1	1
(iii)	(A) Median (B) There is a large outlier or possible outlier of 58 / figure of 58. Just 'outlier' on its own without reference to either 58 or large scores E0 Accept the large outlier affects the mean (more) E1	E1 cao E1indep	2
(iv)	There are $14.75 \times 28 = 413$ messages So total cost = 413×10 pence = £41.30	M1 for 14.75×28 but 413 can also imply the mark A1cao	2
		TOTAL	7
Q2 (i)	$\binom{4}{3} \times 3! = 4 \times 6 = 24$ codes or ${}^4P_3 = 24$ (M2 for 4P_3) Or $4 \times 3 \times 2 = 24$	M1 for 4 M1 for $\times 6$ A1	3
(ii)	$4^3 = 64$ codes	M1 for 4^3 A1 cao	2
		TOTAL	5
Q3 (i)	Probability = $0.3 \times 0.8 = 0.24$	M1 for 0.8 from (1-0.2) A1	2
(ii)	Either: $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ $= 0.3 + 0.2 - 0.3 \times 0.2$ $= 0.5 - 0.06 = 0.44$ Or: $P(A \cup B) = 0.7 \times 0.2 + 0.3 \times 0.8 + 0.3 \times 0.2$ $= 0.14 + 0.24 + 0.06 = 0.44$ Or: $P(A \cup B) = 1 - P(A' \cap B')$ $= 1 - 0.7 \times 0.8 = 1 - 0.56 = 0.44$	M1 for adding 0.3 and 0.2 M1 for subtraction of (0.3 \times 0.2) A1 cao M1 either of first terms M1 for last term A1 M1 for 0.7×0.8 or 0.56 M1 for complete method as seen A1	3
(iii)	$P(A B) = \frac{P(A \cap B)}{P(B)} = \frac{0.06}{0.44} = \frac{6}{44} = 0.136$	M1 for numerator of their 0.06 only M1 for 'their 0.44' in denominator A1 FT (must be valid p)	3
		TOTAL	8

Q4 (i)	$E(X) = 1 \times 0.2 + 2 \times 0.16 + 3 \times 0.128 + 4 \times 0.512 = 2.952$ Division by 4 or other spurious value at end loses A mark $E(X^2) = 1 \times 0.2 + 4 \times 0.16 + 9 \times 0.128 + 16 \times 0.512 = 10.184$ $\text{Var}(X) = 10.184 - 2.952^2 = 1.47$ (to 3 s.f.)	M1 for $\sum rp$ (at least 3 terms correct) A1 cao M1 for $\sum x^2p$ at least 3 terms correct M1 for $E(X^2) - E(X)^2$ Provided ans > 0 A1 FT their $E(X)$ but not a wrong $E(X^2)$	5
(ii)	Expected cost = $2.952 \times \text{£}45000 = \text{£}133000$ (3sf)	B1 FT (no extra multiples / divisors introduced at this stage)	1
(iii)		G1 labelled linear scales G1 height of lines	2
		TOTAL	8
Q5 (i)	Impossible because the competition would have finished as soon as Sophie had won the first 2 matches	E1	1
(ii)	SS, JSS, JSJSS	B1, B1, B1 (-1 each error or omission)	3
(iii)	$0.7^2 + 0.3 \times 0.7^2 + 0.7 \times 0.3 \times 0.7^2 = 0.7399$ or $0.74(0)$ $\{0.49 + 0.147 + 0.1029 = 0.7399\}$	M1 for any correct term M1 for any other correct term M1 for sum of all three correct terms A1 cao	4
		TOTAL	8

Section B																			
Q6 (i)	$\text{Mean} = \frac{180.6}{12} = 15.05 \text{ or } 15.1$ $S_{xx} = 3107.56 - \frac{180.6^2}{12} \text{ or } 3107.56 - 12(\text{their } 15.05)^2 = (389.53)$ $s = \sqrt{\frac{389.53}{11}} = 5.95 \text{ or better}$ NB Accept answers seen without working (from calculator)	B1 for mean M1 for attempt at S_{xx} A1 cao	3																
(ii)	$\bar{x} + 2s = 15.05 + 2 \times 5.95 = 26.95$ $\bar{x} - 2s = 15.05 - 2 \times 5.95 = 3.15$ So no outliers	M1 for attempt at either M1 for both A1 for limits and conclusion FT their mean and sd	3																
(iii)	New mean = $1.8 \times 15.05 + 32 = 59.1$ New s = $1.8 \times 5.95 = 10.7$	B1FT M1 A1FT	3																
(iv)	New York has a higher mean or 'is on average' higher (oe) New York has greater spread /range /variation or SD (oe)	E1FT using $^{\circ}F$ (\bar{x} dep) E1FT using $^{\circ}F$ (σ dep)	2																
(v)	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td style="padding: 2px;">Upper bound</td> <td style="padding: 2px;">(70)</td> <td style="padding: 2px;">100</td> <td style="padding: 2px;">110</td> <td style="padding: 2px;">120</td> <td style="padding: 2px;">150</td> <td style="padding: 2px;">170</td> <td style="padding: 2px;">190</td> </tr> <tr> <td style="padding: 2px;">Cumulative frequency</td> <td style="padding: 2px;">(0)</td> <td style="padding: 2px;">6</td> <td style="padding: 2px;">14</td> <td style="padding: 2px;">24</td> <td style="padding: 2px;">35</td> <td style="padding: 2px;">45</td> <td style="padding: 2px;">48</td> </tr> </table> 	Upper bound	(70)	100	110	120	150	170	190	Cumulative frequency	(0)	6	14	24	35	45	48	B1 for all correct cumulative frequencies (may be implied from graph). Ignore cf of 0 at this stage G1 for linear scales (linear from 70 to 190) ignore $x < 70$ vertical: 0 to 50 but not beyond 100 (no inequality scales) G1 for labels G1 for points plotted as (UCB, their cf). <u>Ignore (70,0) at this stage.</u> No mid-point or LCB plots.	5
Upper bound	(70)	100	110	120	150	170	190												
Cumulative frequency	(0)	6	14	24	35	45	48												
(vi)	NB all G marks dep on attempt at cumulative frequencies. NB All G marks dep on attempt at cumulative frequencies Line on graph at cf = 43.2(soi) or used 90th percentile = 166	G1 for joining all of 'their points'(line or smooth curve) AND now including (70,0) M1 for use of 43.2 A1FT but dep on 3rd G mark earned	2																
		TOTAL	18																


Question 1

(i)	x is independent, y is dependent since the values of x are chosen by the student but the values of y are dependent on x	B1 E1 dep E1 dep	3
(ii)	$\bar{x} = 2.5, \bar{y} = 80.63$ $b = \frac{S_{xy}}{S_{xx}} = \frac{2530.3 - 30 \times 967.6/12}{90 - 30^2/12} = \frac{111.3}{15} = 7.42$ OR $b = \frac{2530.3/12 - 2.50 \times 80.63}{90/12 - 2.50^2} = \frac{9.275}{1.25} = 7.42$ Hence least squares regression line is: $y - \bar{y} = b(x - \bar{x})$ $\Rightarrow y - 80.63 = 7.42(x - 2.5)$ $\Rightarrow y = 7.42x + 62.08$	B1 for \bar{x} and \bar{y} used (SOI) M1 for attempt at gradient (b) A1 for 7.42 cao M1 for equation of line A1 FT ($b > 0$) for complete equation	5
(iii)	(A) For $x = 1.2$, predicted growth $= 7.42 \times 1.2 + 62.08 = 71.0$ (B) For $x = 4.3$, predicted growth $= 7.42 \times 4.3 + 62.08 = 94.0$ Valid relevant comments relating to the predictions such as : Comment re interpolation/extrapolation Comment relating to the fact that $x = 4.3$ is only just beyond the existing data. Comment relating to size of residuals near each predicted value (need not use word 'residual')	M1 for at least one prediction attempted. A1 for both answers (FT their equation if $b > 0$) E1 (first comment) E1 (second comment)	4
(iv)	$x = 3 \Rightarrow$ predicted $y = 7.42 \times 3 + 62.08 = 84.3$ Residual = $80 - 84.3 = -4.3$	M1 for prediction M1 for subtraction A1 FT ($b > 0$)	3
(v)	This point is a long way from the regression line. The line may be valid for the range used in the experiment but then the relationship may break down for higher concentrations, or the relationship may be non linear.	E1 E1 for valid in range E1 for <i>either</i> 'may break down' <i>or</i> 'could be non linear' <i>or</i> other relevant comment	3
			18

Question 2

(i)	Binomial (94,0.1)	B1 for binomial B1 dep for parameters	2
(ii)	n is large and p is small	B1, B1 Allow appropriate numerical ranges	2
(iii)	$\lambda = 94 \times 0.1 = 9.4$ (A) $P(X = 4) = e^{-9.4} \frac{9.4^4}{4!} = 0.0269$ (3 s.f.) or from tables = 0.0429 – 0.0160 = 0.0269 <i>cao</i> (B) Using tables: $P(X \geq 4) = 1 - P(X \leq 3)$ = 1 – 0.0160 = 0.9840 <i>cao</i>	B1 for mean M1 for calculation or use of tables A1 M1 for attempt to find $P(X \geq 4)$ A1 <i>cao</i>	5
(iv)	P(sufficient rooms throughout August) = $0.9840^{31} = 0.6065$	M1 A1 FT	2
(v)	(A) $31 \times 94 = 2914$ Binomial (2914,0.1) (B) Use Normal approx with $\mu = np = 2914 \times 0.1 = 291.4$ $\sigma^2 = npq = 2914 \times 0.1 \times 0.9 = 262.26$ $P(X \leq 300.5) = P\left(Z \leq \frac{300.5 - 291.4}{\sqrt{262.26}}\right)$ = $P(Z \leq 0.5619) = \Phi(0.5619) = 0.7130$	B1 for binomial B1 dep, for parameters B1 B1 B1 for continuity corr. M1 for probability using correct tail A1 <i>cao</i> , (but FT wrong or omitted CC)	5
			18

Question 3

(i)	$X \sim N(56, 6.5^2)$ $P(52.5 < X < 57.5) = P\left(\frac{52.5 - 56}{6.5} < Z < \frac{57.5 - 56}{6.5}\right)$ $= P(-0.538 < Z < 0.231)$ $= \Phi(0.231) - (1 - \Phi(0.538))$ $= 0.5914 - (1 - 0.7046)$ $= 0.5914 - 0.2954$ $= 0.2960 \text{ (4 s.f.) or } 0.296 \text{ (to 3 s.f.)}$	<p>M1 for standardizing</p> <p>A1 for -0.538 and 0.231</p> <p>M1 for prob. with tables and correct structure</p> <p>A1 CAO (min 3 s.f., to include use of difference column)</p>	4
(ii)	$P(\text{5-year-old} < 62) = P\left(Z < \frac{62 - 56}{6.5}\right)$ $= \Phi(0.923) = 0.8220$ $P(\text{young adult} < 62) = P\left(Z < \frac{62 - 68}{10}\right)$ $= \Phi(-0.6) = 1 - 0.7257 = 0.2743$ $P(\text{One over, one under})$ $= 0.8220 \times 0.7257 + 0.1780 \times 0.2743$ $= 0.645$	<p>B1 for 0.8220 or 0.1780</p> <p>B1 for 0.2743 or 0.7257</p> <p>M1 for either product</p> <p>M1 for sum of both products</p> <p>A1 CAO</p>	5
(iii)		<p>G1 for shape</p> <p>G1 for means, shown explicitly or by scale</p> <p>G1 for lower max height in young adults</p> <p>G1 for greater variance in young adults</p>	4
(iv)	$Y \sim N(82, \sigma^2)$ <p>From tables $\Phi^{-1}(0.88) = 1.175$</p> $\frac{62 - 82}{\sigma} = -1.175$ $-20 = -1.175 \sigma$ $\sigma = 17.0$	<p>B1 for 1.175 seen</p> <p>M1 for equation in σ with z-value</p> <p>M1 for correct handling of LH tail</p> <p>A1 cao</p>	4
			17

Question 4

<p>(i)</p> <p>H_0: no association between sex and subject; H_1: some association between sex and subject;</p> <table border="1" data-bbox="247 347 973 560"> <thead> <tr> <th>OBS</th> <th>Maths</th> <th>English</th> <th>Both</th> <th>Neither</th> <th>Row sum</th> </tr> </thead> <tbody> <tr> <td>Male</td> <td>38</td> <td>19</td> <td>6</td> <td>32</td> <td>95</td> </tr> <tr> <td>Female</td> <td>42</td> <td>55</td> <td>9</td> <td>49</td> <td>155</td> </tr> <tr> <td>Col sum</td> <td>80</td> <td>74</td> <td>15</td> <td>81</td> <td>250</td> </tr> </tbody> </table> <table border="1" data-bbox="247 627 973 840"> <thead> <tr> <th>EXP</th> <th>Maths</th> <th>English</th> <th>Both</th> <th>Neither</th> <th>Row sum</th> </tr> </thead> <tbody> <tr> <td>Male</td> <td>30.40</td> <td>28.12</td> <td>5.70</td> <td>30.78</td> <td>95</td> </tr> <tr> <td>Female</td> <td>49.60</td> <td>45.88</td> <td>9.30</td> <td>50.22</td> <td>155</td> </tr> <tr> <td>Col sum</td> <td>80</td> <td>74</td> <td>15</td> <td>81</td> <td>250</td> </tr> </tbody> </table> <table border="1" data-bbox="247 907 973 1019"> <thead> <tr> <th>CONT</th> <th>Maths</th> <th>English</th> <th>Both</th> <th>Neither</th> </tr> </thead> <tbody> <tr> <td>Male</td> <td>1.900</td> <td>2.958</td> <td>0.016</td> <td>0.048</td> </tr> <tr> <td>Female</td> <td>1.165</td> <td>1.813</td> <td>0.010</td> <td>0.030</td> </tr> </tbody> </table> <p>$\chi^2 = 7.94$</p> <p>Refer to χ^2_3 Critical value at 5% level = 7.815 Result is significant There is evidence to suggest that there is some association between sex and subject choice. NB if H_0 H_1 reversed, or 'correlation' mentioned, do not award first B1 or final E1</p>	OBS	Maths	English	Both	Neither	Row sum	Male	38	19	6	32	95	Female	42	55	9	49	155	Col sum	80	74	15	81	250	EXP	Maths	English	Both	Neither	Row sum	Male	30.40	28.12	5.70	30.78	95	Female	49.60	45.88	9.30	50.22	155	Col sum	80	74	15	81	250	CONT	Maths	English	Both	Neither	Male	1.900	2.958	0.016	0.048	Female	1.165	1.813	0.010	0.030	<p>B1</p> <p>M1 A2 for expected values (allow A1 for at least one row or column correct)</p> <p>M1 for valid attempt at $(O-E)^2/E$ A1 NB These M1 A1 marks cannot be implied by a correct final value of χ^2</p> <p>M1 for summation A1 cao for χ^2</p> <p>B1 for 3 deg of f B1 CAO for cv</p> <p>B1</p> <p>E1</p>	<p>1</p> <p>7</p> <p>4</p>
OBS	Maths	English	Both	Neither	Row sum																																																												
Male	38	19	6	32	95																																																												
Female	42	55	9	49	155																																																												
Col sum	80	74	15	81	250																																																												
EXP	Maths	English	Both	Neither	Row sum																																																												
Male	30.40	28.12	5.70	30.78	95																																																												
Female	49.60	45.88	9.30	50.22	155																																																												
Col sum	80	74	15	81	250																																																												
CONT	Maths	English	Both	Neither																																																													
Male	1.900	2.958	0.016	0.048																																																													
Female	1.165	1.813	0.010	0.030																																																													
<p>(ii)</p> <p>$H_0: \mu = 67.4$; $H_1: \mu > 67.4$ Where μ denotes the mean score of the population of students taught with the new method.</p> <p>Test statistic = $\frac{68.3 - 67.4}{8.9/\sqrt{12}} = \frac{0.9}{2.57} = 0.35$</p> <p>10% level 1 tailed critical value of z = 1.282 0.35 < 1.282 so not significant. There is insufficient evidence to reject H_0 There is insufficient evidence to conclude that the mean score is increased by the new teaching method.</p>	<p>B1 for both correct</p> <p>B1 for definition of μ</p> <p>M1</p> <p>A1 cao</p> <p>B1 for 1.282</p> <p>M1 for comparison</p> <p>A1 for conclusion in words and in context</p>	<p>7</p> <p>19</p>																																																															

4768

Statistics 3


Q1	$P(T > t) = \frac{k}{t^2}, \quad t \geq 1,$																																				
(i)	$F(t) = P(T < t) = 1 - P(T > t)$ $\therefore F(t) = 1 - \frac{k}{t^2}$ $F(1) = 0$ $\therefore 1 - \frac{k}{1^2} = 0$ $\therefore k = 1$	M1 M1 A1	Use of $1 - P(\dots)$. Beware: answer given.	3																																	
(ii)	$f(t) = \frac{d F(t)}{d t}$ $= \frac{2}{t^3}$	M1 A1	Attempt to differentiate c's cdf. (For $t \geq 1$, but condone absence of this.) Ft c's cdf provided answer sensible.	2																																	
(iii)	$\mu = \int_1^{\infty} t f(t) dt = \int_1^{\infty} \frac{2}{t^2} dt$ $= \left[\frac{-2}{t} \right]_1^{\infty}$ $= 0 - (-2) = 2$	M1 A1 A1	Correct form of integral for the mean, with correct limits. Ft c's pdf. Correctly integrated. Ft c's pdf. Correct use of limits leading to correct value. Ft c's pdf provided answer sensible.	3																																	
(b)	$H_0: m = 5.4$ $H_1: m \neq 5.4$ where m is the population median time for the task. <table border="1" data-bbox="239 1249 670 1668"> <thead> <tr> <th>Times</th> <th>- 5.4</th> <th>Rank of diff </th> </tr> </thead> <tbody> <tr><td>6.4</td><td>1.0</td><td>8</td></tr> <tr><td>5.9</td><td>0.5</td><td>5</td></tr> <tr><td>5.0</td><td>-0.4</td><td>4</td></tr> <tr><td>6.2</td><td>0.8</td><td>7</td></tr> <tr><td>6.8</td><td>1.4</td><td>10</td></tr> <tr><td>6.0</td><td>0.6</td><td>6</td></tr> <tr><td>5.2</td><td>-0.2</td><td>2</td></tr> <tr><td>6.5</td><td>1.1</td><td>9</td></tr> <tr><td>5.7</td><td>0.3</td><td>3</td></tr> <tr><td>5.3</td><td>-0.1</td><td>1</td></tr> </tbody> </table> $W_- = 1 + 2 + 4 = 7$ (or $W_+ = 3 + 5 + 6 + 7 + 8 + 9 + 10 = 48$) Refer to tables of Wilcoxon single sample (/paired) statistic for $n = 10$. Lower (or upper if 48 used) double-tailed 5% point is 8 (or 47 if 48 used). Result is significant. Seems that the median time is no longer as previously thought.	Times	- 5.4	Rank of diff	6.4	1.0	8	5.9	0.5	5	5.0	-0.4	4	6.2	0.8	7	6.8	1.4	10	6.0	0.6	6	5.2	-0.2	2	6.5	1.1	9	5.7	0.3	3	5.3	-0.1	1	B1 B1 M1 M1 A1 B1 M1 A1 A1	Both hypotheses. Hypotheses in words only must include "population". For adequate verbal definition. for subtracting 5.4. for ranks. FT if ranks wrong. No ft from here if wrong. i.e. a 2-tail test. No ft from here if wrong. ft only c's test statistic. ft only c's test statistic.	10
Times	- 5.4	Rank of diff																																			
6.4	1.0	8																																			
5.9	0.5	5																																			
5.0	-0.4	4																																			
6.2	0.8	7																																			
6.8	1.4	10																																			
6.0	0.6	6																																			
5.2	-0.2	2																																			
6.5	1.1	9																																			
5.7	0.3	3																																			
5.3	-0.1	1																																			

Q2	$X \sim N(260, \sigma = 24)$		When a candidate's answers suggest that (s)he appears to have neglected to use the difference columns of the Normal distribution tables penalise the first occurrence only.	
(i)	$P(X < 300) = P\left(Z < \frac{300 - 260}{24} = 1.6667\right)$ $= 0.9522$	M1 A1 A1	For standardising. Award once, here or elsewhere.	3
(ii)	$Y \sim N(260 \times 0.6 = 156, 24^2 \times 0.6^2 = 207.36)$ $P(Y > 175) = P\left(Z > \frac{175 - 156}{14.4} = 1.3194\right)$ $= 1 - 0.9063 = 0.0937$	B1 B1 A1	Mean. Variance. Accept sd (= 14.4). c.a.o.	3
(iii)	$Y_1 + Y_2 + Y_3 + Y_4 \sim N(624, 829.44)$ $P(\text{this} < 600) = P\left(Z < \frac{600 - 624}{28.8} = -0.8333\right)$ $= 1 - 0.7976 = 0.2024$	B1 B1 A1	Mean. Ft mean of (ii). Variance. Accept sd (= 28.8). Ft variance of (ii). c.a.o.	3
(iv)	Require w such that $0.975 = P(\text{above} > w) = P\left(Z > \frac{w - 624}{28.8}\right)$ $= P(Z > -1.96)$ $\therefore w - 624 = 28.8 \times -1.96 \Rightarrow w = 567.5(52)$	M1 B1 A1	Formulation of requirement. - 1.96 Ft parameters of (iii).	3
(v)	$On \sim N(150, \sigma = 18)$ $X_1 + X_2 + X_3 + On_1 + On_2 \sim N(1080, 2376)$ $P(\text{this} > 1000) = P\left(Z > \frac{1000 - 1080}{48.744} = -1.6412\right)$ $= 0.9496$	B1 B1 A1	Mean. Variance. Accept sd (= 48.744). c.a.o.	3
(vi)	Given $\bar{x} = 252.4$ $s_{n-1} = 24.6$ CI is given by $252.4 \pm 2.576 \times \frac{24.6}{\sqrt{100}}$ $= 252.4 \pm 6.33(6) = (246.0(63), 258.7(36))$	M1 B1 A1	Correct use of 252.4 and $24.6/\sqrt{100}$. For 2.576. c.a.o. Must be expressed as an interval.	3
				18


Q3				
(i)	<p>A t test should be used because the sample is small, the population variance is unknown, the background population is Normal</p>	E1 E1 E1	3	
(ii)	<p>$H_0: \mu = 380$ $H_1: \mu < 380$</p> <p>where μ is the mean temperature in the chamber.</p> <p>$\bar{x} = 373.825$ $s_{n-1} = 9.368$</p> <p>Test statistic is $\frac{373.825 - 380}{\frac{9.368}{\sqrt{12}}}$</p> <p style="text-align: center;">$= -2.283(359).$</p> <p>Refer to t_{11}. Single-tailed 5% point is -1.796.</p> <p>Significant. Seems mean temperature in the chamber has fallen.</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p>	<p>Both hypotheses. Hypotheses in words only must include "population".</p> <p>For adequate verbal definition. Allow absence of "population" if correct notation μ is used, but do NOT allow "$\bar{X} = \dots$" or similar unless \bar{X} is clearly and explicitly stated to be a <u>population</u> mean.</p> <p>$s_n = 8.969$ but do <u>NOT</u> allow this here or in construction of test statistic, but FT from there.</p> <p>Allow c's \bar{x} and/or s_{n-1}. Allow alternative: $380 + (c's - 1.796) \times \frac{9.368}{\sqrt{12}}$ ($= 375.143$) for subsequent comparison with \bar{x}. (Or $\bar{x} - (c's - 1.796) \times \frac{9.368}{\sqrt{12}}$ ($= 378.681$) for comparison with 380.)</p> <p>c.a.o. but ft from here in any case if wrong. Use of $380 - \bar{x}$ scores M1A0, but ft.</p> <p>No ft from here if wrong. Must be minus 1.796 unless absolute values are being compared. No ft from here if wrong.</p> <p>ft only c's test statistic.</p> <p>ft only c's test statistic.</p>	9
(iii)	<p>CI is given by</p> $373.825 \pm 2.201 \times \frac{9.368}{\sqrt{12}}$ <p>$= 373.825 \pm 5.952 = (367.87(3), 379.77(7))$</p>	<p>M1</p> <p>B1</p> <p>M1</p> <p>A1</p>	<p>c.a.o. Must be expressed as an interval. ZERO/4 if not same distribution as test. Same wrong distribution scores maximum M1B0M1A0. Recovery to t_{11} is OK.</p>	4
(iv)	<p>Advantage: greater certainty. Disadvantage: less precision.</p>	E1 E1	2 18	

Q4																									
(a) (i)	$\bar{x} = \frac{1125}{500} = 2.25$ For binomial $E(X) = n \times p$ $\therefore \hat{p} = \frac{2.25}{5} = 0.45$	B1 M1 A1	Use of mean of binomial distribution. May be implicit. Beware: answer given.	3																					
(ii)	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td>f_o</td> <td>32</td> <td>110</td> <td>154</td> <td>125</td> <td>63</td> <td>16</td> </tr> <tr> <td>f_e (calc)</td> <td>25.164</td> <td>102.944</td> <td>168.455</td> <td>137.827</td> <td>56.384</td> <td>9.226</td> </tr> <tr> <td>f_e (tables)</td> <td>25.15</td> <td>102.95</td> <td>168.45</td> <td>137.85</td> <td>56.35</td> <td>9.25</td> </tr> </table> $\chi^2 = 1.8571 + 0.4836 + 1.2404 + 1.1938 + 0.7763 + 4.9737$ $= 10.52(49)$ Refer to χ_4^2 . Upper 5% point is 9.488. Significant. Suggests binomial model does not fit. The model appears to overestimate in the middle and to underestimate at the tails. The biggest discrepancy is at $X = 5$. A binomial model assumes all trials are independent with a constant probability of "success". It seems unlikely that there will be independence within families and/or that p will be the same for all families.	f_o	32	110	154	125	63	16	f_e (calc)	25.164	102.944	168.455	137.827	56.384	9.226	f_e (tables)	25.15	102.95	168.45	137.85	56.35	9.25	M1 A1 M1 A1 M1 A1 A1 A1 E1 E1 E2	Calculation of expected frequencies. All correct. Or using tables: $1.8657 + 0.4828 + 1.2396 + 1.1978 + 0.7848 + 4.9257$ c.a.o. Or using tables: 10.49(64) Allow correct df (= cells – 2) from wrongly grouped or ungrouped table, and FT. Otherwise, no FT if wrong. No ft from here if wrong. ft only c's test statistic. ft only c's test statistic. Accept also any other sensible comment e.g. at 2.5% significance, the result would NOT have been significant. (E2, 1, 0) Any sensible comment which addresses independence and constant p .	12
f_o	32	110	154	125	63	16																			
f_e (calc)	25.164	102.944	168.455	137.827	56.384	9.226																			
f_e (tables)	25.15	102.95	168.45	137.85	56.35	9.25																			
(b)	She should try to choose a simple random sample which would involve establishing a sampling frame and using some form of random number generator.	E1 E1 E1	Allow sensible discussion of practical limitations of choosing a random sample. Allow other sensible suggestions. E.g systematic sample - choosing every tenth family; stratified sample - by the number of girls in a family.	3																					
				18																					

1

(i) 6 routes $M \rightarrow A \rightarrow I \rightarrow T \rightarrow Pi \rightarrow C$ $M \rightarrow A \rightarrow I \rightarrow T \rightarrow Pi \rightarrow R \rightarrow C$ $M \rightarrow A \rightarrow I \rightarrow T \rightarrow Pi \rightarrow H \rightarrow R \rightarrow C$ $M \rightarrow V \rightarrow I \rightarrow T \rightarrow Pi \rightarrow C$ $M \rightarrow V \rightarrow I \rightarrow T \rightarrow Pi \rightarrow R \rightarrow C$ $M \rightarrow V \rightarrow I \rightarrow T \rightarrow Pi \rightarrow H \rightarrow R \rightarrow C$	B1 B1
(ii) 6 routes $M \rightarrow A \rightarrow I \rightarrow Pa \rightarrow Pi \rightarrow C$ $M \rightarrow A \rightarrow I \rightarrow Pa \rightarrow Pi \rightarrow R \rightarrow C$ $M \rightarrow A \rightarrow I \rightarrow Pa \rightarrow Pi \rightarrow H \rightarrow R \rightarrow C$ $M \rightarrow V \rightarrow I \rightarrow Pa \rightarrow Pi \rightarrow C$ $M \rightarrow V \rightarrow I \rightarrow Pa \rightarrow Pi \rightarrow R \rightarrow C$ $M \rightarrow V \rightarrow I \rightarrow Pa \rightarrow Pi \rightarrow H \rightarrow R \rightarrow C$	B1 B1
(iii) $M \rightarrow V \rightarrow I \rightarrow Pa \rightarrow Pi \rightarrow H \rightarrow R \rightarrow Pi \rightarrow C$ 	B1
(iv) e.g. $P \rightarrow T \rightarrow I \rightarrow V \rightarrow M \rightarrow A \rightarrow I \rightarrow Pa \rightarrow P \rightarrow H \rightarrow R \rightarrow C \rightarrow P \rightarrow R$	M1 ends at R A2 (-1 each error/omission)

2. y



(i)	lines	B1	x
	shading	B1	graph or sim. eqns
	(3.6, 0.6)	M1 A1	
	25.8 at (3.6, 0.6) versus 21 and 24 (or profit line)		
(ii)	25 at (3, 1)	B1	B1

3.

$y = 2008$	
$c = 2008/100 = 20$	
$n = 2008 - 19 \times (2008/19) = 2008 - 19 \times (105) = 13$	
$k = 3/25 = 0$	B1
$i = 20 - 5 - 20 / 3 + 19 \times 13 + 15 = 271$	B1
$i = 1$	
$i = 1 - 0 = 1$	B1
$j = 2008 + 502 + 1 + 2 - 20 + 5 = 2498$	
$j = 6$	B1
$p = -5$	B1
$m = 3$	B1
$d = 23$	B1
So 23 rd March	B1

4.

(i) e.g.	0–3→brown 4–7→blue 8–9→green	M1 A1 proportions OK A1 efficient																																	
(ii) e.g.	0–1→brown 2–5→blue 6–7→green 8–9→reject	M1 some rejected A2 proportions OK (–1 each error) A1 efficient																																	
(iii) e.g.	Eye colours <table border="1" data-bbox="245 658 807 848"> <tbody> <tr> <td>Parent 1</td> <td>brown</td> <td>brown</td> <td>brown</td> <td>blue</td> </tr> <tr> <td>Parent 2</td> <td>brown</td> <td>blue</td> <td>brown</td> <td>blue</td> </tr> <tr> <td>Offspring</td> <td>brown</td> <td>brown</td> <td>brown</td> <td>brown</td> </tr> </tbody> </table> <table border="1" data-bbox="245 882 850 1072"> <tbody> <tr> <td>brown</td> <td>green</td> <td>blue</td> <td>green</td> <td>brown</td> <td>brown</td> </tr> <tr> <td>brown</td> <td>blue</td> <td>brown</td> <td>green</td> <td>brown</td> <td>green</td> </tr> <tr> <td>brown</td> <td>blue</td> <td>brown</td> <td>green</td> <td>brown</td> <td>blue</td> </tr> </tbody> </table>	Parent 1	brown	brown	brown	blue	Parent 2	brown	blue	brown	blue	Offspring	brown	brown	brown	brown	brown	green	blue	green	brown	brown	brown	blue	brown	green	brown	green	brown	blue	brown	green	brown	blue	B1 br/br→br (4 times) B1 br/gr→bl B1 gr/gr→gr M1 br/bl rule A1 application A1 application B1 bl/bl application M1 gr/bl rule A1 application
Parent 1	brown	brown	brown	blue																															
Parent 2	brown	blue	brown	blue																															
Offspring	brown	brown	brown	brown																															
brown	green	blue	green	brown	brown																														
brown	blue	brown	green	brown	green																														
brown	blue	brown	green	brown	blue																														

5.

<p>(i)&(ii) e.g.</p> <p>time – 55 weeks critical – A; B; F; G; J</p> <p>(iii) 50 weeks (49 weeks if G crashed rather than H)</p> <p>(iv) E – 1 week F – 3 weeks J – 2 weeks (G – 1 week, if crashed)</p> <p>(v) £115000 (£121000)</p>	<p>M1 sca (activity on arc) A1 dummy activities + E and F A1 A, B, C, D A1 G, H, I, J</p> <p>M1 forward pass A1</p> <p>M1 backward pass A1</p> <p>B1 cao B1 cao</p> <p>B1</p> <p>M1 A3</p> <p>A1</p>
---	--

6.

(i) e.g.

0.3 0.1 0.1 0.2 0.2
Total length = 2.2 km

0.2
0.2
0.1
0.1

M1 connecting tree
A1 DE
A1 FC, FG
A1 AD, DI, FH
A1 2 of length 0.4

M1 A1

(ii) Prim: connect in nearest to connected set
Kruskal: Shortest arc s.t. no cycles

(iii)

5 0.7
0.7 B
8 1.1
1.1 C
6 0.9
0.9 F
9 1.1
1.5 1.1 G
1 0 A
2 0.3
0.3 D
3 0.4
0.4 E
7 1.0
1.0 H
4 0.6
0.6 I

0.3 0.1 0.1 0.2 0.2
Arcs used: AD, DE, EF, FG, DI, IH, AB or DB, FC or BC
Total length = 2.7 km (AB&FC) or 2.9 km (AB&BC) or 2.4 km (DB&FC) or 2.6 km (DB&BC)

0.2
0.2
0.1
0.1

M1 Dijkstra
A1 working values (see vertex G)
A1 order of labelling
A1 labels

M1 arcs counted
A1 only once
A1

4776 Numerical Methods

1	x	2	3	root = $(2 \times 0.03 - 3 \times 0.24) / (0.03 - 0.24)$	[M1A1]
	f(x)	0.24	0.03	= 3.142857	[A1]
	Eg: graph showing turning point at x = 3 with root some way to the left or the right.				[G2]
[TOTAL 5]					

2	x	f(x)			
	0	1			
	1	0.333333	T1 =	0.666667	[M1A1]
	0.5	0.477592	M =	0.477592	[M1A1]
			hence	$T2 = (T1 + M)/2 =$	0.572129 [M1A1]
			and	$S = (T1 + 2*M)/3 =$	0.540617 [M1A1]
[TOTAL 8]					

3	x	0	1	3	
	f(x)	2	2.57	3.85	
					3 terms: [M1]
					form: [M1]
					use x=2: [M1]
	f(2) =	$2(2-1)(2-3)/(0-1)(0-3) + 2.57(2-0)(2-3)/(1-0)(1-3) + 3.85(2-0)(2-1)/(3-0)(3-1)$			[A1A1A1]
	=	3.186667	(3.19)		[A1]
[TOTAL 7]					

4	x	1.5	2		
	$x^3(2-x)-1$	0.6875	-1	change of sign, so root (<i>may be implied</i>)	[M1A1]
	a	b	x	$x^3(2-x)-1$	mpe
	1.5	2	1.75	0.339844	0.25 [M1A1]
	1.75	2	1.875	-0.17603	0.125 [A1]
	1.75	1.875	1.8125		0.0625 [A1]
	4 further iterations reqd: mpe 0.0325, 0.015625, 0.0078125, 0.00390625				[M1A1]
[TOTAL 8]					

5	Sketch showing curve, tangent, chord, h. Makes clear that tangent and chord have substantially different gradients.				[G3]
	h	0	0.1	0.01	0.001
	g(2 + h)	3.61	3.849	3.633	3.612
	est g'(2)		2.39	2.3	2
	Clear loss of significant figures as h is reduced				[M1A1A1A1]
					[E1]
[TOTAL 8]					

4776

Mark Scheme

January 200

6 (i)	x	f(x)	Δf	$\Delta^2 f$	$\Delta^3 f$	
	3	1				
	4	3	2			
	5	-1	-4	-6		
	6	-10	-9	-5	1	

[M1A1A1]

quadratic = $1 + 2(x-3) - 6(x-3)(x-4)/2$ [M1A1]
 = $1 + 2x - 6 - 3x^2 + 21x - 36$ [A1]
 = $-3x^2 + 23x - 41$ [A1]

$q'(x) = -6x + 23 = 0$ at $x = 23/6 (= 3.833\dots)$ [M1A1]

$q(x) = 0$ at $x = 4.847(127)$; also at 2.81954 - not reqd. [M1A1]

$q(6) = -11$ (or point out that the second differences not constant) [A1]
 [subtotal 12]

(ii) cubic est = $1 + 2(4.5-3) - 6(4.5-3)(4.5-4)/2 + 1(4.5-3)(4.5-4)(4.5-5)/6$ [M1A1A1]
 = 1.6875 [A1]

$S = 1.5/3 (1 + 4 \times 1.6875 - 10) = -1.125$ [M1A1]
 [subtotal 6]

[TOTAL 18]

7 (i) mpe 0.000 000 5 [B1]
 mppe 0.000 000 5 / 2.506 628 [M1A1]
 = 1.99×10^{-7} [subtotal 3]

(ii) mpe $1000 \times 0.000 000 5 = 0.000 5$ [M1A1]
 In practice the positive and negative errors will tend to cancel out [E1]
 [subtotal 3]

(iii) mpe $1000 \times 0.000 001 = 0.001$ [M1A1]
 In practice $1000 \times 0.000 000 5 = 0.000 5$ [M1A1]
 because average error in chopping will be 0.000 000 5 [E1]
 [subtotal 5]

(iv) L to R: 1 (or 1.000 000) [B1]
 R to L: 1.000 001 [B1]
 L to R requires 8 sf, (R to L doesn't) [E1]
 [subtotal 3]

(v) Reverse order more accurate [E1]
 as that way allows the very small terms at the end of the series
 to contribute to the sum. [E1]

The spreadsheet is likely to work to greater accuracy [E1]
 The spreadsheet works to more sf than are displayed [E1]
 [subtotal 4]

[TOTAL 18]

Grade Thresholds

**Advanced GCE (Subject) (Aggregation Code(s))
January 2008 Examination Series**

Unit Threshold Marks

Unit		Maximum Mark	A	B	C	D	E	U
All units	UMS	100	80	70	60	50	40	0
4751	Raw	72	54	46	38	31	24	0
4752	Raw	72	55	48	41	34	28	0
4753	Raw	72	57	50	43	36	28	0
4753/02	Raw	18	15	13	11	9	8	0
4754	Raw	90	77	68	59	50	41	0
4755	Raw	72	55	47	39	32	25	0
4756	Raw	72	59	51	44	37	30	0
4758	Raw	72	62	54	46	38	30	0
4758/02	Raw	18	15	13	11	9	8	0
4761	Raw	72	60	52	44	37	30	0
4762	Raw	72	61	53	45	37	30	0
4763	Raw	72	58	51	44	37	30	0
4766/ G241	Raw	72	56	49	42	35	28	0
4767	Raw	72	62	54	46	38	31	0
4768	Raw	72	54	47	40	33	27	0
4771	Raw	72	60	53	46	39	33	0
4776	Raw	72	58	50	42	35	27	0
4776/02	Raw	18	14	12	10	8	7	0

Specification Aggregation Results

Overall threshold marks in UMS (ie after conversion of raw marks to uniform marks)

	Maximum Mark	A	B	C	D	E	U
7895-7898	600	480	420	360	300	240	0
3895-3898	300	240	210	180	150	120	0

The cumulative percentage of candidates awarded each grade was as follows:

	A	B	C	D	E	U	Total Number of Candidates
7895	25.5	50.0	75.5	85.9	95.3	100	106
7896	42.9	85.7	85.7	85.7	85.7	100	7
7897							0
7898							0
3895	22.7	40.7	59.3	77.8	94.8	100	383
3896	80	80	95	95	100	100	20
3897	0	100	100	100	100	100	1
3898	56.4	76.9	87.2	97.4	97.4	100	39

556 candidates aggregated this series

For a description of how UMS marks are calculated see:

http://www.ocr.org.uk/learners/ums_results.html

Statistics are correct at the time of publication.

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)
Telephone: 01223 553998
Facsimile: 01223 552627
Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity



OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553