

# Mathematics (MEI)

Advanced GCE A2 7895-8

Advanced Subsidiary GCE AS 3895-8

## Mark Schemes for the Units

---

**June 2008**

**3895-8/7895-8/MS/R/08**

OCR (Oxford, Cambridge and RSA Examinations) is a unitary awarding body, established by the University of Cambridge Local Examinations Syndicate and the RSA Examinations Board in January 1998. OCR provides a full range of GCSE, A level, GNVQ, Key Skills and other qualifications for schools and colleges in the United Kingdom, including those previously provided by MEG and OCEAC. It is also responsible for developing new syllabuses to meet national requirements and the needs of students and teachers.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2008

Any enquiries about publications should be addressed to:

OCR Publications  
PO Box 5050  
Annesley  
NOTTINGHAM  
NG15 0DL

Telephone: 0870 770 6622  
Facsimile: 01223 552310  
E-mail: [publications@ocr.org.uk](mailto:publications@ocr.org.uk)

## CONTENTS

**Advanced GCE Further Mathematics (MEI) (7896)**  
**Advanced GCE Further Mathematics (Additional) (MEI) (7897)**  
**Advanced GCE Mathematics (MEI) (7895)**  
**Advanced GCE Pure Mathematics (MEI) (7898)**

**Advanced Subsidiary GCE Further Mathematics (MEI) (3896)**  
**Advanced Subsidiary GCE Further Mathematics (Additional) (MEI) (3897)**  
**Advanced Subsidiary GCE Mathematics (MEI) (3895)**  
**Advanced Subsidiary GCE Pure Mathematics (MEI) (3898)**

### MARK SCHEME FOR THE UNITS

<b>Unit/Content</b>	<b>Page</b>
4751 (C1) Introduction to Advanced Mathematics	1
4752 (C2) Concepts for Advanced Mathematics	6
4753 (C3) Methods for Advanced Mathematics	8
4754 (C4) Applications of Advanced Mathematics	12
4755 (FP1) Further Concepts for Advanced Mathematics	19
4756 (FP2) Further Methods for Advanced Mathematics	25
4757 (FP3) Further Applications of Advanced Mathematics	31
4758 Differential Equations	38
4761 Mechanics 1	42
4762 Mechanics 2	47
4763 Mechanics 3	51
4764 Mechanics 4	57
4766 Statistics 1	60
4767 Statistics 2	67
4768 Statistics 3	71
4769 Statistics 4	77
4771 Decision Mathematics 1	83
4772 Decision Mathematics 2	88
4773 Decision Mathematics Computation	93
4776 Numerical Methods	98
4777 Numerical Computation	101
Grade Thresholds	105

# 4751 (C1) Introduction to Advanced Mathematics

## Section A

1	$x > 6/4$ o.e. isw	2	M1 for $4x > 6$ or for $6/4$ o.e. found or for their final ans ft their $4x > k$ or $kx > 6$	2
2	(i) (0, 4) and (6, 0)	2	1 each; allow $x = 0, y = 4$ etc; condone $x = 6, y = 4$ isw but 0 for (6, 4) with no working	4
	(ii) $-4/6$ o.e. or ft their (i) isw	2	1 for $-\frac{4}{6}x$ or $4/-6$ or $4/6$ o.e. or ft (accept 0.67 or better) 0 for just rearranging to $y = -\frac{2}{3}x + 4$	
3	(i) 0 or $-3/2$ o.e.	2	1 each	5
	(ii) $k < -9/8$ o.e. www	3	M2 for $3^2(-)(-8k) < 0$ o.e. or $-9/8$ found or M1 for attempted use of $b^2 - 4ac$ (may be in quadratic formula); SC: allow M1 for $9 - 8k < 0$ and M1 ft for $k > 9/8$	
4	(i) T (ii) E (iii) T (iv) F	3	3 for all correct, 2 for 3 correct. 1 for 2 correct	3
5	$y(x - 2) = (x + 3)$	M1	for multiplying by $x - 2$ ; condone missing brackets	4
	$xy - 2y = x + 3$ or ft [ft from earlier errors if of comparable difficulty – no ft if there are no $xy$ terms]	M1	for expanding bracket and being at stage ready to collect $x$ terms	
	$xy - x = 2y + 3$ or ft	M1	for collecting $x$ and 'other' terms on opposite sides of eqn	
	$[x = ] \frac{2y+3}{y-1}$ o.e. or ft	M1	for factorising and division	
	<u>alt method:</u> $y = 1 + \frac{5}{x-2}$	M1	for either method: award 4 marks only if fully correct	
	$y - 1 = \frac{5}{x-2}$	M1		
$x - 2 = \frac{5}{y-1}$	M1			
$x = 2 + \frac{5}{y-1}$	M1			

6	(i) 5 www  (ii) $8x^{10}y^{13}z^4$ or $2^3x^{10}y^{13}z^4$	2  3	allow 2 for $\pm 5$ ; M1 for $25^{1/2}$ seen or for $1/5$ seen or for using $25^{1/2} = 5$ with another error (ie M1 for coping correctly with fraction and negative index or with square root)  mark final answer; B2 for 3 elements correct, B1 for 2 elements correct; condone multn signs included, but -1 from total earned if addn signs	5
7	(i) $\frac{5-\sqrt{3}}{22}$ or $\frac{5+(-1)\sqrt{3}}{22}$ or $\frac{5-1\sqrt{3}}{22}$  (ii) $37 - 12\sqrt{7}$ isw www	2  3	or $a = 5, b = -1, c = 22$ ; M1 for attempt to multiply numerator and denominator by $5 - \sqrt{3}$  2 for 37 and 1 for $-12\sqrt{7}$ or M1 for 3 correct terms from $9 - 6\sqrt{7} - 6\sqrt{7} + 28$ or $9 - 3\sqrt{28} - 3\sqrt{28} + 28$ or $9 - \sqrt{252} - \sqrt{252} + 28$ o.e. eg using $2\sqrt{63}$ or M2 for $9 - 12\sqrt{7} + 28$ or $9 - 6\sqrt{28} + 28$ or $9 - 2\sqrt{252} + 28$ or $9 - \sqrt{1008} + 28$ o.e.; 3 for $37 - \sqrt{1008}$ but not other equivs	5
8	-2000 www	4	M3 for $10 \times 5^2 \times (-2[x])^3$ o.e. or M2 for two of these elements or M1 for 10 or $(5 \times 4 \times 3)/(3 \times 2 \times 1)$ o.e. used [ ${}^5C_3$ is not sufficient] or for 1 5 10 10 5 1 seen;  or B3 for 2000;  condone $x^3$ in ans;  equivs: M3 for e.g $5^5 \times 10 \times \left(-\frac{2}{5}[x]\right)^3$  o.e. [ $5^5$ may be outside a bracket for whole expansion of all terms], M2 for two of these elements etc  similarly for factor of 2 taken out at start	4
9	$(y - 3)(y - 4) [= 0]$  $y = 3$ or 4 cao  $x = \pm\sqrt{3}$ or $\pm 2$ cao	M1  A1  B2	for factors giving two terms correct or attempt at quadratic formula or completing square  or B2 (both roots needed)  B1 for 2 roots correct or ft their y (condone $\sqrt{3}$ and $\sqrt{4}$ for B1)	4

## Section B

10	i	$(x - 3)^2 - 7$	3	mark final answer; 1 for $a = 3$ , 2 for $b = 7$ or M1 for $-3^2 + 2$ ; bod 3 for $(x - 3) - 7$	3
	ii	$(3, -7)$ or ft from (i)	1+1		2
	iii	sketch of quadratic correct way up and through $(0, 2)$	G1	accept $(0, 2)$ o.e. seen in this part [eg in table] if 2 not marked as intercept on graph	2
		t.p. correct or ft from (ii)	G1	accept 3 and $-7$ marked on axes level with turning pt., or better; no ft for $(0, 2)$ as min	
iv	$x^2 - 6x + 2 = 2x - 14$ o.e.	M1	or their (i) = $2x - 14$	5	
	$x^2 - 8x + 16 [= 0]$	M1	dep on first M1; condone one error		
	$(x - 4)^2 [= 0]$	M1	or correct use of formula, giving equal roots; allow $(x + 4)^2$ o.e. ft $x^2 + 8x + 16$		
	$x = 4, y = -6$	A1	if M0M0M0, allow SC2 for showing $(4, -6)$ is on both graphs (need to go on to show line is tgt to earn more)		
		equal/repeated roots [implies tgt] - must be explicitly stated; condone 'only one root [so tgt]' or 'line meets curve only once, so tgt' or 'line touches curve only once' etc]	A1	or for use of calculus to show grad of line and curve are same when $x = 4$	
					12

11	i	f(-4) used	M1		2	
		$-128 + 112 + 28 - 12 [= 0]$	A1	or B2 for $(x + 4)(2x^2 - x - 3)$ here; or correct division with no remainder		
	ii	division of $f(x)$ by $(x + 4)$	M1	as far as $2x^3 + 8x^2$ in working, or two terms of $2x^2 - x - 3$ obtained by inspection etc (may be earned in (i)), or $f(-1) = 0$ found	4	
		$2x^2 - x - 3$	A1	$2x^2 - x - 3$ seen implies M1A1		
		$(x + 1)(2x - 3)$	A1			
		$[f(x) =] (x + 4)(x + 1)(2x - 3)$	A1	or B4; allow final A1 ft their factors if M1A1A0 earned		
	iii	sketch of cubic correct way up	G1	ignore any graph of $y = f(x - 4)$	3	
		through $-12$ shown on $y$ axis	G1	or coords stated near graph		
		roots $-4, -1, 1.5$ or ft shown on $x$ axis	G1	or coords stated near graph  if no curve drawn, but intercepts marked on axes, can earn max of G0G1G1		
	iv	$x(x - 3)(2[x - 4] - 3)$ o.e. or $x(x - 3)(x - 5.5)$ or ft their factors	M1	or $2(x - 4)^3 + 7(x - 4)^2 - 7(x - 4) - 12$ or stating roots are 0, 3 and 5.5 or ft; condone one error eg $2x - 7$ not $2x - 11$	3	
correct expansion of one pair of brackets ft from their factors		M1	or for correct expn of $(x - 4)^3$ [allow unsimplified]; or for showing $g(0) = g(3) = g(5.5) = 0$ in given ans $g(x)$			
correct completion to given answer		M1	allow M2 for working backwards from given answer to $x(x - 3)(2x - 11)$ and M1 for full completion with factors or roots			
					3	12

12	i	grad AB = $\frac{9-1}{3--1}$ or 2	M1		3
		$y - 9 = 2(x - 3)$ or $y - 1 = 2(x + 1)$	M1	ft their $m$ , or subst coords of A or B in $y = \text{their } m x + c$	
		$y = 2x + 3$ o.e.	A1	or B3	
	ii	mid pt of AB = (1, 5)	M1	condone not stated explicitly, but used in eqn	4
		grad perp = $-1/\text{grad AB}$	M1	soi by use eg in eqn	
		$y - 5 = -\frac{1}{2}(x - 1)$ o.e. or ft [no ft for just grad AB used]	M1	ft their grad and/or midpt, but M0 if their midpt not used; allow M1 for $y = -\frac{1}{2}x + c$ and then their midpt subst	
		at least one correct interim step towards given answer $2y + x = 11$ , and correct completion NB ans $2y + x = 11$ given	M1	no ft; correct eqn only	
		<u>alt method working back from ans:</u>		mark one method or the other, to benefit of cand, not a mixture	
		$y = \frac{11-x}{2}$ o.e.	M1		
	iii	grad perp = $-1/\text{grad AB}$ and showing/stating same as given line	M1	eg stating $-\frac{1}{2} \times 2 = -1$	2
		finding intn of their $y = 2x + 3$ and $2y + x = 11$ [= (1, 5)]	M1	or showing that (1, 5) is on $2y + x = 11$ , having found (1, 5) first	
		showing midpt of AB is (1, 5)	M1	[for both methods: for M4 must be fully correct]	
showing $(-1 - 5)^2 + (1 - 3)^2 = 40$		M1	at least one interim step needed for each mark; M0 for just $6^2 + 2^2 = 40$		
showing B to centre = $\sqrt{40}$ or verifying that (3, 9) fits given circle		M1	with no other evidence such as a first line of working or a diagram; condone marks earned in reverse order		
iv	$(x - 5)^2 + 3^2 = 40$	M1	for subst $y = 0$ in circle eqn	3	
	$(x - 5)^2 = 31$	M1	condone slip on rhs; or for rearrangement to zero (condone one error) <u>and</u> attempt at quad. formula [allow M1 M0 for $(x - 5)^2 = 40$ or for $(x - 5)^2 + 3^2 = 0$ ]		
	$x = 5 \pm \sqrt{31}$ or $\frac{10 \pm \sqrt{124}}{2}$ isw	A1	or $5 \pm \frac{\sqrt{124}}{2}$		



# 4752 (C2) Concepts for Advanced Mathematics

## Section A

1	210 c.a.o.	2	1 for $\pi$ rads = $180^\circ$ soi	2
2	(i) $5.4 \times 10^{-3}$ , 0.0054 or $\frac{27}{5000}$ (ii) 6 www	1 2	M1 for $S = 5.4 / (1 - 0.1)$	3
3	stretch, parallel to the $y$ axis, sf 3	2	1 for stretch plus one other element correct	2
4	[f'(x) =] $12 - 3x^2$ their $f'(x) > 0$ or $= 0$ soi $-2 < x < 2$	B1 M1 A1	condone $-2 \leq x \leq 2$ or "between -2 and 2"	3
5	(i) grad of chord = $(2^{3.1} - 2^3)/0.1$ o.e. = 5.74 c.a.o. (ii) correct use of A and C where for C, $2.9 < x < 3.1$ answer in range (5.36, 5.74)	M1 A1  M1 A1	or chord with ends $x = 3 \pm h$ , where $0 < h \leq 0.1$ s.c.1 for consistent use of reciprocal of gradient formula in parts (i) and (ii)	4
6	[y =] $kx^{3/2}$ [+ c] $k = 4$ subst of (9, 105) in their eqn with c  or $c = -3$	M1 A1 M1  A1	may appear at any stage must have c; must have attempted integration	4
7	sector area = 28.8 or $\frac{144}{5}$ [cm <sup>2</sup> ] c.a.o. area of triangle = $\frac{1}{2} \times 6^2 \times \sin 1.6$ o.e. their sector – their triangle s.o.i. 10.8 to 10.81 [cm <sup>2</sup> ]	2 M1  M1 A1	M1 for $\frac{1}{2} \times 6^2 \times 1.6$  must both be areas leading to a positive answer	5
8	$a + 10d = 1$ or $121 = 5.5(2a+10d)$ $5(2a + 9d) = 120$ o.e. $a = 21$ s.o.i. www and $d = -2$ s.o.i. www 4th term is 15	M1 M1 A1 A1 A1	or $121 = 5.5(a + 1)$ gets M2 eg $2a + 9d = 24$	5
9	$x \log 5 = \log 235$ or $x = \frac{\log 235}{\log 5}$  3.39	M1  A2	or $x = \log_5 235$  A1 for 3.4 or versions of 3.392...	3
10	$2(1 - \cos^2 \theta) = \cos \theta + 2$ $-2 \cos^2 \theta = \cos \theta$ s.o.i. valid attempt at solving their quadratic in $\cos \theta$ $\cos \theta = -\frac{1}{2}$ www $\theta = 90, 270, 120, 240$	M1 A1 DM1  A1 A1	for $1 - \cos^2 \theta = \sin^2 \theta$ substituted graphic calc method: allow M3 for intersection of $y = 2 \sin^2 \theta$ and $y = \cos \theta + 2$ and A2 for all four roots. All four answers correct but unsupported scores B2. 120 and 240 only: B1.	5

18

18


## Section B

11	i	$(x + 5)(x - 2)(x + 2)$	2	M1 for a $(x + 5)(x - 2)(x + 2)$	2
	ii	$[(x + 2)](x^2 + 3x - 10)$	M1	for correct expansion of one pair of their brackets	2
		$x^3 + 3x^2 - 10x + 2x^2 + 6x - 20$ o.e.	M1	for clear expansion of correct factors – accept given answer from $(x + 5)(x^2 - 4)$ as first step	
	iii	$y' = 3x^2 + 10x - 4$ their $3x^2 + 10x - 4 = 0$ s.o.i. $x = 0.36\dots$ from formula o.e.	M2 M1 A1	M1 if one error or M1 for substitution of 0.4 if trying to obtain 0, and A1 for correct demonstration of sign change	6
$(-3.7, 12.6)$		B1+1			
iv	$(-1.8, 12.6)$	B1+1	accept $(-1.9, 12.6)$ or f.t. ( $\frac{1}{2}$ their max x, their max y)	2	
12	i	Area = $(-0.136$ seen $[m^2]$ www	4	M3 for $0.1/2 \times (0.14 + 0.16 + 2[0.22 + 0.31 + 0.36 + 0.32])$ M2 for one slip; M1 for two slips must be positive	5
		Volume = $0.34 [m^3]$ or ft from their area $\times 2.5$	1		
	ii	$2x^4 - x^3 - 0.25x^2 - 0.15x$ o.e. value at 0.5 [- value at 0] $= -0.1375$ area of cross section (of trough) or area between curve and x-axis $0.34375$ r.o.t. to 3 or more sf $[m^3]$ $m^3$ seen in (i) or (ii)	M2 M1 A1  E1 B1 U1	M1 for 2 terms correct dep on integral attempted must have neg sign	7
13	i	$\log P = \log a + b \log t$ www	1	must be with correct equation condone omission of base	3
		comparison with $y = mx + c$	1		
		intercept = $\log_{10} a$	1		
	ii	$\log t$ 0 0.78 1.15 1.18 1.20	1 1	accept to 2 or more dp	4
$\log P$ 1.49 1.64 1.75 1.74 1.76		1 1			
iii	plots f.t. ruled line of best fit gradient rounding to 0.22 or 0.23 $a = 10^{1.49}$ s.o.i. $P = 31t^m$ allow the form $P = 10^{0.22 \log t + 1.49}$	2 1 1	M1 for y step / x-step accept 1.47 – 1.50 for intercept accept answers that round to 30 – 32, their positive m	4	
iv	answer rounds in range 60 to 63	1		1	


# 4753 (C3) Methods for Advanced Mathematics


## Section A

<p><b>1</b> <math> 2x-1  \leq 3</math>  <math>\Rightarrow -3 \leq 2x-1 \leq 3</math>  <math>\Rightarrow -2 \leq 2x \leq 4</math>  <math>\Rightarrow -1 \leq x \leq 2</math>  <i>or</i>  <math>(2x-1)^2 \leq 9</math>  <math>\Rightarrow 4x^2 - 4x - 8 \leq 0</math>  <math>\Rightarrow (4)(x+1)(x-2) \leq 0</math>  <math>\Rightarrow -1 \leq x \leq 2</math></p>	<p>M1 A1 M1 A1  M1 A1 A1 A1 [4]</p>	<p><math>2x-1 \leq 3</math> (or =)  <math>x \leq 2</math>  <math>2x-1 \geq -3</math> (or =)  <math>x \geq -1</math>             squaring and forming quadratic = 0 (or <math>\leq</math>)            factorising or solving to get <math>x = -1, 2</math>  <math>x \geq -1</math>  <math>x \leq 2</math> (www)</p>
<p><b>2</b> Let <math>u = x</math>, <math>dv/dx = e^{3x} \Rightarrow v = e^{3x}/3</math>  <math>\Rightarrow \int x e^{3x} dx = \frac{1}{3} x e^{3x} - \int \frac{1}{3} e^{3x} \cdot 1 dx</math>  <math>= \frac{1}{3} x e^{3x} - \frac{1}{9} e^{3x} + c</math></p>	<p>M1 A1 A1 B1 [4]</p>	<p>parts with <math>u = x</math>, <math>dv/dx = e^{3x} \Rightarrow v</math>   <math>= \frac{1}{3} x e^{3x} - \frac{1}{9} e^{3x} + c</math></p>
<p><b>3 (i)</b> <math>f(-x) = f(x)</math> Symmetrical about Oy.</p>	<p>B1 B1 [2]</p>	
<p><b>(ii)</b> (A) even (B) neither (C) odd</p>	<p>B1 B1 B1 [3]</p>	
<p><b>4</b> Let <math>u = x^2 + 2 \Rightarrow du = 2x dx</math>  <math>\int_1^4 \frac{x}{x^2+2} dx = \int_3^{18} \frac{1/2}{u} du</math>  <math>= \frac{1}{2} [\ln u]_3^{18}</math>  <math>= \frac{1}{2} (\ln 18 - \ln 3)</math>  <math>= \frac{1}{2} \ln(18/3)</math>  <math>= \frac{1}{2} \ln 6^*</math></p>	<p>M1 A1 M1 E1 [4]</p>	<p><math>\int \frac{1/2}{u} du</math> or <math>k \ln(x^2 + 2)</math>  <math>\frac{1}{2} \ln u</math> or <math>\frac{1}{2} \ln(x^2 + 2)</math>            substituting correct limits (<math>u</math> or <math>x</math>)            must show working for <math>\ln 6</math></p>
<p><b>5</b> <math>y = x^2 \ln x</math>  <math>\Rightarrow \frac{dy}{dx} = x^2 \cdot \frac{1}{x} + 2x \ln x</math>  <math>= x + 2x \ln x</math>  <math>dy/dx = 0</math> when <math>x + 2x \ln x = 0</math>  <math>\Rightarrow x(1 + 2 \ln x) = 0</math>  <math>\Rightarrow \ln x = -\frac{1}{2}</math>  <math>\Rightarrow x = e^{-\frac{1}{2}} = 1/\sqrt{e}^*</math></p>	<p>M1 B1 A1  M1 M1 E1 [6]</p>	<p>product rule  <math>d/dx (\ln x) = 1/x</math> soi            oe             their deriv = 0 or attempt to verify  <math>\ln x = -\frac{1}{2} \Rightarrow x = e^{-\frac{1}{2}}</math> or <math>\ln(1/\sqrt{e}) = -\frac{1}{2}</math></p>

<b>6(i)</b> Initial mass = $20 + 30 e^0 = 50$ grams Long term mass = 20 grams	M1A1 B1 [3]	
<b>(ii)</b> $30 = 20 + 30 e^{-0.1t}$ $\Rightarrow e^{-0.1t} = 1/3$ $\Rightarrow -0.1t = \ln(1/3) = -1.0986\dots$ $\Rightarrow t = 11.0$ mins	M1  M1  A1 [3]	anti-logging correctly  11, 11.0, 10.99, 10.986 (not more than 3 d.p)
<b>(iii)</b> 	B1  B1 [2]	correct shape through (0, 50) – ignore negative values of $t$ $\rightarrow 20$ as $t \rightarrow \infty$
<b>7</b> $x^2 + xy + y^2 = 12$ $\Rightarrow 2x + x \frac{dy}{dx} + y + 2y \frac{dy}{dx} = 0$ $\Rightarrow (x + 2y) \frac{dy}{dx} = -2x - y$ $\Rightarrow \frac{dy}{dx} = -\frac{2x + y}{(x + 2y)}$	M1 B1  A1  M1  A1 [5]	Implicit differentiation $x \frac{dy}{dx} + y$ correct equation  collecting terms in $dy/dx$ and factorising  oe cao

Section B

<p><b>8(i)</b> <math>y = 1/(1+\cos\pi/3) = 2/3.</math></p>	<p>B1 [1]</p>	<p>or 0.67 or better</p>
<p><b>(ii)</b> <math>f'(x) = -1(1 + \cos x)^{-2} \cdot -\sin x</math>  <math>= \frac{\sin x}{(1 + \cos x)^2}</math>                  When <math>x = \pi/3</math>, <math>f'(x) = \frac{\sin(\pi/3)}{(1 + \cos(\pi/3))^2}</math>  <math>= \frac{\sqrt{3}/2}{(1 + 1/2)^2} = \frac{\sqrt{3}}{2} \times \frac{4}{9} = \frac{2\sqrt{3}}{9}</math></p>	<p>M1 B1 A1  M1 A1  [5]</p>	<p>chain rule or quotient rule  <math>d/dx (\cos x) = -\sin x</math> so i                  correct expression                   substituting <math>x = \pi/3</math>                   oe or 0.38 or better. (0.385, 0.3849)</p>
<p><b>(iii)</b> deriv = <math>\frac{(1 + \cos x)\cos x - \sin x \cdot (-\sin x)}{(1 + \cos x)^2}</math>  <math>= \frac{\cos x + \cos^2 x + \sin^2 x}{(1 + \cos x)^2}</math>  <math>= \frac{\cos x + 1}{(1 + \cos x)^2}</math>  <math>= \frac{1}{1 + \cos x} *</math>                  Area = <math>\int_0^{\pi/3} \frac{1}{1 + \cos x} dx</math>  <math>= \left[ \frac{\sin x}{1 + \cos x} \right]_0^{\pi/3}</math>  <math>= \frac{\sin \pi/3}{1 + \cos \pi/3} - (-0)</math>  <math>= \frac{\sqrt{3}}{2} \times \frac{2}{3} = \frac{\sqrt{3}}{3}</math></p>	<p>M1  A1 M1dep E1  B1  M1 A1 cao [7]</p>	<p>Quotient or product rule –                  condone <math>uv' - u'v</math> for M1                   correct expression   <math>\cos^2 x + \sin^2 x = 1</math> used dep M1                   www                   substituting limits                   or <math>1/\sqrt{3}</math> - must be exact</p>
<p><b>(iv)</b> <math>y = 1/(1 + \cos x) \quad x \leftrightarrow y</math>  <math>x = 1/(1 + \cos y)</math>  <math>\Rightarrow 1 + \cos y = 1/x</math>  <math>\Rightarrow \cos y = 1/x - 1</math>  <math>\Rightarrow y = \arccos(1/x - 1) *</math>                   Domain is <math>1/2 \leq x \leq 1</math>   </p>	<p>M1  A1 E1  B1  B1  [5]</p>	<p>attempt to invert equation                    www                    reasonable reflection in <math>y = x</math></p>

<p><b>9 (i)</b> <math>y = \sqrt{4 - x^2}</math>  <math>\Rightarrow y^2 = 4 - x^2</math>  <math>\Rightarrow x^2 + y^2 = 4</math>          which is equation of a circle centre O radius 2          Square root does not give negative values, so this is only a semi-circle.</p>	<p>M1 A1 B1 [3]</p>	<p>squaring <math>x^2 + y^2 = 4</math> + comment (correct) oe, e.g. f is a function and therefore single valued</p>
<p><b>(ii)</b> (A) Grad of OP = <math>b/a</math>  <math>\Rightarrow</math> grad of tangent = <math>-\frac{a}{b}</math></p> <p>(B) <math>f(x) = \frac{1}{2}(4 - x^2)^{-1/2} \cdot (-2x)</math>  <math>= -\frac{x}{\sqrt{4 - x^2}}</math>  <math>\Rightarrow f'(a) = -\frac{a}{\sqrt{4 - a^2}}</math></p> <p>(C) <math>b = \sqrt{4 - a^2}</math>          so <math>f'(a) = -\frac{a}{b}</math> as before</p>	<p>M1 A1 M1 A1 B1 E1 [6]</p>	<p>chain rule or implicit differentiation oe substituting <math>a</math> into their <math>f'(x)</math></p>
<p><b>(iii)</b> Translation through <math>\begin{pmatrix} 2 \\ 0 \end{pmatrix}</math> followed by stretch scale factor 3 in <math>y</math>-direction</p> 	<p>M1 A1 M1 A1 A1 [6]</p>	<p>Translation in <math>x</math>-direction through <math>\begin{pmatrix} 2 \\ 0 \end{pmatrix}</math> or 2 to right ('shift', 'move' M1 A0)  <math>\begin{pmatrix} 2 \\ 0 \end{pmatrix}</math> alone is SC1          stretch in <math>y</math>-direction (condone <math>y</math> 'axis') (scale) factor 3          elliptical (or circular) shape through (0, 0) and (4, 0) and (2, 6) (soi)          -1 if whole ellipse shown</p>
<p><b>(iv)</b> <math>y = 3f(x - 2)</math>  <math>= 3\sqrt{4 - (x - 2)^2}</math>  <math>= 3\sqrt{4 - x^2 + 4x - 4}</math>  <math>= 3\sqrt{4x - x^2}</math>  <math>\Rightarrow y^2 = 9(4x - x^2)</math>  <math>\Rightarrow 9x^2 + y^2 = 36x</math> *</p>	<p>M1 A1 E1 [3]</p>	<p>or substituting <math>3\sqrt{4 - (x - 2)^2}</math> oe for <math>y</math> in <math>9x^2 + y^2</math>  <math>4x - x^2</math>          www</p>

# 4754 (C4) Applications of Advanced Mathematics

## Section A

<b>1</b> $\frac{x}{x^2-4} + \frac{2}{x+2} = \frac{x}{(x-2)(x+2)} + \frac{2}{x+2}$ $= \frac{x+2(x-2)}{(x+2)(x-2)}$ $= \frac{3x-4}{(x+2)(x-2)}$	M1 M1 A1 [3]	combining fractions correctly factorising and cancelling (may be $3x^2+2x-8$ )
<b>2</b> $V = \int_0^1 \pi y^2 dx = \int_0^1 \pi(1+e^{2x}) dx$ $= \pi \left[ x + \frac{1}{2} e^{2x} \right]_0^1$ $= \pi \left( 1 + \frac{1}{2} e^2 - \frac{1}{2} \right)$ $= \frac{1}{2} \pi (1+e^2) *$	M1 B1 M1 E1 [4]	must be $\pi x$ their $y^2$ in terms of $x$ $\left[ x + \frac{1}{2} e^{2x} \right]$ only substituting both $x$ limits in a function of $x$ www
<b>3</b> $\cos 2\theta = \sin \theta$ $\Rightarrow 1 - 2\sin^2 \theta = \sin \theta$ $\Rightarrow 1 - \sin \theta - 2\sin^2 \theta = 0$ $\Rightarrow (1 - 2\sin \theta)(1 + \sin \theta) = 0$ $\Rightarrow \sin \theta = \frac{1}{2} \text{ or } -1$ $\Rightarrow \theta = \pi/6, 5\pi/6, 3\pi/2$	M1 M1 A1 M1 A1 A2,1,0 [7]	$\cos 2\theta = 1 - 2\sin^2 \theta$ oe substituted forming quadratic (in one variable) = 0 correct quadratic www factorising or solving quadratic $\frac{1}{2}, -1$ oe www cao penalise extra solutions in the range
<b>4</b> $\sec \theta = x/2, \tan \theta = y/3$ $\sec^2 \theta = 1 + \tan^2 \theta$ $\Rightarrow x^2/4 = 1 + y^2/9$ $\Rightarrow x^2/4 - y^2/9 = 1 *$ <p><b>OR</b> <math>x^2/4 - y^2/9 = 4\sec^2 \theta/4 - 9\tan^2 \theta/9</math>  <math>= \sec^2 \theta - \tan^2 \theta = 1</math></p>	M1 M1 E1 [3]	$\sec^2 \theta = 1 + \tan^2 \theta$ used (oe, e.g. converting to sines and cosines and using $\cos^2 \theta + \sin^2 \theta = 1$ ) eliminating $\theta$ (or $x$ and $y$ ) www
<b>5(i)</b> $dx/du = 2u, dy/du = 6u^2$ $\Rightarrow \frac{dy}{dx} = \frac{dy/du}{dx/du} = \frac{6u^2}{2u} = 3u$ <p><b>OR</b> <math>y = 2(x-1)^{3/2}, dy/dx = 3(x-1)^{1/2} = 3u</math></p>	B1 M1 A1 [3]	both $2u$ and $6u^2$ B1 ( $y=f(x)$ ), M1 differentiation, A1
<b>(ii)</b> At (5, 16), $u = 2$ $\Rightarrow dy/dx = 6$	M1 A1 [2]	cao

<p>6(i) <math>(1+4x^2)^{-\frac{1}{2}} = 1 - \frac{1}{2} \cdot 4x^2 + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!} (4x^2)^2 + \dots</math>  <math>= 1 - 2x^2 + 6x^4 + \dots</math></p> <p>Valid for <math>-1 &lt; 4x^2 &lt; 1 \Rightarrow -\frac{1}{2} &lt; x &lt; \frac{1}{2}</math></p>	<p>M1 A1 A1 M1A1 [5]</p>	<p>binomial expansion with <math>p = -1/2</math></p> <p><math>1 - 2x^2 \dots</math> <math>+ 6x^4</math></p>
<p>(ii) <math>\frac{1-x^2}{\sqrt{1+4x^2}} = (1-x^2)(1-2x^2+6x^4+\dots)</math>  <math>= 1 - 2x^2 + 6x^4 - x^2 + 2x^4 + \dots</math>  <math>= 1 - 3x^2 + 8x^4 + \dots</math></p>	<p>M1 A1 A1 [3]</p>	<p>substituting their <math>1 - 2x^2 + 6x^4 + \dots</math> and expanding</p> <p>ft their expansion (of three terms)</p> <p>cao</p>
<p>7 <math>\sqrt{3} \sin x - \cos x = R \sin(x - \alpha)</math>  <math>= R(\sin x \cos \alpha - \cos x \sin \alpha)</math>  <math>\Rightarrow \sqrt{3} = R \cos \alpha, 1 = R \sin \alpha</math>  <math>\Rightarrow R^2 = 3 + 1 = 4 \Rightarrow R = 2</math>  <math>\tan \alpha = 1/\sqrt{3}</math>  <math>\Rightarrow \alpha = \pi/6</math>  <math>\Rightarrow y = 2 \sin(x - \pi/6)</math></p> <p>Max when <math>x - \pi/6 = \pi/2 \Rightarrow x = \pi/6 + \pi/2 = 2\pi/3</math>  max value <math>y = 2</math></p> <p>So maximum is <math>(2\pi/3, 2)</math></p>	<p>M1 B1 M1 A1  B1 B1  [6]</p>	<p>correct pairs soi  <math>R = 2</math>  ft  cao www</p> <p>cao  ft their <math>R</math></p> <p>SC B1 <math>(2, 2\pi/3)</math> no working</p>



## Section B

<p><b>8(i)</b> At A: <math>3 \times 0 + 2 \times 0 + 20 \times (-15) + 300 = 0</math>          At B: <math>3 \times 100 + 2 \times 0 + 20 \times (-30) + 300 = 0</math>          At C: <math>3 \times 0 + 2 \times 100 + 20 \times (-25) + 300 = 0</math>          So ABC has equation <math>3x + 2y + 20z + 300 = 0</math></p>	<p>M1          A2,1,0          [3]</p>	<p>substituting co-ords into equation of plane...          for ABC  <b>OR</b> using two vectors in the plane form vector          product M1A1 then  <math>3x + 2y + 20z = c = -300</math> A1  <b>OR</b> using vector equation of plane M1, elim          both parameters M1, A1</p>
<p><b>(ii)</b> <math>\overline{DE} = \begin{pmatrix} 100 \\ 0 \\ -10 \end{pmatrix}</math> <math>\overline{DF} = \begin{pmatrix} 0 \\ 100 \\ 5 \end{pmatrix}</math></p> <p><math>\begin{pmatrix} 100 \\ 0 \\ -10 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ -1 \\ 20 \end{pmatrix} = 100 \times 2 + 0 \times -1 + -10 \times 20 = 200 - 200 = 0</math></p> <p><math>\begin{pmatrix} 0 \\ 100 \\ 5 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ -1 \\ 20 \end{pmatrix} = 0 \times 2 + 100 \times -1 + 5 \times 20 = -100 + 100 = 0</math></p> <p>Equation of plane is <math>2x - y + 20z = c</math>          At D (say) <math>c = 20 \times -40 = -800</math>          So equation is <math>2x - y + 20z + 800 = 0</math></p>	<p>B1B1          B1          B1          M1          A1          [6]</p>	<p>need evaluation          need evaluation</p>
<p><b>(iii)</b> Angle is <math>\theta</math>, where</p> $\cos \theta = \frac{\begin{pmatrix} 2 \\ -1 \\ 20 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 2 \\ 20 \end{pmatrix}}{\sqrt{2^2 + (-1)^2 + 20^2} \sqrt{3^2 + 2^2 + 20^2}} = \frac{404}{\sqrt{405} \sqrt{413}}$ <p><math>\Rightarrow \theta = 8.95^\circ</math></p>	<p>M1          A1          A1          A1cao          [4]</p>	<p>formula with correct vectors          top          bottom          (or 0.156 radians)</p>
<p><b>(iv)</b> RS: <math>\mathbf{r} = \begin{pmatrix} 15 \\ 34 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 3 \\ 2 \\ 20 \end{pmatrix}</math></p> $= \begin{pmatrix} 15 + 3\lambda \\ 34 + 2\lambda \\ 20\lambda \end{pmatrix}$ <p><math>\Rightarrow 3(15 + 3\lambda) + 2(34 + 2\lambda) + 20 \cdot 20\lambda + 300 = 0</math>  <math>\Rightarrow 45 + 9\lambda + 68 + 4\lambda + 400\lambda + 300 = 0</math>  <math>\Rightarrow 413 + 413\lambda = 0</math>  <math>\Rightarrow \lambda = -1</math>          so S is (12, 32, -20)</p>	<p>B1          B1          M1          A1          A1          [5]</p>	<p><math>\begin{pmatrix} 15 \\ 34 \\ 0 \end{pmatrix} + \dots</math>  <math>\dots + \lambda \begin{pmatrix} 3 \\ 2 \\ 20 \end{pmatrix}</math>          solving with plane  <math>\lambda = -1</math>          cao</p>

<p>9(i) <math>v = \int 10e^{-\frac{1}{2}t} dt</math>  <math>= -20e^{-\frac{1}{2}t} + c</math>          when <math>t = 0, v = 0</math>  <math>\Rightarrow 0 = -20 + c</math>  <math>\Rightarrow c = 20</math>          so <math>v = 20 - 20e^{-\frac{1}{2}t}</math></p>	<p>M1 A1 M1  A1 [4]</p>	<p>separate variables and intend to integrate  <math>-20e^{-\frac{1}{2}t}</math>          finding <math>c</math>            cao</p>
<p>(ii) As <math>t \rightarrow \infty</math> <math>e^{-1/2t} \rightarrow 0</math>  <math>\Rightarrow v \rightarrow 20</math>          So long term speed is <math>20 \text{ m s}^{-1}</math></p>	<p>M1  A1 [2]</p>	<p>ft (for their <math>c &gt; 0</math>, found)</p>
<p>(iii) <math>\frac{1}{(w-4)(w+5)} = \frac{A}{w-4} + \frac{B}{w+5}</math>  <math>= \frac{A(w+5) + B(w-4)}{(w-4)(w+5)}</math>  <math>\Rightarrow 1 \equiv A(w+5) + B(w-4)</math>  <math>w = 4: 1 = 9A \Rightarrow A = 1/9</math>  <math>w = -5: 1 = -9B \Rightarrow B = -1/9</math>  <math>\Rightarrow \frac{1}{(w-4)(w+5)} = \frac{1/9}{w-4} - \frac{1/9}{w+5}</math>  <math>= \frac{1}{9(w-4)} - \frac{1}{9(w+5)}</math></p>	<p>M1  M1 A1 A1  [4]</p>	<p>cover up, substitution or equating coeffs  <math>1/9</math>  <math>-1/9</math></p>
<p>(iv) <math>\frac{dw}{dt} = -\frac{1}{2}(w-4)(w+5)</math>  <math>\Rightarrow \int \frac{dw}{(w-4)(w+5)} = \int -\frac{1}{2} dt</math>  <math>\Rightarrow \int \left[ \frac{1}{9(w-4)} - \frac{1}{9(w+5)} \right] dw = \int -\frac{1}{2} dt</math>  <math>\Rightarrow \frac{1}{9} \ln(w-4) - \frac{1}{9} \ln(w+5) = -\frac{1}{2} t + c</math>  <math>\Rightarrow \frac{1}{9} \ln \frac{w-4}{w+5} = -\frac{1}{2} t + c</math>          When <math>t = 0, w = 10 \Rightarrow c = \frac{1}{9} \ln \frac{6}{15} = \frac{1}{9} \ln \frac{2}{5}</math>  <math>\Rightarrow \ln \frac{w-4}{w+5} = -\frac{9}{2} t + \ln \frac{2}{5}</math>  <math>\Rightarrow \frac{w-4}{w+5} = e^{\frac{9}{2}t + \ln \frac{2}{5}} = \frac{2}{5} e^{\frac{9}{2}t} = 0.4e^{-4.5t} *</math></p>	<p>M1  M1  A1ft  M1  M1  E1 [6]</p>	<p>separating variables          substituting their partial fractions            integrating correctly (condone absence of <math>c</math>)            correctly evaluating <math>c</math> (at any stage)            combining lns (at any stage)            www</p>
<p>(v) As <math>t \rightarrow \infty</math> <math>e^{-4.5t} \rightarrow 0</math>  <math>\Rightarrow w - 4 \rightarrow 0</math>          So long term speed is <math>4 \text{ m s}^{-1}</math>.</p>	<p>M1 A1 [2]</p>	

Comprehension

1. (i)

2	1	3
3	2	1
1	3	2

B1  
cao

(ii)

2	3	1
3	1	2
1	2	3

B1  
cao

2. Dividing the grid up into four 2 x 2 blocks gives

1	2	3	4
3	1	4	2
2	4	1	3
4	3	2	1

Lines drawn on diagram or reference to 2 x 2 blocks. M1

One (or more) block does not contain all 4 of the symbols 1, 2, 3 and 4. oe. E1

3.

1	2	3	4
4	3	1	2
2	1	4	3
3	4	2	1

Many possible answers Row 2 correct

Rest correct B1  
B1

4. Either

4	2	3	1
		2	4
		4	2
2	4	1	3

Or

4	2	3	1
		2	4
		4	2
2	4	1	3

B2

5. In the top row there are 9 ways of allocating a symbol to the left cell, then 8 for the next, 7 for the next and so on down to 1 for the right cell, giving

M1

$$9 \times 8 \times 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1 = 9! \text{ ways.}$$

E1

So there must be  $9! \times$  the number of ways of completing the rest of the puzzle.

6.

(i)

Block side length, $b$	Sudoku, $s \times s$	$M$
1	$1 \times 1$	-
2	$4 \times 4$	12
3	$9 \times 9$	77
4	$16 \times 16$	252
5	$25 \times 25$	621

$25 \times 25$  B1

77, 252 and 621 B1

(ii)

$$M = b^4 - 4$$

$b^4$  B1

- 4 B1

7.

(i)

There are neither 3s nor 5s among the givens.

M1

So they are interchangeable and therefore there is no unique solution

E1

(ii)

The missing symbols form a  $3 \times 3$  embedded Latin square.


M1

There is not a unique arrangement of the numbers 1, 2 and 3 in this square.

E1

[18]


## 4755 (FP1) Further Concepts for Advanced Mathematics


Qu	Answer	Mark	Comment
<b>Section A</b>			
1(i)	$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$	B1	
1(ii)	$\begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}$	B1	
1(iii)	$\begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} -3 & 0 \\ 0 & 3 \end{pmatrix}$	M1 A1 <b>[4]</b>	Multiplication, or other valid method (may be implied) c.a.o.
2		B3  B3  B1  <b>[7]</b>	Circle, B1; centre $-3+2j$ , B1; radius = 2, B1  Line parallel to real axis, B1; through $(0, 2)$ , B1; correct half line, B1  Points $-1+2j$ and $-5+2j$ indicated c.a.o.
3	$\begin{pmatrix} -1 & -1 \\ 2 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix}$ $\Rightarrow -x - y = x, 2x + 2y = y$ $\Rightarrow y = -2x$	M1  M1 B1 <b>[3]</b>	For $\begin{pmatrix} -1 & -1 \\ 2 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix}$
4	$3x^3 - x^2 + 2 \equiv A(x-1)^3 + (x^3 + Bx^2 + Cx + D)$ $\equiv Ax^3 - 3Ax^2 + 3Ax - A + x^3 + Bx^2 + Cx + D$ $\equiv (A+1)x^3 + (B-3A)x^2 + (3A+C)x + (D-A)$ $\Rightarrow A=2, B=5, C=-6, D=4$	M1  B4  <b>[5]</b>	Attempt to compare coefficients  One for each correct value

<p>5(i)</p> $\mathbf{AB} = \begin{pmatrix} 7 & 0 & 0 \\ 0 & 7 & 0 \\ 0 & 0 & 7 \end{pmatrix}$ <p>5(ii)</p> $\mathbf{A}^{-1} = \frac{1}{7} \begin{pmatrix} -1 & 0 & 2 \\ 14 & -14 & 7 \\ -5 & 7 & -4 \end{pmatrix}$		<p>B3 [3]</p> <p>M1 A1 [2]</p>	<p>Minus 1 each error to minimum of 0</p> <p>Use of B c.a.o.</p>
<p>6</p> $w = 2x \Rightarrow x = \frac{w}{2}$ $\Rightarrow 2\left(\frac{w}{2}\right)^3 + \left(\frac{w}{2}\right)^2 - 3\left(\frac{w}{2}\right) + 1 = 0$ $\Rightarrow w^3 + w^2 - 6w + 4 = 0$		<p>B1</p> <p>M1 A1</p> <p>A2</p> <p>[5]</p>	<p>Substitution. For substitution <math>x = 2w</math> give B0 but then follow through for a maximum of 3 marks</p> <p>Substitute into cubic Correct substitution</p> <p>Minus 1 for each error (including '= 0' missing), to a minimum of 0 Give full credit for integer multiple of equation</p>
<p>6</p> <p><b>OR</b></p> $\alpha + \beta + \gamma = -\frac{1}{2}$ $\alpha\beta + \alpha\gamma + \beta\gamma = -\frac{3}{2}$ $\alpha\beta\gamma = -\frac{1}{2}$ <p>Let new roots be <math>k, l, m</math> then</p> $k + l + m = 2(\alpha + \beta + \gamma) = -1 = \frac{-B}{A}$ $kl + km + lm = 4(\alpha\beta + \alpha\gamma + \beta\gamma) = -6 = \frac{C}{A}$ $klm = 8\alpha\beta\gamma = -4 = \frac{-D}{A}$ $\Rightarrow w^3 + w^2 - 6w + 4 = 0$		<p>B1</p> <p>M1 A1 A2</p> <p>[5]</p>	<p>All three</p> <p>Attempt to use sums and products of roots of original equation to find sums and products of roots in related equation Sums and products all correct</p> <p>ft their coefficients; minus one for each error (including '= 0' missing), to minimum of 0 Give full credit for integer multiple of equation</p>

7(i)	$\frac{1}{3r-1} - \frac{1}{3r+2} \equiv \frac{3r+2-(3r-1)}{(3r-1)(3r+2)}$ $\equiv \frac{3}{(3r-1)(3r+2)}$	M1	Attempt at correct method
7(ii)	$\sum_{r=1}^n \frac{1}{(3r-1)(3r+2)} = \frac{1}{3} \sum_{r=1}^n \left[ \frac{1}{3r-1} - \frac{1}{3r+2} \right]$ $= \frac{1}{3} \left[ \left( \frac{1}{2} - \frac{1}{5} \right) + \left( \frac{1}{5} - \frac{1}{8} \right) + \dots + \left( \frac{1}{3n-1} - \frac{1}{3n+2} \right) \right]$ $= \frac{1}{3} \left[ \frac{1}{2} - \frac{1}{3n+2} \right]$	M1 A1 M1 A2 [5]	Correct, without fudging [2] Attempt to use identity Terms in full (at least two) Attempt at cancelling A1 if factor of $\frac{1}{3}$ missing, A1 max if answer not in terms of $n$
<b>Section A Total: 36</b>			




Section B			
8(i)	$x = 3, x = -2, y = 2$	B1 B1 B1 [3]	
8(ii)	Large positive $x, y \rightarrow 2^+$ (e.g. consider $x = 100$ ) Large negative $x, y \rightarrow 2^-$ (e.g. consider $x = -100$ )	M1 B1 B1 [3]	Evidence of method required
8(iii)	Curve Central and RH branches correct Asymptotes correct and labelled LH branch correct, with clear minimum	B1 B1 B1 [3]	
			
8(iv)	$-2 < x < 3$ $x \neq 0$	B2 B1 [3]	B2 max if any inclusive inequalities appear B3 for $-2 < x < 0$ and $0 < x < 3$ ,

<p>9(i)</p>	<p><math>2+2j</math> and <math>-1-j</math></p>	<p>B2 [2]</p>	<p>1 mark for each</p>
<p>9(ii)</p>		<p>B2 [2]</p>	<p>1 mark for each correct pair</p>
<p>9(iii)</p>	<p><math>(x-2-2j)(x-2+2j)(x+1+j)(x+1-j)</math></p> <p><math>= (x^2 - 4x + 8)(x^2 + 2x + 2)</math></p> <p><math>= x^4 + 2x^3 + 2x^2 - 4x^3 - 8x^2 - 8x + 8x^2 + 16x + 16</math></p> <p><math>= x^4 - 2x^3 + 2x^2 + 8x + 16</math></p> <p><math>\Rightarrow A = -2, B = 2, C = 8, D = 16</math></p> <p><b>OR</b></p> <p><math>\sum \alpha = 2</math>  <math>\alpha\beta\gamma\delta = 16</math>  <math>\sum \alpha\beta = \alpha\alpha^* + \alpha\beta + \alpha\beta^* + \beta\beta^* + \beta\alpha^* + \beta^*\alpha^*</math>  <math>\sum \alpha\beta\gamma = \alpha\alpha^*\beta + \alpha\alpha^*\beta^* + \alpha\beta\beta^* + \alpha^*\beta\beta^*</math></p> <p><math>\sum \alpha\beta = 2, \sum \alpha\beta\gamma = -8</math></p> <p><math>A = -2, B = 2, C = 8, D = 16</math></p> <p><b>OR</b></p> <p>Attempt to substitute in one root          Attempt to substitute in a second root</p> <p>Equating real and imaginary parts to 0          Attempt to solve simultaneous equations</p> <p><math>A = -2, B = 2, C = 8, D = 16</math></p>	<p>M1 B2 A1 M1 A2 [7] B1 B1 M1 M1 A1 A2 [7] M1 M1 A1 M1 M1 A2 [7]</p>	<p>Attempt to use factor theorem          Correct factors, minus 1 each error          B1 if only errors are sign errors          One correct quadratic with real coefficients (may be implied)          Expanding          Minus 1 each error, A1 if only errors are sign errors          Both correct          Minus 1 each error, A1 if only errors are sign errors          Both correct          Minus 1 each error, A1 if only errors are sign errors</p>

Qu	Answer	Mark	Comment
<b>Section B (continued)</b>			
10(i)	$\sum_{r=1}^n r^2(r+1) = \sum_{r=1}^n r^3 + \sum_{r=1}^n r^2$ $= \frac{1}{4}n^2(n+1)^2 + \frac{1}{6}n(n+1)(2n+1)$ $= \frac{1}{12}n(n+1)[3n(n+1) + 2(2n+1)]$ $= \frac{1}{12}n(n+1)(3n^2 + 7n + 2)$ $= \frac{1}{12}n(n+1)(n+2)(3n+1)$	M1 B1 M1 A1 E1  <b>[5]</b>	Separation of sums (may be implied) One mark for both parts Attempt to factorise (at least two linear algebraic factors) Correct Complete, convincing argument
10(ii)	$\sum_{r=1}^n r^2(r+1) = \frac{1}{12}n(n+1)(n+2)(3n+1)$ <p><math>n = 1</math>, LHS = RHS = 2</p> <p>Assume true for <math>n = k</math></p> $\sum_{r=1}^k r^2(r+1) = \frac{1}{12}k(k+1)(k+2)(3k+1)$ $\sum_{r=1}^{k+1} r^2(r+1)$ $= \frac{1}{12}k(k+1)(k+2)(3k+1) + (k+1)^2(k+2)$ $= \frac{1}{12}(k+1)(k+2)[k(3k+1) + 12(k+1)]$ $= \frac{1}{12}(k+1)(k+2)(3k^2 + 13k + 12)$ $= \frac{1}{12}(k+1)(k+2)(k+3)(3k+4)$ $= \frac{1}{12}(k+1)((k+1)+1)((k+1)+2)(3(k+1)+1)$ <p>But this is the given result with <math>k + 1</math> replacing <math>k</math>. Therefore if it is true for <math>k</math> it is true for <math>k + 1</math>. Since it is true for <math>k = 1</math>, it is true for <math>k = 1, 2, 3</math> and so true for all positive integers.</p>	B1 E1  B1 M1 A1 A1  E1 E1  <b>[8]</b>	2 must be seen Assuming true for $k$  ( $k + 1$ )th term Attempt to factorise Correct Complete convincing argument Dependent on previous A1 and previous E1 Dependent on first B1 and previous E1
			<b>Section B Total: 36</b>
			<b>Total: 72</b>


## 4756 (FP2) Further Methods for Advanced Mathematics

1(a)(i)	$x = r \cos \theta, \quad y = r \sin \theta$ $(r^2 \cos^2 \theta + r^2 \sin^2 \theta)^2 = 3(r \cos \theta)(r \sin \theta)^2$ $r^4 = 3r^3 \cos \theta \sin^2 \theta$ $r = 3 \cos \theta \sin^2 \theta$	M1 A1  A1 ag <b>3</b>	(M0 for $x = \cos \theta, y = \sin \theta$ )
(ii)		B1  B1  B1 <b>3</b>	Loop in 1st quadrant  Loop in 4th quadrant  Fully correct curve <i>Curve may be drawn using continuous or broken lines in any combination</i>
(b)	$\int_0^1 \frac{1}{\sqrt{4-3x^2}} dx = \left[ \frac{1}{\sqrt{3}} \arcsin \frac{\sqrt{3}x}{2} \right]_0^1$ $= \frac{1}{\sqrt{3}} \arcsin \frac{\sqrt{3}}{2}$ $= \frac{\pi}{3\sqrt{3}}$ <p>OR</p> <p>Put <math>\sqrt{3}x = 2 \sin \theta</math></p> $\int_0^1 \frac{1}{\sqrt{4-3x^2}} dx = \int_0^{\frac{\pi}{3}} \frac{1}{\sqrt{3}} d\theta$ $= \frac{\pi}{3\sqrt{3}}$	M1 A1A1  M1 A1 <b>5</b>	For arcsin For $\frac{1}{\sqrt{3}}$ and $\frac{\sqrt{3}x}{2}$  Exact numerical value <i>Dependent on first M1</i> (M1A0 for $60/\sqrt{3}$ )
(c)(i)	$\ln(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \frac{1}{5}x^5 - \dots$ $\ln(1-x) = -x - \frac{1}{2}x^2 - \frac{1}{3}x^3 - \frac{1}{4}x^4 - \frac{1}{5}x^5 - \dots$	B1  B1 <b>2</b>	<i>Accept unsimplified forms</i>
(ii)	$\ln\left(\frac{1+x}{1-x}\right) = \ln(1+x) - \ln(1-x)$ $= 2x + \frac{2}{3}x^3 + \frac{2}{5}x^5 + \dots$	M1  A1 <b>2</b>	Obtained from two correct series <i>Terms need not be added</i> If M0, then B1 for $2x + \frac{2}{3}x^3 + \frac{2}{5}x^5$


(iii)	$\sum_{r=0}^{\infty} \frac{1}{(2r+1)4^r} = 1 + \frac{1}{3 \times 4} + \frac{1}{5 \times 4^2} + \dots$ $= 2 \times \frac{1}{2} + \frac{2}{3} \times \left(\frac{1}{2}\right)^3 + \frac{2}{5} \times \left(\frac{1}{2}\right)^5 + \dots$ $= \ln\left(\frac{1+\frac{1}{2}}{1-\frac{1}{2}}\right) = \ln 3$	B1 B1 B1 ag <b>3</b>	<i>Terms need not be added</i> For $x = \frac{1}{2}$ seen or implied Satisfactory completion
2 (i)	$ z  = 8, \quad \arg z = \frac{1}{4}\pi$  $ z^*  = 8, \quad \arg z^* = -\frac{1}{4}\pi$ $ z w  = 8 \times 8 = 64$ $\arg(z w) = \frac{1}{4}\pi + \frac{7}{12}\pi = \frac{5}{6}\pi$  $\left \frac{z}{w}\right  = \frac{8}{8} = 1$  $\arg\left(\frac{z}{w}\right) = \frac{1}{4}\pi - \frac{7}{12}\pi = -\frac{1}{3}\pi$	B1B1  B1 ft B1 ft B1 ft  B1 ft  B1 ft <b>7</b>	<i>Must be given separately</i> <i>Remainder may be given in exponential or <math>r c j s \theta</math> form</i> (B0 for $\frac{7}{4}\pi$ )  (B0 if left as $8/8$ )
(ii)	$\frac{z}{w} = \cos\left(-\frac{1}{3}\pi\right) + j \sin\left(-\frac{1}{3}\pi\right)$ $= \frac{1}{2} - \frac{\sqrt{3}}{2}j$ $a = \frac{1}{2}, \quad b = -\frac{1}{2}\sqrt{3}$	M1 A1 <b>2</b>	If M0, then B1B1 for $\frac{1}{2}$ and $-\frac{\sqrt{3}}{2}$
(iii)	$r = \sqrt[3]{8} = 2$ $\theta = \frac{1}{12}\pi$ $\theta = \frac{\pi}{12} + \frac{2k\pi}{3}$ $\theta = -\frac{7}{12}\pi, \quad \frac{3}{4}\pi$	B1 ft B1  M1 A1 <b>4</b>	Accept $\sqrt[3]{8}$  Implied by one further correct (ft) value <i>Ignore values outside the required range</i>
(iv)	$w^* = 8e^{-\frac{7}{12}\pi j}, \quad \text{so } 2e^{-\frac{7}{12}\pi j} = \frac{1}{4}w^*$ $k_1 = \frac{1}{4}$  $z^* = 8e^{-\frac{1}{4}\pi j} = -8e^{\frac{3}{4}\pi j}$  So $2e^{\frac{3}{4}\pi j} = -\frac{1}{4}z^*$ $k_2 = -\frac{1}{4}$  $jw = 8e^{(\frac{1}{2}\pi + \frac{7}{12}\pi)j} = 8e^{\frac{13}{12}\pi j}$ $= -8e^{\frac{1}{12}\pi j}, \quad \text{so } 2e^{\frac{1}{12}\pi j} = -\frac{1}{4}jw$ $k_3 = -\frac{1}{4}$	B1 ft   M1  A1 ft  M1  A1 ft <b>5</b>	Matching $w^*$ to a cube root with argument $-\frac{7}{12}\pi$ and $k_1 = \frac{1}{4}$ or ft is $\frac{r}{8}$  Matching $z^*$ to a cube root with argument $\frac{3}{4}\pi$ <i>May be implied</i> ft is $-\frac{r}{ z^* }$  Matching $jw$ to a cube root with argument $\frac{1}{12}\pi$ <i>May be implied</i> OR M1 for $\arg(jw) = \frac{1}{2}\pi + \arg w$ ( <i>implied by <math>\frac{13}{12}\pi</math> or <math>-\frac{11}{12}\pi</math></i> ) ft is $-\frac{r}{8}$

<b>3 (i)</b>	$\mathbf{Q}^{-1} = \frac{1}{k-3} \begin{pmatrix} -1 & k+2 & -1 \\ 1 & 4-3k & k-2 \\ 1 & -5 & 1 \end{pmatrix}$ <p>When <math>k = 4</math>, <math>\mathbf{Q}^{-1} = \begin{pmatrix} -1 &amp; 6 &amp; -1 \\ 1 &amp; -8 &amp; 2 \\ 1 &amp; -5 &amp; 1 \end{pmatrix}</math></p>	M1 A1 M1 A1 M1 A1	Evaluation of determinant <i>(must involve k)</i> For $(k-3)$ Finding at least four cofactors <i>(including one involving k)</i> Six signed cofactors correct <i>(including one involving k)</i> Transposing and dividing by det <i>Dependent on previous M1M1</i> $\mathbf{Q}^{-1}$ correct (in terms of $k$ ) and <b>6</b> result for $k = 4$ stated After 0, SC1 for $\mathbf{Q}^{-1}$ when $k = 4$ obtained correctly with some working
<b>(ii)</b>	$\mathbf{P} = \begin{pmatrix} 2 & -1 & 4 \\ 1 & 0 & 1 \\ 3 & 1 & 2 \end{pmatrix}, \quad \mathbf{D} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 3 \end{pmatrix}$ $\mathbf{M} = \mathbf{PDP}^{-1}$ $= \begin{pmatrix} 2 & -1 & 4 \\ 1 & 0 & 1 \\ 3 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} -1 & 6 & -1 \\ 1 & -8 & 2 \\ 1 & -5 & 1 \end{pmatrix}$ $= \begin{pmatrix} 2 & 1 & 12 \\ 1 & 0 & 3 \\ 3 & -1 & 6 \end{pmatrix} \begin{pmatrix} -1 & 6 & -1 \\ 1 & -8 & 2 \\ 1 & -5 & 1 \end{pmatrix}$ $= \begin{pmatrix} 11 & -56 & 12 \\ 2 & -9 & 2 \\ 2 & -4 & 1 \end{pmatrix}$	B1B1 B2 M1 A2	For B2, order must be consistent Give B1 for $\mathbf{M} = \mathbf{P}^{-1} \mathbf{D} \mathbf{P}$ $\text{or } \begin{pmatrix} 2 & -1 & 4 \\ 1 & 0 & 1 \\ 3 & 1 & 2 \end{pmatrix} \begin{pmatrix} -1 & 6 & -1 \\ -1 & 8 & -2 \\ 3 & -15 & 3 \end{pmatrix}$ Good attempt at multiplying two matrices (no more than 3 errors), leaving third matrix in correct position <b>7</b> Give A1 for five elements correct Correct $\mathbf{M}$ implies B2M1A2 5-8 elements correct implies B2M1A1
<b>(iii)</b>	Characteristic equation is $(\lambda-1)(\lambda+1)(\lambda-3) = 0$ $\lambda^3 - 3\lambda^2 - \lambda + 3 = 0$ $\mathbf{M}^3 = 3\mathbf{M}^2 + \mathbf{M} - 3\mathbf{I}$ $\mathbf{M}^4 = 3\mathbf{M}^3 + \mathbf{M}^2 - 3\mathbf{M}$ $= 3(3\mathbf{M}^2 + \mathbf{M} - 3\mathbf{I}) + \mathbf{M}^2 - 3\mathbf{M}$ $= 10\mathbf{M}^2 - 9\mathbf{I}$ $a = 10, b = 0, c = -9$	B1 M1 A1 M1 A1	In any correct form <i>(Condone omission of =0)</i> $\mathbf{M}$ satisfies the characteristic equation Correct expanded form <i>(Condone omission of I)</i> <b>5</b>

<b>4 (i)</b>	$\cosh^2 x = \left[\frac{1}{2}(e^x + e^{-x})\right]^2 = \frac{1}{4}(e^{2x} + 2 + e^{-2x})$ $\sinh^2 x = \left[\frac{1}{2}(e^x - e^{-x})\right]^2 = \frac{1}{4}(e^{2x} - 2 + e^{-2x})$ $\cosh^2 x - \sinh^2 x = \frac{1}{4}(2 + 2) = 1$ <p>OR</p> $\cosh x + \sinh x = \frac{1}{2}(e^x + e^{-x}) + \frac{1}{2}(e^x - e^{-x}) = e^x \quad \text{B1}$ $\cosh x - \sinh x = \frac{1}{2}(e^x + e^{-x}) - \frac{1}{2}(e^x - e^{-x}) = e^{-x} \quad \text{B1}$ $\cosh^2 x - \sinh^2 x = e^x \times e^{-x} = 1 \quad \text{B1}$	B1 B1 B1 ag <b>3</b>	For completion
<b>(ii)</b>	$4(1 + \sinh^2 x) + 9 \sinh x = 13$ $4 \sinh^2 x + 9 \sinh x - 9 = 0$ $\sinh x = \frac{3}{4}, -3$ $x = \ln 2, \ln(-3 + \sqrt{10})$ <p>OR</p> $2e^{4x} + 9e^{3x} - 22e^{2x} - 9e^x + 2 = 0$ $(2e^{2x} - 3e^x - 2)(e^{2x} + 6e^x - 1) = 0 \quad \text{M1}$ $e^x = 2, -3 + \sqrt{10} \quad \text{M1}$ $x = \ln 2, \ln(-3 + \sqrt{10}) \quad \text{A1A1}$	M1 M1 A1A1 A1A1 ft <b>6</b>	(M0 for $1 - \sinh^2 x$ ) Obtaining a value for $\sinh x$ Exact logarithmic form <i>Dep on M1M1</i> Max A1 if any extra values given Quadratic and / or linear factors Obtaining a value for $e^x$ Ignore extra values <i>Dependent on M1M1</i> Max A1 if any extra values given <i>Just <math>x = \ln 2</math> earns M0M1A1A0A0A0</i>
<b>(iii)</b>	$\frac{dy}{dx} = 8 \cosh x \sinh x + 9 \cosh x$ $= \cosh x(8 \sinh x + 9)$ $= 0 \text{ only when } \sinh x = -\frac{9}{8}$ $\cosh^2 x = 1 + \left(-\frac{9}{8}\right)^2 = \frac{145}{64}$ $y = 4 \times \frac{145}{64} + 9 \times \left(-\frac{9}{8}\right) = -\frac{17}{16}$	B1 B1 M1 A1 <b>4</b>	Any correct form <i>or <math>y = (2 \sinh x + \frac{9}{4})^2 + \dots (-\frac{17}{16})</math></i> Correctly showing there is only one solution Exact evaluation of $y$ or $\cosh^2 x$ or $\cosh 2x$ Give B2 (replacing M1A1) for -1.06 or better
<b>(iv)</b>	$\int_0^{\ln 2} (2 + 2 \cosh 2x + 9 \sinh x) dx$ $= \left[ 2x + \sinh 2x + 9 \cosh x \right]_0^{\ln 2}$ $= \left\{ 2 \ln 2 + \frac{1}{2} \left( 4 - \frac{1}{4} \right) + \frac{9}{2} \left( 2 + \frac{1}{2} \right) \right\} - 9$ $= 2 \ln 2 + \frac{33}{8}$	M1 A2 M1 A1 ag <b>5</b>	Expressing in integrable form Give A1 for two terms correct $\sinh(2 \ln 2) = \frac{1}{2} \left( 4 - \frac{1}{4} \right)$ <i>Must see both terms for M1</i> <i>Must also see <math>\cosh(\ln 2) = \frac{1}{2} \left( 2 + \frac{1}{2} \right)</math> for A1</i>

	<p>OR <math>\int_0^{\ln 2} (e^{2x} + 2 + e^{-2x} + \frac{9}{2}(e^x - e^{-x})) dx</math> M1</p> <p><math>= \left[ \frac{1}{2}e^{2x} + 2x - \frac{1}{2}e^{-2x} + \frac{9}{2}e^x + \frac{9}{2}e^{-x} \right]_0^{\ln 2}</math> A2</p> <p><math>= \left( 2 + 2\ln 2 - \frac{1}{8} + 9 + \frac{9}{4} \right) - \left( \frac{1}{2} - \frac{1}{2} + \frac{9}{2} + \frac{9}{2} \right)</math> M1</p> <p><math>= 2\ln 2 + \frac{33}{8}</math> A1 ag</p>		<p>Expanded exponential form (M0 if the 2 is omitted)</p> <p>Give A1 for three terms correct</p> <p><math>e^{2\ln 2} = 4</math> and <math>e^{-2\ln 2} = \frac{1}{4}</math> both seen</p> <p><b>Must also see</b>  <math>e^{\ln 2} = 2</math> and <math>e^{-\ln 2} = \frac{1}{2}</math>  <b>for A1</b></p>
5 (i)	<p><math>\lambda = 0.5</math>                      <math>\lambda = 3</math>                      <math>\lambda = 5</math></p> 	B1B1B1 3	
(ii)	Ellipse	B1 1	
(iii)	<p><math>y = \sqrt{2} \cos(\theta - \frac{1}{4}\pi)</math></p> <p>Maximum <math>y = \sqrt{2}</math> when <math>\theta = \frac{1}{4}\pi</math></p>	M1 A1 ag 2	or $\sqrt{2} \sin(\theta + \frac{1}{4}\pi)$
	<p>OR <math>\frac{dy}{d\theta} = -\sin \theta + \cos \theta = 0</math> when <math>\theta = \frac{1}{4}\pi</math> M1</p> <p><math>y = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \sqrt{2}</math> A1</p>		
(iv)	<p><math>x^2 + y^2 = \lambda^2 \cos^2 \theta - 2 \cos \theta \sin \theta + \frac{1}{\lambda^2} \sin^2 \theta</math></p> <p><math>+ \cos^2 \theta + 2 \cos \theta \sin \theta + \sin^2 \theta</math></p> <p><math>= (\lambda^2 + 1)(1 - \sin^2 \theta) + (\frac{1}{\lambda^2} + 1) \sin^2 \theta</math></p> <p><math>= 1 + \lambda^2 + (\frac{1}{\lambda^2} - \lambda^2) \sin^2 \theta</math></p> <p>When <math>\sin^2 \theta = 0</math>, <math>x^2 + y^2 = 1 + \lambda^2</math></p> <p>When <math>\sin^2 \theta = 1</math>, <math>x^2 + y^2 = 1 + \frac{1}{\lambda^2}</math></p> <p>Since <math>0 \leq \sin^2 \theta \leq 1</math>, distance from O,</p> <p><math>\sqrt{x^2 + y^2}</math>, is between <math>\sqrt{1 + \frac{1}{\lambda^2}}</math> and <math>\sqrt{1 + \lambda^2}</math></p>	M1 M1 A1 ag M1 M1 A1 ag 6	Using $\cos^2 \theta = 1 - \sin^2 \theta$
(v)	<p>When <math>\lambda = 1</math>, <math>x^2 + y^2 = 2</math></p> <p>Curve is a circle (centre O) with radius <math>\sqrt{2}</math></p>	M1 A1 2	



(vi)		B4 4	<p>A, E at maximum distance from O        C, G at minimum distance from O        B, F are stationary points        D, H are on the x-axis</p> <p>Give <math>\frac{1}{2}</math> mark for each point, then round down</p> <p>Special properties must be clear from diagram, or stated</p> <p><i>Max 3 if curve is not the correct shape</i></p>
------	---	---------	---

## 4757 (FP3) Further Applications of Advanced Mathematics

1 (i)	$\overline{AB} \times \overline{AC} = \begin{pmatrix} 6 \\ 8 \\ 5 \end{pmatrix} \times \begin{pmatrix} 10 \\ -5 \\ 1 \end{pmatrix} = \begin{pmatrix} 33 \\ 44 \\ -110 \end{pmatrix}$ <p>ABC is <math>3x + 4y - 10z = -9 + 20 - 20</math>  <math>3x + 4y - 10z + 9 = 0</math></p>	B2  M1 A1  <b>4</b>	<p><i>Ignore subsequent working</i>          Give B1 for one element correct          SC1 for minus the correct vector</p> <p>For <math>3x + 4y - 10z</math>          Accept <math>33x + 44y - 110z = -99</math> etc</p>
(ii)	<p>Distance is <math>\frac{3 \times 5 + 4 \times 4 - 10 \times 8 + 9}{\sqrt{3^2 + 4^2 + 10^2}}</math></p> $= (-) \frac{40}{\sqrt{125}} \quad (= \frac{8}{\sqrt{5}})$	M1 A1 ft  A1  <b>3</b>	<p>Using distance formula (or other complete method)</p> <p><i>Condone negative answer</i>          Accept a.r.t. 3.58</p>
(iii)	$\overline{AB} \times \overline{CD} = \begin{pmatrix} 6 \\ 8 \\ 5 \end{pmatrix} \times \begin{pmatrix} -2 \\ 4 \\ 5 \end{pmatrix} = \begin{pmatrix} 20 \\ -40 \\ 40 \end{pmatrix} \quad [ = 20 \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix} ]$ $\text{Distance is } \overline{AC} \cdot \hat{n} = \frac{\begin{pmatrix} 10 \\ -5 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}}{\sqrt{1^2 + 2^2 + 2^2}}$ $= \frac{22}{3}$	M1  A1  M1  A1  <b>4</b>	<p>Evaluating <math>\overline{AB} \times \overline{CD}</math> or method for finding end-points of common perp PQ</p> <p>or P <math>(\frac{3}{2}, 11, \frac{23}{4})</math> &amp;          Q <math>(\frac{71}{18}, \frac{55}{9}, \frac{383}{36})</math>          or <math>\overline{PQ} = (\frac{22}{9}, -\frac{44}{9}, \frac{44}{9})</math></p>
(iv)	<p>Volume is <math>\frac{1}{6}(\overline{AB} \times \overline{AC}) \cdot \overline{AD}</math></p> $= \frac{1}{6} \begin{pmatrix} 33 \\ 44 \\ -110 \end{pmatrix} \cdot \begin{pmatrix} 8 \\ -1 \\ 6 \end{pmatrix}$ $= (-) \frac{220}{3}$	M1 A1  M1  A1  <b>4</b>	<p>Scalar triple product</p> <p>Accept a.r.t. 73.3</p>
(v)	<p>E is <math>(-3 + 10\lambda, 5 - 5\lambda, 2 + \lambda)</math>  <math>3(-3 + 10\lambda) - 2(2 + \lambda) + 5 = 0</math>  <math>\lambda = \frac{2}{7}</math></p> <p>F is <math>(-3 + 8\mu, 5 - \mu, 2 + 6\mu)</math>  <math>3(-3 + 8\mu) - 2(2 + 6\mu) + 5 = 0</math>  <math>\mu = \frac{2}{3}</math></p> <p>Since <math>0 &lt; \lambda &lt; 1</math>, E is between A and C          Since <math>0 &lt; \mu &lt; 1</math>, F is between A and D</p>	M1  A1  M1  A1  B1  <b>5</b>	

<b>(vi)</b>	$V_{ABEF} = \frac{1}{6}(\overline{AB} \times \overline{AE}) \cdot \overline{AF}$ $= \frac{1}{6} \lambda \mu (\overline{AB} \times \overline{AC}) \cdot \overline{AD}$ $= \lambda \mu V_{ABCD}$ $= \frac{4}{21} V_{ABCD}$ <p>Ratio of volumes is <math>\frac{4}{21} : \frac{17}{21}</math></p> $= 4 : 17$	M1 A1 M1 A1 ag	$(13\frac{61}{63})$ ft if numerical  Finding ratio of volumes of two parts 4 SC1 for 4 : 17 deduced from $\frac{4}{21}$ without working
<b>2 (i)</b>	$\frac{\partial g}{\partial x} = 6z - 2(x + 2y + 3z) = -2x - 4y$ $\frac{\partial g}{\partial y} = -4(x + 2y + 3z)$ $\frac{\partial g}{\partial z} = 6x - 6(x + 2y + 3z) = -12y - 18z$	M1 A1 A1 A1	Partial differentiation Any correct form, ISW  4
<b>(ii)</b>	<p>At P, <math>\frac{\partial g}{\partial x} = 16</math>, <math>\frac{\partial g}{\partial y} = -4</math>, <math>\frac{\partial g}{\partial z} = 36</math></p> <p>Normal line is <math>\mathbf{r} = \begin{pmatrix} 7 \\ -7.5 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ -1 \\ 9 \end{pmatrix}</math></p>	M1 A1 A1 ft	Evaluating partial derivatives at P All correct  3 Condone omission of 'r = '
<b>(iii)</b>	$\delta g \approx 16 \delta x - 4 \delta y + 36 \delta z$ <p>If <math>\overline{PQ} = \lambda \begin{pmatrix} 4 \\ -1 \\ 9 \end{pmatrix}</math>,</p> $\delta g \approx 16(4\lambda) - 4(-\lambda) + 36(9\lambda) \quad (= 392\lambda)$ <p><math>h = \delta g</math>, so <math>h \approx 392\lambda</math></p> $\overline{PQ} \approx \frac{h}{392} \begin{pmatrix} 4 \\ -1 \\ 9 \end{pmatrix}, \text{ so } \mathbf{n} = \frac{1}{392} \begin{pmatrix} 4 \\ -1 \\ 9 \end{pmatrix}$	M1 M1 A1 ft M1 A1	Alternative: M3 for substituting $x = 7 + 4\lambda$ , ... into $g = 125 + h$ and neglecting $\lambda^2$ A1 ft for linear equation in $\lambda$ and $h$ A1 for n correct  5
<b>(iv)</b>	<p>Require <math>\frac{\partial g}{\partial x} = \frac{\partial g}{\partial y} = 0</math></p> <p><math>-2x - 4y = 0</math> and <math>x + 2y + 3z = 0</math></p> <p><math>x + 2y = 0</math> and <math>z = 0</math></p> <p><math>g(x, y, z) = 0 - 0^2 = 0 \neq 125</math></p> <p>Hence there is no such point on S</p>	M1 M1 M1 A1	Useful manipulation using both eqns  Showing there is no such point on S 4 Fully correct proof
<b>(v)</b>	<p>Require <math>\frac{\partial g}{\partial z} = 0</math></p> <p>and <math>\frac{\partial g}{\partial y} = 5 \frac{\partial g}{\partial x}</math></p> <p><math>-4x - 8y - 12z = 5(-2x - 4y)</math></p>	M1 M1 M1	Implied by $\frac{\partial g}{\partial x} = \lambda$ , $\frac{\partial g}{\partial y} = 5\lambda$  This M1 can be awarded for $-2x - 4y = 1$ and $-4x - 8y - 12z = 5$

	$y = -\frac{3}{2}z$ and $x = 5z$  $6(5z)z - (5z)^2 = 125$ $z = \pm 5$  Points are $(25, -7.5, 5)$ and $(-25, 7.5, -5)$	A1  M1 M1 A1 A1 ft  <b>8</b>	$or\ z = -\frac{2}{3}y$ and $x = -\frac{10}{3}y$ $or\ y = -\frac{3}{10}x$ and $z = \frac{1}{5}x$ $or\ x = -\frac{5}{4}\lambda$ , $y = \frac{3}{8}\lambda$ , $z = -\frac{1}{4}\lambda$ $or\ x : y : z = 10 : -3 : 2$ Substituting into $g(x, y, z) = 125$ Obtaining one value of $x, y, z$ or $\lambda$ <i>Dependent on previous M1</i> <i>ft is minus the other point,</i> <i>provided all M marks have been earned</i>
<b>3 (i)</b>	$\dot{x}^2 + \dot{y}^2 = (24t^2)^2 + (18t - 8t^3)^2$ $= 576t^4 + 324t^2 - 288t^4 + 64t^6$ $= 324t^2 + 288t^4 + 64t^6$ $= (18t + 8t^3)^2$ Arc length is $\int_0^2 (18t + 8t^3) dt$ $= [9t^2 + 2t^4]_0^2$ $= 68$	B1  M1 A1 ag  M1  A1 A1  <b>6</b>	<i>Note</i> $\int_0^2 (18t + 8t^3) dt = [9t^2 + 2t^4]_0^2 = 68$ <i>earns M1A0A0</i>
<b>(ii)</b>	Curved surface area is $\int 2\pi y ds$ $= \int_0^2 2\pi(9t^2 - 2t^4)(18t + 8t^3) dt$ $= \int_0^2 \pi(324t^3 + 72t^5 - 32t^7) dt$ $= \pi [81t^4 + 12t^6 - 4t^8]_0^2$ $= 1040\pi \quad (\approx 3267)$	M1  M1 A1  M1  M1  A1  <b>6</b>	Using $ds = (18t + 8t^3) dt$ Correct integral expression including limits ( <i>may be implied by later work</i> )
<b>(iii)</b>	$\kappa = \frac{\dot{x}\ddot{y} - \ddot{x}y}{(\dot{x}^2 + \dot{y}^2)^{\frac{3}{2}}} = \frac{(24t^2)(18 - 24t^2) - (48t)(18t - 8t^3)}{(18t + 8t^3)^3}$ $= \frac{48t^2(9 - 12t^2 - 18 + 8t^2)}{8t^3(9 + 4t^2)^3} = \frac{-48t^2(9 + 4t^2)}{8t^3(9 + 4t^2)^3}$ $= \frac{-6}{t(4t^2 + 9)^2}$	M1 A1A1  M1  A1 ag  <b>5</b>	Using formula for $\kappa$ (or $\rho$ ) For numerator and denominator  Simplifying the numerator

(iv)	<p>When <math>t=1</math>, <math>x=8</math>, <math>y=7</math>, <math>\kappa=-\frac{6}{169}</math></p> $\rho = (-) \frac{169}{6}$ $\frac{dy}{dx} = \frac{\dot{y}}{\dot{x}} = \frac{18t-8t^3}{24t^2} = \frac{10}{24}$ $\hat{\mathbf{n}} = \begin{pmatrix} \frac{5}{13} \\ -\frac{12}{13} \end{pmatrix}$ $\mathbf{c} = \begin{pmatrix} 8 \\ 7 \end{pmatrix} + \frac{169}{6} \begin{pmatrix} \frac{5}{13} \\ -\frac{12}{13} \end{pmatrix}$ <p>Centre of curvature is <math>(18\frac{5}{6}, -19)</math></p>	<p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1A1</p>	<p>Finding gradient (or tangent vector)</p> <p>Finding direction of the normal</p> <p>Correct unit normal (either direction)</p> <p><b>7</b></p>
4 (i)	<p><b>Commutative:</b> <math>x * y = y * x</math> (for all <math>x, y</math>)</p> <p><b>Associative:</b> <math>(x * y) * z = x * (y * z)</math></p> <p>(for all <math>x, y, z</math>)</p>	<p>B1</p> <p>B2</p>	<p>Accept e.g. 'Order does not matter'</p> <p><b>3</b> Give B1 for a partial explanation, e.g. 'Position of brackets does not matter'</p>
(ii)	$2(x + \frac{1}{2})(y + \frac{1}{2}) - \frac{1}{2} = 2xy + x + y + \frac{1}{2} - \frac{1}{2}$ $= 2xy + x + y = x * y$	<p>B1 ag</p>	<p><i>Intermediate step required</i></p> <p><b>1</b></p>
(iii)(A)	<p>If <math>x, y \in S</math> then <math>x &gt; -\frac{1}{2}</math> and <math>y &gt; -\frac{1}{2}</math></p> <p><math>x + \frac{1}{2} &gt; 0</math> and <math>y + \frac{1}{2} &gt; 0</math>, so <math>2(x + \frac{1}{2})(y + \frac{1}{2}) &gt; 0</math></p> <p><math>2(x + \frac{1}{2})(y + \frac{1}{2}) - \frac{1}{2} &gt; -\frac{1}{2}</math>, so <math>x * y \in S</math></p>	<p>M1</p> <p>A1</p> <p>A1</p>	<p><b>3</b></p>
(B)	<p><b>0 is the identity</b> since <math>0 * x = 0 + x + 0 = x</math></p> <p>If <math>x \in S</math> and <math>x * y = 0</math> then</p> $2xy + x + y = 0$ $y = \frac{-x}{2x+1}$ $y + \frac{1}{2} = \frac{1}{2(2x+1)} > 0 \quad (\text{since } x > -\frac{1}{2})$ <p>so <math>y \in S</math></p> <p>S is closed and associative; there is an identity; and every element of S has an inverse in S</p>	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>or <math>2(x + \frac{1}{2})(y + \frac{1}{2}) - \frac{1}{2} = 0</math></p> <p>or <math>y + \frac{1}{2} = \frac{1}{4(x + \frac{1}{2})}</math></p> <p><i>Dependent on M1A1M1</i></p> <p><b>6</b></p>
(iv)	<p>If <math>x * x = 0</math>, <math>2x^2 + x + x = 0</math></p> $x = 0 \text{ or } -1$ <p>0 is the identity (and has order 1)</p> <p>-1 is not in S</p>	<p>M1</p> <p>A1</p> <p>A1</p>	<p><b>3</b></p>

<b>(v)</b>	$4 * 6 = 48 + 4 + 6 = 58$ $= 56 + 2 = 7 \times 8 + 2$ So $4 \circ 6 = 2$							B1  B1 ag  <b>2</b>		
<b>(vi)</b>	Element	0	1	2	4	5	6		B3  <b>3</b>	Give B2 for 4 correct B1 for 2 correct
<b>(vii)</b>	$\{0\}, G$ $\{0, 6\}$ $\{0, 4, 5\}$							B1 B1 B1  <b>3</b>	<i>Condone omission of G</i>  If more than 2 non-trivial subgroups are given, deduct 1 mark (from final B1B1) for each non-trivial subgroup in excess of 2	

## Pre-multiplication by transition matrix



5 (i)	$P = \begin{pmatrix} 0.1 & 0.7 & 0.1 \\ 0.4 & 0.2 & 0.6 \\ 0.5 & 0.1 & 0.3 \end{pmatrix}$	B2 <b>2</b>	Give B1 for two columns correct
(ii)	$P^6 \begin{pmatrix} \frac{1}{3} \\ \frac{1}{3} \\ \frac{1}{3} \end{pmatrix} = \begin{pmatrix} 0.328864 \\ 0.381536 \\ 0.2896 \end{pmatrix}$ <p><math>P(B \text{ used on 7th day}) = 0.3815</math></p>	M1 M1 M1 A1 <b>4</b>	Using $P^6$ (or $P^7$ ) For matrix of initial probabilities For evaluating matrix product <i>Accept 0.381 to 0.382</i>
(iii)	$0.328864 \times 0.1 + 0.381536 \times 0.2 + 0.2896 \times 0.3$ $= 0.1961$	M1 M1 A1 <b>3</b>	Using diagonal elements from $P$ Correct method <i>Accept a.r.t. 0.196</i>
(iv)	$P^3 = \begin{pmatrix} 0.352 & 0.328 & 0.304 \\ 0.364 & 0.404 & 0.372 \\ 0.284 & 0.268 & 0.324 \end{pmatrix}$ $0.328864 \times 0.352 + 0.381536 \times 0.404 + 0.2896 \times 0.324$ $= 0.3637$	M1 M1 M1 A1 <b>4</b>	For evaluating $P^3$ Using diagonal elements from $P^3$ Correct method <i>Accept a.r.t. 0.364</i>
(v)	$Q = \begin{pmatrix} 0.3289 & 0.3289 & 0.3289 \\ 0.3816 & 0.3816 & 0.3816 \\ 0.2895 & 0.2895 & 0.2895 \end{pmatrix}$ <p>0.3289, 0.3816, 0.2895 are the long-run probabilities for the routes A, B, C</p>	B1 B1 B1 B1 <b>4</b>	<i>Deduct 1 if not given as a (3×3) matrix</i> <i>Deduct 1 if not 4 dp</i>  <i>Accept 'equilibrium probabilities'</i>
(vi)	$\begin{pmatrix} 0.1 & 0.7 & a \\ 0.4 & 0.2 & b \\ 0.5 & 0.1 & c \end{pmatrix} \begin{pmatrix} 0.4 \\ 0.2 \\ 0.4 \end{pmatrix} = \begin{pmatrix} 0.4 \\ 0.2 \\ 0.4 \end{pmatrix}$ <p><math>0.04 + 0.14 + 0.4a = 0.4</math>, so <math>a = 0.55</math>  <math>0.16 + 0.04 + 0.4b = 0.2</math>, so <math>b = 0</math>  <math>0.2 + 0.02 + 0.4c = 0.4</math>, so <math>c = 0.45</math></p> <p>After C, routes A, B, C will be used with probabilities 0.55, 0, 0.45</p>	M1 M1 A2 <b>4</b>	Obtaining a value for a, b or c  Give A1 for one correct
(vii)	$0.4 \times 0.1 + 0.2 \times 0.2 + 0.4 \times 0.45$ $= 0.26$	M1 M1 A1 <b>3</b>	Using long-run probs 0.4, 0.2, 0.4 Using diag elements from new matrix


## Post-multiplication by transition matrix


5 (i)	$P = \begin{pmatrix} 0.1 & 0.4 & 0.5 \\ 0.7 & 0.2 & 0.1 \\ 0.1 & 0.6 & 0.3 \end{pmatrix}$	B2 <b>2</b>	Give B1 for two rows correct
(ii)	$\left(\frac{1}{3} \quad \frac{1}{3} \quad \frac{1}{3}\right) P^6 = (0.328864 \quad 0.381536 \quad 0.2896)$ $P(B \text{ used on 7th day}) = 0.3815$	M1 M1 M1 A1 <b>4</b>	Using $P^6$ (or $P^7$ ) For matrix of initial probabilities For evaluating matrix product <i>Accept 0.381 to 0.382</i>
(iii)	$0.328864 \times 0.1 + 0.381536 \times 0.2 + 0.2896 \times 0.3$ $= 0.1961$	M1 M1 A1 <b>3</b>	Using diagonal elements from $P$ Correct method <i>Accept a.r.t. 0.196</i>
(iv)	$P^3 = \begin{pmatrix} 0.352 & 0.364 & 0.284 \\ 0.328 & 0.404 & 0.268 \\ 0.304 & 0.372 & 0.324 \end{pmatrix}$ $0.328864 \times 0.352 + 0.381536 \times 0.404 + 0.2896 \times 0.324$ $= 0.3637$	M1 M1 M1 A1 <b>4</b>	For evaluating $P^3$ Using diagonal elements from $P^3$ Correct method <i>Accept a.r.t. 0.364</i>
(v)	$Q = \begin{pmatrix} 0.3289 & 0.3816 & 0.2895 \\ 0.3289 & 0.3816 & 0.2895 \\ 0.3289 & 0.3816 & 0.2895 \end{pmatrix}$ <p>0.3289, 0.3816, 0.2895 are the long-run probabilities for the routes A, B, C</p>	B1B1B1  B1 <b>4</b>	<i>Deduct 1 if not given as a (3×3) matrix</i> <i>Deduct 1 if not 4 dp</i>  <i>Accept 'equilibrium probabilities'</i>
(vi)	$(0.4 \quad 0.2 \quad 0.4) \begin{pmatrix} 0.1 & 0.4 & 0.5 \\ 0.7 & 0.2 & 0.1 \\ a & b & c \end{pmatrix} = (0.4 \quad 0.2 \quad 0.4)$ <p>0.04 + 0.14 + 0.4a = 0.4, so <math>a = 0.55</math> 0.16 + 0.04 + 0.4b = 0.2, so <math>b = 0</math> 0.2 + 0.02 + 0.4c = 0.4, so <math>c = 0.45</math></p> <p>After C, routes A, B, C will be used with probabilities 0.55, 0, 0.45</p>	M1  M1  A2 <b>4</b>	Obtaining a value for a, b or c  Give A1 for one correct
(vii)	$0.4 \times 0.1 + 0.2 \times 0.2 + 0.4 \times 0.45$ $= 0.26$	M1 M1 A1 <b>3</b>	Using long-run probs 0.4, 0.2, 0.4 Using diag elements from new matrix



# 4758 Differential Equations

1 (i)	$2\ddot{x} = 2g - 8(x + 0.25g) - 2kv$ Weight positive as down, tension negative as up. Resistance negative as opposes motion. $\Rightarrow \ddot{x} + k\dot{x} + 4x = 0$	M1 N2L equation with all forces using given expressions for tension and resistance B1 B1 E1 Must follow correct N2L equation	4
(ii)	$x = A \cos 2t + B \sin 2t$ $t = 0, x = 0.1 \Rightarrow A = 0.1$ $\dot{x} = -2A \sin 2t + 2B \cos 2t$ so $t = 0, \dot{x} = 0 \Rightarrow B = 0$ $x = 0.1 \cos 2t$	B1 M1 Find the coefficient of cos M1 Find the coefficient of sin A1 cao	4
(iii)	$\alpha^2 + 2\alpha + 4 = 0$ $\alpha = -1 \pm \sqrt{3}j$ $x = e^{-t} (C \cos \sqrt{3}t + D \sin \sqrt{3}t)$ $t = 0, x = 0.1 \Rightarrow C = 0.1$ $\dot{x} = -e^{-t} (C \cos \sqrt{3}t + D \sin \sqrt{3}t)$ $+ e^{-t} (-\sqrt{3}C \sin \sqrt{3}t + \sqrt{3}D \cos \sqrt{3}t)$ $0 = -C + \sqrt{3}D$ $D = \frac{0.1}{\sqrt{3}}$ $x = 0.1e^{-t} \left( \cos \sqrt{3}t + \frac{1}{\sqrt{3}} \sin \sqrt{3}t \right)$ 	M1 Auxiliary equation A1 M1 CF for complex roots F1 CF for their roots M1 Condition on $x$ M1 Differentiate (product rule) M1 Condition on $\dot{x}$ A1 cao B1 Curve through (0,0.1) with zero gradient B1 Oscillating B1 Asymptote $x = 0$	11
(iv)	$k^2 - 4 \cdot 1 \cdot 4 > 0$ (As $k$ is positive) $k > 4$ 	M1 Use of discriminant A1 Correct inequality A1 Accept $k < -4$ in addition (but not $k > -4$ ) B1 Curve through (0,0.1) B1 Decays without oscillating (at most one intercept with positive $t$ axis)	5

2 (i) $x = Ae^{-2t}$ $t = 0, x = 8 \Rightarrow A = 8$ $x = 8e^{-2t}$	M1 Any valid method M1 Condition on $x$ A1	3
(ii) $\dot{y} + y = 16e^{-2t}$ $\alpha + 1 = 0 \Rightarrow \alpha = -1$ CF $y = Be^{-t}$ PI $y = ae^{-2t}$ $-2ae^{-2t} + ae^{-2t} = 16e^{-2t}$ $a = -16$ GS $y = -16e^{-2t} + Be^{-t}$ $t = 0, y = 0 \Rightarrow B = 16$ $y = 16(e^{-t} - e^{-2t})$ <b>Alternative mark scheme for first 7 marks:</b> $I = e^t$ $d(y e^t)/dt = 16e^{-t}$ $y e^t = -16e^{-t} + B$ $y = -16e^{-2t} + Be^{-t}$	M1 Substitute for $x$ M1 Auxiliary equation A1 B1 M1 Differentiate and substitute A1 cao F1 Their PI + CF (with one arbitrary constant) M1 Condition on $y$ F1 Follow a non-trivial GS  M1 Substitute for $x$ M1 Attempt integrating factor A1 IF correct B1 M1 Integrate A1 cao F1 Divide by their $I$ (must divide constant)	9
(iii) $y = 16e^{-t}(1 - e^{-t})$ $16e^{-t} > 0$ and $t > 0 \Rightarrow e^{-t} < 1$ hence $y > 0$ 	M1 Or equivalent (NB $e^{-t} > e^{-2t}$ needs justifying) E1 Complete argument B1 Starts at origin B1 General shape consistent with their solution and $y > 0$ B1 Tends to zero	5
(iv) $\frac{d}{dt}(x + y + z) = (-2x) + (2x - y) + (y) = 0$ $\Rightarrow x + y + z = c$ Hence initial conditions $\Rightarrow x + y + z = 8$ $z = 8 - x - y$ $z = 8(1 - 2e^{-t} + e^{-2t}) = 8(1 - e^{-t})^2$	M1 Consider sum of DE's E1 E1 M1 Substitute for $x$ and $y$ and find $z$ E1 Convincingly shown ( $x, y$ must be correct)	5
(v) $0.99 \times 8 = 8(1 - e^{-t})^2$ $t = -0.690638$ or $5.29581$ 99% is $Z$ after 5.30 hours	B1 Correct equation (any form) B1 Accept value in [5.29, 5.3]	2

<p>3</p> <p>(i) <math>\dot{y} + \frac{k}{t}y = 1</math></p> $I = \exp\left(\int \frac{k}{t} dt\right) = \exp(k \ln t) = t^k$ $t^k \dot{y} + kt^{k-1}y = t^k$ $\frac{d}{dt}(yt^k) = t^k$ $yt^k = \int t^k dt$ $= \frac{1}{k+1}t^{k+1} + A$ $y = \frac{1}{k+1}t + At^{-k}$ $t = 1, y = 0 \Rightarrow 0 = \frac{1}{k+1} + A \Rightarrow A = -\frac{1}{k+1}$ $y = \frac{1}{k+1}(t - t^{-k})$	<p>M1 Divide by <math>t</math> (condone LHS only)</p> <p>M1 Attempt integrating factor</p> <p>A1 Integrating factor</p> <p>F1 Multiply DE by their <math>I</math></p> <p>M1 LHS</p> <p>M1 Integrate</p> <p>A1 cao (including constant)</p> <p>F1 Divide by their <math>I</math> (must divide constant)</p> <p>M1 Use condition</p> <p>F1 Follow a non-trivial GS</p>	10												
<p>(ii) <math>y = \frac{1}{3}(t - t^{-2})</math></p> 	<p>B1 Shape consistent with their solution for <math>t \geq 1</math></p> <p>B1 Passes through (1, 0)</p> <p>B1 Behaviour for large <math>t</math></p>	3												
<p>(iii) <math>yt^{-1} = \int t^{-1} dt</math></p> $= \ln t + B$ $y = t(\ln t + B)$ $t = 1, y = 0 \Rightarrow B = 0 \Rightarrow y = t \ln t$	<p>M1 Follow their (i)</p> <p>A1 cao</p> <p>F1 Divide by their <math>I</math> (must divide constant)</p> <p>A1 cao</p>	4												
<p>(iv) <math>\frac{dy}{dt} = 1 + t^{-1} \sin y</math></p> <table border="1" data-bbox="287 1344 734 1478"> <thead> <tr> <th><math>t</math></th> <th><math>y</math></th> <th><math>dy/dt</math></th> </tr> </thead> <tbody> <tr> <td>1</td> <td>0</td> <td>1</td> </tr> <tr> <td>1.1</td> <td>0.1</td> <td>1.0908</td> </tr> <tr> <td>1.2</td> <td>0.2091</td> <td></td> </tr> </tbody> </table>	$t$	$y$	$dy/dt$	1	0	1	1.1	0.1	1.0908	1.2	0.2091		<p>M1 Rearrange DE (may be implied)</p> <p>M1 Use algorithm</p> <p>A1 <math>y(1.1)</math></p> <p>A1 <math>y(1.2)</math></p>	4
$t$	$y$	$dy/dt$												
1	0	1												
1.1	0.1	1.0908												
1.2	0.2091													
<p>(v) 0.2138 as smaller step size Decreasing step length has increased estimate. Assuming this estimate is more accurate, decreasing step length further will increase estimate further, so true value likely to be greater. Hence underestimates.</p> <p><i>Alternative mark scheme for last 2 marks: dy/dt seems to be increasing, hence Euler's method will underestimate true value + sketch (or explanation).</i></p>	<p>B1 Must give reason</p> <p>M1 Identify effect of decreasing step length</p> <p>A1 Convincing argument</p> <p>M1 Identify derivative increasing</p> <p>A1 Convincing argument</p>	3												

4	$\ddot{x} = 4\dot{x} - 6\dot{y} - 9\cos t$	M1	Differentiate first equation	
(i)	$= 4\dot{x} - 6(3x - 5y - 7\sin t) - 9\cos t$	M1	Substitute for $\dot{y}$	
	$y = \frac{1}{6}(4x - \dot{x} - 9\sin t)$	M1	$y$ in terms of $x, \dot{x}$	
	$\ddot{x} = 4\dot{x} - 18x + 5(4x - \dot{x} - 9\sin t) + 42\sin t - 9\cos t$	M1	Substitute for $y$	
	$\ddot{x} + \dot{x} - 2x = -3\sin t - 9\cos t$	E1	LHS	
		E1	RHS	6
(ii)	$\alpha^2 + \alpha - 2 = 0$	M1	Auxiliary equation	
	$\alpha = 1$ or $-2$	A1		
	CF $x = Ae^t + Be^{-2t}$	F1	CF for their roots	
	PI $x = a\cos t + b\sin t$	B1	PI of this form	
	$(-ac - bs) + (-as + bc) - 2(ac + bs) = -3s - 9c$	M1	Differentiate twice and substitute	
	$-a + b - 2a = -9$	M1	Compare coefficients (2 equations)	
	$-b - a - 2b = -3$	M1	Solve (2 equations)	
	$\Rightarrow a = 3, b = 0$	A1		
	$x = 3\cos t + Ae^t + Be^{-2t}$	F1	Their PI + CF (with two arbitrary constants)	9
(iii)	$y = \frac{1}{6}(4x - \dot{x} - 9\sin t)$	M1	$y$ in terms of $x, \dot{x}$	
	$= \frac{1}{6}(12\cos t + 4Ae^t + 4Be^{-2t} + 3\sin t - Ae^t + 2Be^{-2t} - 9\sin t)$	M1	Differentiate $x$ and substitute	
	$y = 2\cos t - \sin t + \frac{1}{2}Ae^t + Be^{-2t}$	A1	Constants must correspond with those in $x$	3
(iv)	$x$ bounded $\Rightarrow A = 0$	M1	Identify coefficient of exponentially growing term must be zero	
	$\Rightarrow y$ bounded	E1	Complete argument	2
(v)	$t = 0, y = 0 \Rightarrow 0 = B + 2 \Rightarrow B = -2$	M1	Condition on $y$	
	$x = 3\cos t - 2e^{-2t}, y = 2\cos t - \sin t - 2e^{-2t}$	F1	Follow their (non-trivial) general solutions	
	$x = 3\cos t$	A1	cao	
	$y = 2\cos t - \sin t$	A1	cao	4

## 4761 Mechanics 1

Q 1		mark	comment	sub
(i)	$N2L \uparrow 1000 - 100 \times 9.8 = 100a$ $a = 0.2$ so $0.2 \text{ m s}^{-2}$ upwards	M1 B1 A1	N2L. Accept $F = mga$ and no weight Weight correct (including sign). Allow if seen. Accept $\pm 0.2$ . Ignore units and direction	3
(ii)	$T_{BA} - 980 = 100 \times 0.8$ so tension is 1060 N	M1 A1	N2L. $F = ma$ . Weight present, no extras. Accept sign errors.	2
(iii)	$T_{BA} \cos 30 = 1060$  $T_{BA} = 1223.98\dots$ so 1220 N (3 s. f.)	M1 A1 A1	Attempt to resolve their (ii). Do not award for <b>their</b> 1060 resolved unless all forces present and all resolutions needed are attempted. If start again allow no weight. Allow $\sin \leftrightarrow \cos$ . No extra forces. Condone sign errors FT <b>their</b> 1060 only cao	3
		8		

Q 2		mark	comment	sub
(i)		B1	Sketch. O, i, j and r (only require correct quadrant.) Vectors must have arrows. Need not label r.	1
(ii)	$\sqrt{4^2 + (-5)^2}$ $= \sqrt{41}$ or 6.4031... so 6.40 (3 s. f.) Need $180 - \arctan\left(\frac{4}{5}\right)$ 141.340 so $141^\circ$	M1 A1 M1 A1	Accept $\sqrt{4^2 - 5^2}$ Or equivalent. Award for $\arctan\left(\pm\frac{4}{5}\right)$ or $\arctan\left(\pm\frac{5}{4}\right)$ or equivalent seen without 180 or 90. cao	4
(iii)	$12i - 15j$ or $\begin{pmatrix} 12 \\ -15 \end{pmatrix}$	B1	Do not award for magnitude given as the answer. Penalise spurious notation by 1 mark at most once in paper	1
		6		

Q 3		mark	comment	sub
(i)	$\mathbf{F} = 5 \begin{pmatrix} -1 \\ 2 \end{pmatrix} = \begin{pmatrix} -5 \\ 10 \end{pmatrix}$ so $\begin{pmatrix} -5 \\ 10 \end{pmatrix}$ N	M1 A1	Penalise spurious notation by 1 mark at most once in paper  Use of N2L in vector form  Ignore units. [Award 2 for answer seen] [SC1 for $\sqrt{125}$ or equiv seen]	2
(ii)	$\mathbf{s} = \begin{pmatrix} -2 \\ 3 \end{pmatrix} + 4 \begin{pmatrix} 4 \\ 5 \end{pmatrix} + \frac{1}{2} \times 4^2 \times \begin{pmatrix} -1 \\ 2 \end{pmatrix}$  $\mathbf{s} = \begin{pmatrix} 6 \\ 39 \end{pmatrix}$ so $\begin{pmatrix} 6 \\ 39 \end{pmatrix}$ m	M1 A1 B1	Use of $\mathbf{s} = t\mathbf{u} + 0.5t^2\mathbf{a}$ or integration of $\mathbf{a}$ . Allow $\mathbf{s}_0$ omitted. If integrated need to consider $\mathbf{v}$ when $t = 0$ Correctly evaluated; accept $\mathbf{s}_0$ omitted.  Correctly adding $\mathbf{s}_0$ to a vector (FT). Ignore units.  [NB $\begin{pmatrix} 8 \\ 36 \end{pmatrix}$ seen scores M1 A1]	3
		5		


Q 4		mark	comment	sub
(i)	The distance travelled by P is $0.5 \times 0.5 \times t^2$ The distance travelled by Q is $10t$	B1 B1	Accept $10t + 125$ if used correctly below.	2
(ii)	Meet when $0.25t^2 = 125 + 10t$  so $t^2 - 40t - 500 = 0$ Solving  $t = 50$ (or -10) Distance is $0.25 \times 50^2 = 625$ m	M1 F1  M1 A1 A1	Allow <b>their</b> wrong expressions for P and Q distances Allow $\pm 125$ or 125 omitted Award for <b>their</b> expressions as long as one is quadratic and one linear. Must have 125 with correct sign.  Accept any method that yields (smaller) + ve root of their 3 term quadratic  cao Allow -ve root not mentioned cao [SC2 400 m seen]	5
		7		

Q 5	mark	comment	sub
<p><b>either</b> Overall, N2L → <math>135 - 9 = (5 + 4)a</math></p> <p><math>a = 14</math> so <math>14 \text{ m s}^{-2}</math></p> <p>For A, N2L → <math>T - 9 = 4 \times 14</math> so 65 N</p> <p><b>or</b> <math>135 - T = 5a</math></p> <p><math>T - 9 = 4a</math> Solving <math>T = 65</math> so 65 N</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>Use of N2L. Allow <math>F = mga</math> but no extra forces. Allow 9 omitted.</p> <p>N2L on A or B with correct mass. <math>F = ma</math>. All relevant forces and no extras. cao</p> <p>* 1 equation in <math>T</math> and <math>a</math>. Allow sign errors. Allow <math>F = mga</math></p> <p>Both equations correct and consistent</p> <p>Dependent on M* solving for <math>T</math>. cao.</p>	4
	4		

Q 6	mark	comment	sub
<p>(i) <math>40 \times 0.6t - 5t^2</math></p> <p><math>= 24t - 5t^2</math></p>	<p>M1</p> <p>A1</p>	<p>Use of <math>s = ut + 0.5at^2</math> with <math>a = \pm 9.8, \pm 10</math>. Accept 40 or <math>40 \times 0.8</math> for 'u'.</p> <p>Any form</p>	2
<p>(ii) <b>either</b> Need zero vertical distance so <math>24t - 5t^2 = 0</math></p> <p>so <math>t = 0</math> or <math>t = 4.8</math></p> <p><b>or</b> Time to highest point, <math>T</math></p> <p><math>0 = 40 \times 0.6 - 10T</math> so <math>T = 2.4</math> and time of flight is 4.8</p> <p>range is <math>40 \times 0.8 \times 4.8 = 153.6</math></p> <p>so 154 m (3 s. f.)</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>Equate <b>their</b> <math>y</math> to zero. With fresh start must have correct <math>y</math>. Accept no reference to <math>t = 0</math> and the other root in any form. FT <b>their</b> <math>y</math> if gives <math>t &gt; 0</math></p> <p>Allow use of <math>u = 40</math> and <math>40 \times 0.8</math>. Award even if half range found.</p> <p>May be awarded for doubling half range later.</p> <p>Horiz cpt. Accept 0.6 instead of 0.8 only if consistent with expression in (i). FT <b>their</b> <math>t</math>.</p> <p>cao [NB Use of half range or half time to get 76.8... (<math>g = 10</math>) or 78.36... (<math>g = 9.8</math>) scores 2] [If range formula used: M1 sensible attempt at substitution; allow <math>\sin 2\alpha</math> wrong B1 <math>\sin 2\alpha</math> correct A1 all correct A1 cao]</p>	4
	6		


Q 7		mark	comment	sub
(i)	Continuous string: smooth ring: light string	E1 E1	One reason Another reason	2
(ii)	Resolve $\leftarrow$ : $60\cos\alpha - 60\cos\beta = 0$  (so $\cos\alpha = \cos\beta$ ) and so $\alpha = \beta$	M1 E1	[(ii) and (iii) may be argued using Lami or triangle of forces] Resolution and an equation or equivalent. Accept $s \leftrightarrow c$ . Accept a <i>correct</i> equation seen without method stated. Accept the use of 'T' instead of '60'. Shown. Must have stated method (allow $\rightarrow$ seen).	2
(iii)	Resolve $\uparrow$  $2 \times 60 \times \sin\alpha - 8g = 0$  so $\alpha = 40.7933\dots$ so $40.8^\circ$ (3 s. f.)	M1 B1 B1 A1 A1	Resolution and an equation. Accept $s \leftrightarrow c$ . Do not award for resolution that cannot give solution (e.g. horizontal) Both strings used (accept use of half weight), seen in an equation  $\sin\alpha$ or equivalent seen in an equation All correct	5
(iv)	Resolve $\rightarrow$ $10 + T_{QC} \cos 25 - T_{PC} \cos 45 = 0$  Resolve $\uparrow T_{PC} \sin 45 + T_{QC} \sin 25 - 8g = 0$  Solving  $T_{CQ} = 51.4701\dots$ so 51.5 N (3 s. f.) $T_{CP} = 80.1120\dots$ so 80.1 N (3 s. f.)	M1 M1 A1 M1 A1 M1 A1 F1	Recognise strings have different tensions. Resolution and an equation. Accept $s \leftrightarrow c$ . No extra forces. All forces present. Allow sign errors. Correct. Any form. Resolution and an equation. Accept $s \leftrightarrow c$ . No extra forces. All forces present. Allow sign errors. Correct. Any form. * A method that leads to at least one solution of a pair of simultaneous equations.  cao either tension other tension. Allow FT only if M1* awarded [Scale drawing: 1 <sup>st</sup> M1 then A1, A1 for answers correct to 2 s.f.]	8
		17		




Q 8		mark	comment	sub
(i)	10	B1		1
(ii)	$v = 36 + 6t - 6t^2$	M1 A1	Attempt at differentiation	2
(iii)	$a = 6 - 12t$	M1 F1	Attempt at differentiation	2
(iv)	Take $a = 0$ so $t = 0.5$ and $v = 37.5$ so $37.5 \text{ m s}^{-1}$	M1 A1 A1	Allow table if maximum indicated or implied FT <b>their a</b> cao Accept no justification given that this is maximum	3
(v)	<b>either</b> Solving $36 + 6t - 6t^2 = 0$  so $t = -2$ or $t = 3$ <b>or</b> Sub the values in the expression for $v$ Both shown to be zero A quadratic so the only roots <b>then</b> $x(-2) = -34$ $x(3) = 91$	M1 B1 E1  M1 E1 B1  B1 B1	A method for two roots using <b>their v</b> Factorization or formula or ... of <b>their</b> expression Shown  Allow just 1 substitution shown Both shown Must be a clear argument  cao cao	5
(vi)	$ x(3) - x(0)  +  x(4) - x(3) $ $=  91 - 10  +  74 - 91 $ $= 98$ so 98 m	M1 A1 A1	Considering two parts Either correct cao [SC 1 for $s(4) - s(0) = 64$ ]	3
(vii)	At the SP of $v$  $x(-2) = -34$ i.e. $< 0$ and $x(3) = 91$ i.e. $> 0$ Also $x(-4) = 42 > 0$ and $x(6) = -98 < 0$    so three times	M1    B1  B1	Or any other valid argument e.g. find all the zeros, sketch, consider sign changes. Must have some working. If only a sketch, must have correct shape.  Doing appropriate calculations e.g. find all 3 zeros; sketch cubic reasonably (showing 3 roots); sign changes in range  3 times seen	3
		19		

## 4762 Mechanics 2

Q 1	mark	comment	sub
(a) (i) In $i$ direction: $6u - 12 = 18$ so $u = 5$ i.e. $5i \text{ m s}^{-1}$  <b>either</b> In $i$ direction: $0.5v + 12 = 0.5 \times 11$  $v = -13$ so $-13i \text{ m s}^{-1}$ <b>or</b> $6 \times 5 + 0.5v = 6 \times 3 + 0.5 \times 11$ $v = -13$ so $-13i \text{ m s}^{-1}$	M1 E1  M1 B1 A1  M1 A1 A1	Use of I-M Accept $6u - 12 = 18$ as total working. Accept 5 instead of $5i$ .  Use of I-M Use of $+12i$ or equivalent Accept direction indicated by any means  PCLM Allow only sign errors Accept direction indicated by any means	5
(ii) Using NEL: $\frac{11-3}{-13-5} = -e$  $e = \frac{4}{9}$ (0.4)	M1  F1 F1	Use of NEL. Condone sign errors but not reciprocal expression  FT only <b>their</b> $-13$ (even if +ve) FT only <b>their</b> $-13$ and only if $-ve$ (allow 1 s.f. accuracy)	3
(iii) In $i$ direction: $-2 \times 7 = 0.5v - 0.5 \times 11$  $v = -17$ so $-17i \text{ m s}^{-1}$  <b>or</b> $-2i = 0.5a$ so $a = -4i \text{ m s}^{-2}$ $v = 11i - 4i \times 7$ $v = -17$ so $-17i \text{ m s}^{-1}$	M1 M1 A1 A1  M1 A1 M1 A1	Use of $I = Ft$ Use of $I = m(v - u)$ For $\pm 17$ cao. Direction (indicated by any means)  Use of $F = ma$ For $\pm 4$ Use of $uvast$ cao. Direction (indicated by any means)	4
(b) $u i + ev j$  $\tan \alpha = \frac{v}{u}$ , $\tan \beta = \frac{ev}{u}$  $\tan \beta = e \left( \frac{v}{u} \right) = e \tan \alpha$	B1 B1 M1  B1 E1	For $u$ For $ev$ Use of $\tan$ . Accept reciprocal argument. Accept use of <b>their</b> components  Both correct. Ignore signs. Shown. Accept signs not clearly dealt with.	5
	17		

Q 2	mark	comment	sub
(i) $(2+3 \times 6) \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = 6 \begin{pmatrix} 3 \\ 0 \end{pmatrix} + 6 \begin{pmatrix} 6 \\ 3 \end{pmatrix} + 6 \begin{pmatrix} 3 \\ 6 \end{pmatrix} + 2 \begin{pmatrix} 0 \\ 7 \end{pmatrix}$ $20 \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = \begin{pmatrix} 18+36+18 \\ 18+36+14 \end{pmatrix} = \begin{pmatrix} 72 \\ 68 \end{pmatrix}$ $\bar{x} = 3.6$ $\bar{y} = 3.4$	M1 B1 B1 B1 E1 A1	Method for c.m. Total mass correct For any of the 1 <sup>st</sup> 3 RHS terms For the 4 <sup>th</sup> RHS term cao [If separate cpts, award the 2 <sup>nd</sup> B1 for 2 x- terms correct and 3 <sup>rd</sup> B1 for 2 × 7 in y term]	6
(ii)  $\arctan\left(\frac{3.6}{2+(6-3.4)}\right) = \arctan\left(\frac{3.6}{4.6}\right)$ so 38.047... so 38.0° (3 s. f.)	B1 B1 M1 B1 A1	Diagram showing G vertically below D 3.6 and <b>their</b> 3.4 correctly placed (may be implied) Use of arctan on <b>their</b> lengths. Allow reciprocal of argument. Some attempt to calculate correct lengths needed 2 + (6 – <b>their</b> 3.4) seen cao	5
(iii) <p>moments about D  <math>5 \times 3.6 = 6 \times T_{BP}</math> so tension in BP is 3 N            Resolve vert: <math>3 + T_{DQ} = 5</math>            so tension in DQ is 2 N</p>	M1 F1 M1 F1	moments about D. No extra forces FT <b>their</b> values if calc 2nd Resolve vertically or moments about B. FT <b>their</b> values if calc 2nd	4
(iv) <p>We require x-cpt of c.m. to be zero  <b>either</b>  <math>(20+L)\bar{x} = 20 \times 3.6 - \frac{1}{2}L^2</math>  <b>or</b>  <math>2 \times 6 \times (0.5 \times 6) + 6 \times 6 - 0.5 \times L^2 = 0</math></p> $L = 12$	M1 B1 A1 A1	A method to achieve this with all cpts  For the $0.5 \times L^2$ All correct	4
	19		

Q 3		mark	comment	sub
(a) (i)		B1 B1	Internal forces all present and labelled All forces correct with labels and arrows (Allow the internal forces set as tensions, thrusts or a mixture)	2
(ii)	<p>A <math>\uparrow</math>  <math>T_{AD} \sin 30 - L = 0</math> so <math>T_{AD} = 2L</math> so <math>2L</math> N (T)  A <math>\rightarrow</math> <math>T_{AB} + T_{AD} \cos 30 = 0</math>  so <math>T_{AB} = -\sqrt{3}L</math> so <math>\sqrt{3}L</math> N (C)  B <math>\uparrow</math> <math>T_{BD} \sin 60 - 3L = 0</math>  so <math>T_{BD} = 2\sqrt{3}L</math> so <math>2\sqrt{3}L</math> N (T)  B <math>\rightarrow</math>  <math>T_{BC} + T_{BD} \cos 60 - T_{AB} = 0</math>  so <math>T_{BC} = -2\sqrt{3}L</math> so <math>2\sqrt{3}L</math> N (C)</p>	M1 A1  M1 F1  M1 A1  M1 F1  E1	Equilibrium equation at a pin-joint attempted 1 <sup>st</sup> ans. Accept + or -.  Second equation attempted 2 <sup>nd</sup> ans. FT any previous answer(s) used.  Third equation attempted 3 <sup>rd</sup> ans. FT any previous answer(s) used.  Fourth equation attempted 4 <sup>th</sup> ans. FT any previous answer(s) used.  All T/C consistent [SC 1 all T/C correct WWW]	9
(b)	<p>Leg QR with frictional force <math>F \leftarrow</math>  moments c.w. about R  <math>U \times 2l \sin 60 - Wl \cos 60 = 0</math></p> <p>Horiz equilibrium for QR  <math>F = U</math></p> <p>Hence <math>\frac{1}{2}W = \sqrt{3}F</math>  and so <math>F = \frac{\sqrt{3}}{6}W</math></p>	M1 A1 A1  M1  E1 M1  E1	Accept only 1 leg considered (and without comment)  Suitable moments equation. Allow 1 force omitted a.c. moments c.w. moments  A second correct equation for horizontal or vertical equilibrium to eliminate a force (U or reaction at foot) [Award if correct moments equation containing only W and F]  * This second equation explicitly derived Correct use of 2 <sup>nd</sup> equation with the moments equation  Shown. CWO but do not penalise * again.	7
		18		

Q 4	mark	comment	sub
(a) (i) Tension is perp to the motion of the sphere (so WD, $Fd \cos \theta = 0$ )	E1		1
(ii) Distance dropped is $2 - 2 \cos 40 = 0.467911..$  GPE is $mgh$ so $0.15 \times 9.8 \times 0.467911... = 0.687829... \text{ J}$	M1 E1 M1 B1	Attempt at distance with resolution used. Accept $\sin \leftrightarrow \cos$ Accept seeing $2 - 2 \cos 40$ Any reasonable accuracy	4
(iii) $0.5 \times 0.15 \times v^2 = 0.687829...$ so $v = 3.02837... \text{ so } 3.03 \text{ m s}^{-1} \text{ (3 s. f.)}$	M1 F1	Using KE + GPE constant FT <b>their</b> GPE	2
(iv) $\frac{1}{2} \times 0.15 (v^2 - 2.5^2)$  $= 0.687829... - 0.6 \times \frac{40}{360} \times 2\pi \times 2$  $v = 2.06178... \text{ so } 2.06 \text{ m s}^{-1} \text{ (3 s. f.)}$	M1 B1 M1 A1 A1	Use of W-E equation (allow 1 KE term or GPE term omitted) KE terms correct WD against friction WD against friction correct (allow sign error) cao	5
(b) N2L down slope: $3g \sin 30 - F = 3 \times \frac{1}{8}g$  so $F = \frac{9g}{8} \text{ (= 11.025)}$  $R = 3g \times \frac{\sqrt{3}}{2} \text{ (= 25.4611...)}$  $\mu = \frac{F}{R} = \frac{\sqrt{3}}{4} \text{ (= 0.43301...)}$	M1 A1 A1 B1 M1 E1	Must have attempt at weight component Allow sign errors.  Use of $F = \mu R$ Must be worked precisely	6
	18		

## 4763 Mechanics 3

1(a)(i)	$[ \text{Velocity} ] = \text{LT}^{-1}$ $[ \text{Acceleration} ] = \text{LT}^{-2}$ $[ \text{Force} ] = \text{MLT}^{-2}$	B1 B1 B1 <b>3</b>	<i>(Deduct 1 mark if kg, m, s are consistently used instead of M, L, T)</i>
(ii)	$[ \lambda ] = \frac{[ \text{Force} ]}{[ v^2 ]} = \frac{\text{MLT}^{-2}}{(\text{LT}^{-1})^2}$ $= \text{ML}^{-1}$	M1 A1 cao <b>2</b>	
(iii)	$\left[ \frac{U^2}{2g} \right] = \frac{(\text{LT}^{-1})^2}{\text{LT}^{-2}} = \text{L}$ $\left[ \frac{\lambda U^4}{4mg^2} \right] = \frac{(\text{ML}^{-1})(\text{LT}^{-1})^4}{\text{M}(\text{LT}^{-2})^2}$ $= \frac{\text{ML}^3 \text{T}^{-4}}{\text{ML}^2 \text{T}^{-4}} = \text{L}$ $[ H ] = \text{L}$ ; all 3 terms have the same dimensions	B1 cao M1 A1 cao E1 <b>4</b>	<i>(Condone constants left in)</i>  <i>Dependent on B1M1A1</i>
(iv)	$(\text{ML}^{-1})^2 (\text{LT}^{-1})^\alpha \text{M}^\beta (\text{LT}^{-2})^\gamma = \text{L}$ $\beta = -2$ $-2 + \alpha + \gamma = 1$ $-\alpha - 2\gamma = 0$ $\alpha = 6$ $\gamma = -3$	B1 cao M1 A1 A1 cao A1 cao <b>5</b>	At least one equation in $\alpha, \gamma$ One equation correct

<b>(b)</b>	$EE \text{ is } \frac{1}{2} \times \frac{2060}{24} \times 6^2 \quad (=1545)$ $(PE \text{ gained}) = (EE \text{ lost}) + (KE \text{ lost})$	B1	
		M1	Equation involving PE, EE and KE Can be awarded from start to point where string becomes slack <i>or</i> any complete method (e.g. SHM) for finding $v^2$ at natural length If B0, give A1 for $v^2 = 88.2$ correctly obtained
	$50 \times 9.8 \times h = 1545 + \frac{1}{2} \times 50 \times 12^2$ $490h = 1545 + 3600$ $h = 10.5$	F1	<i>or</i> $0 = 88.2 - 2 \times 9.8 \times s \quad (s = 4.5)$
	$OA = 30 - h = 19.5 \text{ m}$	A1	<b>Notes</b> $\frac{1}{2} \times \frac{2060}{24} \times 6$ <i>used as EE can</i> <b>4 earn B0M1F1A0</b> $\frac{2060}{24} \times 6$ <i>used as EE gets B0M0</i>

<b>2 (i)</b>	$T \cos \alpha = mg$ $3.92 \cos \alpha = 0.3 \times 9.8$ $\cos \alpha = 0.75$ Angle is $41.4^\circ$ (0.723 rad)	M1 A1 <b>2</b>	Resolving vertically (Condone sin / cos mix for M marks throughout this question)
<b>(ii)</b>	$T \sin \alpha = m \frac{v^2}{r}$ $3.92 \sin \alpha = 0.3 \times \frac{v^2}{4.2 \sin \alpha}$ Speed is $4.9 \text{ m s}^{-1}$	M1 B1 A1 A1 <b>4</b>	Force and acceleration towards centre (condone $v^2 / 4.2$ or $4.2\omega^2$ ) For radius is $4.2 \sin \alpha$ (= 2.778) Not awarded for equation in $\omega$ unless $v = (4.2 \sin \alpha)\omega$ also appears
<b>(iii)</b>	$T - mg \cos \theta = m \frac{v^2}{a}$ $T - 0.3 \times 9.8 \times \cos 60^\circ = 0.3 \times \frac{8.4^2}{4.2}$ Tension is 6.51 N	M1 A1 A1 <b>3</b>	Forces and acceleration towards O
<b>(iv)</b>	$\frac{1}{2} mv^2 - mg \times 4.2 \cos \theta = \frac{1}{2} m \times 8.4^2 - mg \times 4.2 \cos 60^\circ$ $v^2 - 82.32 \cos \theta = 70.56 - 41.16$ $v^2 = 29.4 + 82.32 \cos \theta$	M1 M1 A1 E1 <b>4</b>	For $(- )mg \times 4.2 \cos \theta$ in PE Equation involving $\frac{1}{2}mv^2$ and PE
<b>(v)</b>	$(T) - mg \cos \theta = m \frac{v^2}{a}$ $(T) - m \times 9.8 \cos \theta = m \times \frac{29.4 + 82.32 \cos \theta}{4.2}$ String becomes slack when $T = 0$ $-9.8 \cos \theta = 7 + 19.6 \cos \theta$ $\cos \theta = -\frac{7}{29.4}$ $\theta = 104^\circ$ (1.81 rad)	M1 M1 A1 M1 A1 <b>5</b>	Force and acceleration towards O Substituting for $v^2$  <i>Dependent on first M1</i>  No marks for $v = 0 \Rightarrow \theta = 111^\circ$



3 (i)	$T_{PB} = 35(x - 3.2) \quad [= 35x - 112]$ $T_{BQ} = 5(6.5 - x - 1.8)$ $= 5(4.7 - x) \quad [= 23.5 - 5x]$	B1 M1  A1  <b>3</b>	Finding extension of BQ
(ii)	$T_{BQ} + mg - T_{PB} = m \frac{d^2x}{dt^2}$ $5(4.7 - x) + 2.5 \times 9.8 - 35(x - 3.2) = 2.5 \frac{d^2x}{dt^2}$ $160 - 40x = 2.5 \frac{d^2x}{dt^2}$ $\frac{d^2x}{dt^2} = 64 - 16x$	M1  A2   E1  <b>4</b>	Equation of motion (condone one missing force)  Give A1 for three terms correct
(iii)	At the centre, $\frac{d^2x}{dt^2} = 0$ $x = 4$	M1 A1  <b>2</b>	
(iv)	$\omega^2 = 16$ Period is $\frac{2\pi}{\sqrt{16}} = \frac{1}{2}\pi = 1.57 \text{ s}$	M1 A1  <b>2</b>	Seen or implied ( <i>Allow M1 for <math>\omega = 16</math></i> )  Accept $\frac{1}{2}\pi$
(v)	Amplitude $A = 4.4 - 4 = 0.4 \text{ m}$ Maximum speed is $A\omega$ $= 0.4 \times 4 = 1.6 \text{ m s}^{-1}$	B1 ft M1 A1 cao  <b>3</b>	ft is   4.4 - (iii)
(vi)	$x = 4 + 0.4 \cos 4t$ $v = (-) 1.6 \sin 4t$ When $v = 0.9$ , $\sin 4t = -\frac{0.9}{1.6}$ $4t = \pi + 0.5974$  Time is 0.935 s	M1 A1  M1  A1 cao  <b>4</b>	For $v = C \sin \omega t$ or $C \cos \omega t$ <i>This M1A1 can be earned in (v)</i>  Fully correct method for finding the required time e.g. $\frac{1}{4} \arcsin \frac{0.9}{1.6} + \frac{1}{2} \text{ period}$
	OR $0.9^2 = 16(0.4^2 - y^2)$ $y = -0.3307$  $y = 0.4 \cos 4t$ $\cos 4t = -\frac{0.3307}{0.4}$ $4t = \pi + 0.5974$ Time is 0.935 s	M1  A1  M1 A1 cao	Using $v^2 = \omega^2(A^2 - y^2)$ <i>and</i> $y = A \cos \omega t$ or $A \sin \omega t$ For $y = (\pm) 0.331$ <i>and</i> $y = 0.4 \cos 4t$


<b>4</b> <b>(a)(i)</b>	$V = \int \pi x^2 dy = \int_0^8 \pi (4 - \frac{1}{2}y) dy$ $= \pi \left[ 4y - \frac{1}{4}y^2 \right]_0^8 = 16\pi$ $V \bar{y} = \int \pi y x^2 dy$ $= \int_0^8 \pi y (4 - \frac{1}{2}y) dy$ $= \pi \left[ 2y^2 - \frac{1}{6}y^3 \right]_0^8 = \frac{128}{3}\pi$ $\bar{y} = \frac{\frac{128}{3}\pi}{16\pi}$ $= \frac{8}{3} (\approx 2.67)$	M1 A1 M1 A1 A1 M1 A1 <b>7</b>	$\pi$ may be omitted throughout Limits not required for M marks throughout this question  Dependent on M1M1
<b>(ii)</b>	CM is vertically above lower corner  $\tan \theta = \frac{2}{\bar{y}} = \frac{2}{\frac{8}{3}} (= \frac{3}{4})$ $\theta = 36.9^\circ (= 0.6435 \text{ rad})$	M1 M1 A1 A1 <b>4</b>	Trig in a triangle including $\theta$ Dependent on previous M1 Correct expression for $\tan \theta$ or $\tan(90 - \theta)$ Notes $\tan \theta = \frac{2}{\text{cand's } \bar{y}} \text{ implies M1M1A1}$ $\tan \theta = \frac{\text{cand's } \bar{y}}{2} \text{ implies M1M1}$ $\tan \theta = \frac{1}{\text{cand's } \bar{y}} \text{ without further evidence is MOMO}$




## 4764 Mechanics 4

<p>1(i) If <math>\delta m</math> is change in mass over time <math>\delta t</math>            PCLM <math>mv = (m + \delta m)(v + \delta v) +  \delta m (v - u)</math> [N.B. <math>\delta m &lt; 0</math>]  <math>(m + \delta m)\frac{\delta v}{\delta t} + u\frac{\delta m}{\delta t} = 0 \Rightarrow m\frac{dv}{dt} = -u\frac{dm}{dt}</math>  <math>\frac{dm}{dt} = -k \Rightarrow m = m_0 - kt</math>  <math>\Rightarrow (m_0 - kt)\frac{dv}{dt} = uk</math></p>	<p>M1 Change in momentum over time <math>\delta t</math>            M1 Rearrange to produce DE            A1 Accept sign error            M1 Find <math>m</math> in terms of <math>t</math>            E1 Convincingly shown</p>	5
<p>(ii) <math>v = \int \frac{uk}{m_0 - kt} dt</math>  <math>= -u \ln(m_0 - kt) + c</math>  <math>t = 0, v = 0 \Rightarrow c = u \ln m_0</math>  <math>v = u \ln\left(\frac{m_0}{m_0 - kt}\right)</math></p>	<p>M1 Separate and integrate            A1 cao (allow no constant)            M1 Use initial condition            A1 All correct</p>	4
<p>(iii) <math>m = \frac{1}{3}m_0 \Rightarrow m_0 - kt = \frac{1}{3}m_0</math>  <math>\Rightarrow v = u \ln 3</math></p>	<p>M1 Find expression for mass or time            A1 Or <math>t = 2m_0 / 3k</math>            A1</p>	3
<p>2(i) <math>P = Fv</math>  <math>= mv\frac{dv}{dx}</math>  <math>\Rightarrow mv^2\frac{dv}{dx} = m(k^2 - v^2)</math>  <math>\Rightarrow \frac{v^2}{k^2 - v^2}\frac{dv}{dx} = 1</math>  <math>\Rightarrow \left(\frac{k^2}{k^2 - v^2} - 1\right)\frac{dv}{dx} = 1</math>  <math>\int \left(\frac{k^2}{k^2 - v^2} - 1\right) dv = \int dx</math>  <math>\frac{1}{2}k \ln\left(\frac{k+v}{k-v}\right) - v = x + c</math>  <math>x = 0, v = 0 \Rightarrow c = 0</math>  <math>x = \frac{1}{2}k \ln\left(\frac{k+v}{k-v}\right) - v</math></p>	<p>M1 Used, not just quoted            M1 Use N2L and expression for acceleration            A1 Correct DE            M1 Rearrange            E1 Convincingly shown            M1 Separate and integrate            A1 LHS            M1 Use condition            A1 cao</p>	9
<p>(ii) Terminal velocity when acceleration zero  <math>\Rightarrow v = k</math>  <math>v = 0.9k \Rightarrow x = \frac{1}{2}k \ln\left(\frac{1.9}{0.1}\right) - 0.9k = \left(\frac{1}{2}\ln 19 - 0.9\right)k \approx 0.572k</math></p>	<p>M1            A1            F1 Follow their solution to (i)</p>	3

3(i) $M = \int_0^a k(a+r)2\pi r dr$ $= 2k\pi \left[ \frac{1}{2}ar^2 + \frac{1}{3}r^3 \right]_0^a$ $= \frac{5}{3}k\pi a^3$ $I = \int_0^a k(a+r)2\pi r \cdot r^2 dr$ $= 2k\pi \left[ \frac{1}{4}ar^4 + \frac{1}{5}r^5 \right]_0^a$ $= \frac{9}{10}k\pi a^5$ $= \frac{27}{50}Ma^2$	M1 Use circular elements (for $M$ or $I$ ) M1 Integral for mass M1 Integrate (for $M$ or $I$ ) A1 For [...] E1 M1 Integral for $I$ A1 For [...] A1 cao E1 Complete argument (including mass)	9
(ii) $I = 13.5$ $0.625 \times 50 = I\omega$ $\Rightarrow \omega \approx 2.31$	B1 Seen or used (here or later) M1 Use angular momentum M1 Use moment of impulse A1 cao	4
(iii) $\ddot{\theta} = \frac{30 - 2.31}{20} \approx 1.38$ Couple = $I\ddot{\theta}$ $\approx 18.7$	M1 Find angular acceleration M1 Use equation of motion F1 Follow their $\omega$ and $I$	3
(iv) $I\ddot{\theta} = -3\dot{\theta}$ $I \frac{d\dot{\theta}}{dt} = -3\dot{\theta}$ $\int \frac{d\dot{\theta}}{\dot{\theta}} = \int -\frac{3}{I} dt$ $\ln \dot{\theta}  = -\frac{t}{4.5} + c$ $\dot{\theta} = Ae^{-t/4.5}$ $t = 0, \dot{\theta} = 30 \Rightarrow A = 30$ $\dot{\theta} = 30e^{-t/4.5}$	B1 Allow sign error and follow their $I$ (but not $M$ ) M1 Set up DE for $\dot{\theta}$ (first order) M1 Separate and integrate B1 $\ln(\text{multiple of } \dot{\theta})$ seen M1 Rearrange, dealing properly with constant M1 Use condition on $\dot{\theta}$ A1	7
(v) Model predicts $\dot{\theta}$ never zero in finite time.	B1	1

<p>4(i) <math>V = \frac{1}{2} \left( \frac{mg}{10a} \right) (a\theta)^2 + mga \cos \theta</math> (relative to centre of pulley)</p> $\frac{dV}{d\theta} = \frac{1}{2} \left( \frac{mg}{10a} \right) \cdot 2a^2\theta - mga \sin \theta$ $\frac{dV}{d\theta} = mga \left( \frac{1}{10}\theta - \sin \theta \right)$	<p>M1 EPE term</p> <p>B1 Extension = <math>a\theta</math></p> <p>M1 GPE relative to any zero level</p> <p>A1 (<math>\pm</math> constant)</p> <p>M1 Differentiate</p> <p>E1</p>	6
<p>(ii) <math>\theta = 0 \Rightarrow \frac{dV}{d\theta} = mga \left( \frac{1}{10}(0) - \sin 0 \right) = 0</math></p> <p>hence equilibrium</p> $\frac{d^2V}{d\theta^2} = mga \left( \frac{1}{10} - \cos \theta \right)$ $V''(0) = -0.9mga < 0$ <p>hence unstable</p>	<p>M1 Consider value of <math>\frac{dV}{d\theta}</math></p> <p>E1</p> <p>M1 Differentiate again</p> <p>A1</p> <p>M1 Consider sign of <math>V''</math></p> <p>E1 <math>V''</math> must be correct</p>	6
<p>(iii) If the pulley is smooth, then the tension in the string is constant. Hence the EPE term is valid.</p>	<p>B1</p> <p>B1</p>	2
<p>(iv) Equilibrium positions at <math>\theta = 2.8</math>, <math>\theta = 7.1</math> and <math>\theta = 8.4</math></p> <p>From graph, <math>V''(2.8) = mgaf'(2.8) &gt; 0</math> hence stable at <math>\theta = 2.8</math> <math>V''(7.1) = mgaf'(7.1) &lt; 0 \Rightarrow</math> unstable at <math>\theta = 7.1</math> <math>V''(8.4) = mgaf'(8.4) &gt; 0 \Rightarrow</math> stable at <math>\theta = 8.4</math></p>	<p>B1 One correct</p> <p>B1 All three correct, no extras Accept answers in <math>[2.7, 3.0)</math>, <math>[7, 7.2]</math>, <math>[8.3, 8.5]</math></p> <p>M1 Consider sign of <math>V''</math> or <math>f'</math></p> <p>A1</p> <p>A1 Accept no reference to <math>V''</math> for one conclusion but other two must relate to sign of <math>V''</math>, not just <math>f'</math>.</p> <p>A1</p>	6
<p>(v)</p> 	<p>B1 P in approximately correct place</p> <p>B1 B in approximately correct place</p>	2
<p>(vi) If <math>\theta &lt; 0</math> then expression for EPE not valid hence not necessarily an equilibrium position.</p>	<p>M1</p> <p>A1</p>	2

# 4766 Statistics 1

<b>Q1</b> <b>(i)</b>	Mean = 7.35 (or better) Standard deviation: 3.69 – 3.70 (awfw) Allow $s^2 = 13.62$ to 13.68 Allow $\text{rmsd} = 3.64 - 3.66$ (awfw) <b>After B0, B0 scored</b> then if at least 4 correct mid-points seen or used. {1.5, 4, 6, 8.5, 15} Attempt of their mean = $\frac{\sum fx}{44}$ , with $301 \leq fx \leq 346$ and $fx$ strictly from mid-points not class widths or top/lower boundaries.	B2cao $\sum fx = 323.5$ B2cao $\sum fx^2 = 2964.25$ (B1) for variance s.o.i.o (B1) for rmsd (B1) mid-points (B1) $6.84 \leq \text{mean} \leq 7.86$	<b>4</b>
<b>(ii)</b>	Upper limit = $7.35 + 2 \times 3.69 = 14.73$ or 'their sensible mean' + $2 \times$ 'their sensible s.d.' So there could be one or more outliers	M1 ( with s.d. < mean) E1dep on B2, B2 earned and comment	<b>2</b>
<b>TOTAL</b>		<b>6</b>	
<b>Q2</b> <b>(i)</b>	$P(W) \times P(C) = 0.20 \times 0.17 = 0.034$ $P(W \cap C) = 0.06$ (given in the question) Not equal so not independent (Allow $0.20 \times 0.17 \neq 0.06$ or $\neq p(W \cap C)$ so not independent).	M1 for multiplying or 0.034 seen A1 (numerical justification needed)	<b>2</b>
<b>(ii)</b>	<div style="text-align: center;">  </div> <p>The last two G marks are independent of the labels</p>	G1 for two <b>overlapping</b> circles labelled G1 for 0.06 and either 0.14 or 0.11 in the <b>correct places</b> G1 for all 4 <b>correct</b> probs in the <b>correct</b> places (including the 0.69) NB No credit for Karnaugh maps here	<b>3</b>
<b>(iii)</b>	$P(W C) = \frac{P(W \cap C)}{P(C)} = \frac{0.06}{0.17} = \frac{6}{17} = 0.353 \text{ (awrt 0.35)}$	M1 for 0.06 / 0.17 A1 cao	<b>2</b>

(iv)	Children are more likely than adults to be able to speak Welsh or 'proportionally more children speak Welsh than adults'  Do not accept: 'more Welsh children speak Welsh than adults'	E1FT Once the correct idea is seen, apply ISW	1
		<b>TOTAL</b>	<b>8</b>
<b>Q3</b>			
(i)	(A) $0.5 + 0.35 + p + q = 1$ so $p + q = 0.15$  (B) $0 \times 0.5 + 1 \times 0.35 + 2p + 3q = 0.67$ so $2p + 3q = 0.32$  (C) from above $2p + 2q = 0.30$ so $q = 0.02, p = 0.13$	B1 $p + q$ in a correct equation before they reach $p + q = 0.15$  B1 $2p + 3q$ in a correct equation before they reach $2p + 3q = 0.32$  (B1) for any 1 correct answer B2 for both correct answers	1  1  2
(ii)	$E(X^2) = 0 \times 0.5 + 1 \times 0.35 + 4 \times 0.13 + 9 \times 0.02 = 1.05$  $\text{Var}(X) = \text{'their 1.05'} - 0.67^2 = 0.6011$ (awrt 0.6)  (M1, M1 can be earned with their $p^+$ and $q^+$ but not A mark)	M1 $\sum x^2 p$ (at least 2 non zero terms correct) M1dep for $(- 0.67^2)$ , provided $\text{Var}(X) > 0$ A1 cao (No n or n-1 divisors)	3
		<b>TOTAL</b>	<b>7</b>
<b>Q4</b>			
(i)	$X \sim B(8, 0.05)$ (A) $P(X = 0) = 0.95^8 = 0.6634$ 0.663 or better  Or using tables $P(X = 0) = 0.6634$  (B) $P(X = 1) = \binom{8}{1} \times 0.05 \times 0.95^7 = 0.2793$  $P(X > 1) = 1 - (0.6634 + 0.2793) = 0.0573$  Or using tables $P(X > 1) = 1 - 0.9428 = 0.0572$	M1 $0.95^8$ A1 CAO Or B2 (tables)  M1 for $P(X = 1)$ (allow 0.28 or better) M1 for $1 - P(X \leq 1)$ must have both probabilities A1cao (0.0572 - 0.0573)  M1 for $P(X \leq 1)$ 0.9428 M1 for $1 - P(X \leq 1)$ A1 cao (must end in...2)	2  3
(ii)	Expected number of days = $250 \times 0.0572 = 14.3$ awrt	M1 for $250 \times \text{prob}(B)$ A1 FT but no rounding at end	2
		<b>TOTAL</b>	<b>7</b>



<b>Q5</b> <b>(i)</b>	<p>Let <math>p</math> = probability of remembering or naming all items (for population) (whilst listening to music.)  <math>H_0: p = 0.35</math>  <math>H_1: p &gt; 0.35</math></p> <p><math>H_1</math> has this form since the student believes that the probability will be increased/ improved/ got better /gone up.</p>	<p>B1 for definition of <math>p</math>            B1 for <math>H_0</math>            B1 for <math>H_1</math></p> <p>E1dep on <math>p &gt; 0.35</math> in <math>H_0</math>            In words not just because <math>p &gt; 0.35</math></p>	4
<b>(ii)</b>	<p>Let <math>X \sim B(15, 0.35)</math>  <b>Either:</b> <math>P(X \geq 8) = 1 - 0.8868 = 0.1132 &gt; 5\%</math>            Or <math>0.8868 &lt; 95\%</math></p> <p>So not enough evidence to reject <math>H_0</math> (Accept <math>H_0</math>)</p> <p>Conclude that there is not enough evidence to indicate that the probability of remembering all of the items is improved / improved/ got better /gone up. (when listening to music.)</p> <p>-----</p> <p><b>Or:</b></p> <p>Critical region for the test is <math>\{9, 10, 11, 12, 13, 14, 15\}</math>            8 does not lie in the critical region.</p> <p>So not enough evidence to reject <math>H_0</math></p> <p>Conclude that there is not enough evidence to indicate that the probability of remembering all of the items is improved / improved/ got better /gone up. (when listening to music.)</p> <p>-----</p> <p><b>Or:</b></p> <p>The smallest critical region that 8 could fall into is <math>\{8, 9, 10, 11, 12, 13, 14, \text{ and } 15\}</math>. The size of this region is 0.1132</p> <p><math>0.1132 &gt; 5\%</math></p> <p>So not enough evidence to reject <math>H_0</math></p> <p>Conclude that there is not enough evidence to indicate that the probability of remembering all of the items is improved (when listening to music)</p>	<p><b>Either:</b>            M1 for probability (0.1132)            M1dep for comparison            A1dep</p> <p>E1dep on all previous marks for conclusion in context</p> <p>-----</p> <p><b>Or:</b></p> <p>M1 for correct CR (no omissions or additions)            M1dep for 8 does not lie in CR            A1dep</p> <p>E1dep on all previous marks for conclusion in context</p> <p>-----</p> <p><b>Or:</b></p> <p>M1 for CR <math>\{8, 9, \dots, 15\}</math> and size = 0.1132            M1 dep for comparison            A1dep</p> <p>E1dep on all previous marks for conclusion in context</p>	4
		<b>TOTAL</b>	<b>8</b>

	Section B		
<b>Q6</b>			
<b>(i)</b>	<p>(A) <math>P(\text{both rest of UK}) = 0.20 \times 0.20</math>  <math>= 0.04</math></p> <p>(B) <i>Either: All 5 case</i>  <math>P(\text{at least one England}) =</math>  <math>(0.79 \times 0.20) + (0.79 \times 0.01) + (0.20 \times 0.79) + (0.01 \times 0.79) +</math>  <math>(0.79 \times 0.79)</math>  <math>= 0.158 + 0.0079 + 0.158 + 0.0079 + 0.6241 = 0.9559</math></p> <p><i>Or</i></p> <p><math>P(\text{at least one England}) = 1 - P(\text{neither England})</math>  <math>= 1 - (0.21 \times 0.21) = 1 - 0.0441 = 0.9559</math>  <i>or listing all</i>  <math>= 1 - \{ (0.2 \times 0.2) + (0.2 \times 0.01) + (0.01 \times 0.20) + (0.01 \times</math>  <math>0.01) \}</math>  <math>= 1 - (**)</math>  <math>= 1 - \{ 0.04 + 0.002 + 0.002 + 0.0001 \}</math>  <math>= 1 - 0.0441</math>  <math>= 0.9559</math></p> <p><i>Or: All 3 case</i>  <math>P(\text{at least one England}) =</math>  <math>= 0.79 \times 0.21 + 0.21 \times 0.79 + 0.79^2</math>  <math>= 0.1659 + 0.1659 + 0.6241</math>  <math>= 0.9559</math></p> <hr/> <p>(C) <i>Either</i>  <math>0.79 \times 0.79 + 0.79 \times 0.2 + 0.2 \times 0.79 + 0.2 \times 0.2 = 0.9801</math></p> <p><i>Or</i>  <math>0.99 \times 0.99 = 0.9801</math></p> <p><i>Or</i>  <math>1 - \{ 0.79 \times 0.01 + 0.2 \times 0.01 + 0.01 \times 0.79 + 0.01 \times 0.02 +</math>  <math>0.01^2 \} = 1 - 0.0199</math>  <math>= 0.9801</math></p>	<p>M1 for multiplying A1cao</p> <p>M1 for any correct term (3case or 5case) M1 for correct sum of all 3 (or of all 5) with <b>no extras</b> A1cao (condone 0.96 www)</p> <p><i>Or</i> M1 for <math>0.21 \times 0.21</math> or for (**) fully enumerated or 0.0441 seen M1<b>dep</b> for <math>1 - (1^{\text{st}} \text{ part})</math> A1cao</p> <p>See above for 3 case</p> <hr/> <p>M1 for sight of all 4 correct terms summed A1 cao (condone 0.98 www) <i>or</i> M1 for <math>0.99 \times 0.99</math> A1cao <b>Or</b> M1 for everything <math>1 - \{ \dots \}</math> A1cao</p>	<p><b>2</b></p> <p><b>3</b></p> <p><b>2</b></p>
<b>(ii)</b>	<p><math>P(\text{both the rest of the UK   neither overseas})</math>  <math>= \frac{P(\text{the rest of the UK and neither overseas})}{P(\text{neither overseas})}</math>  <math>= \frac{0.04}{0.9801} = 0.0408</math></p> <p>{Watch for: <math>\frac{\text{answer}(A)}{\text{answer}(C)}</math> as evidence of method (<math>p &lt; 1</math>)}</p>	<p>M1 for numerator of 0.04 or 'their answer to (i)(A)'</p> <p>M1 for denominator of 0.9801 or 'their answer to (i) (C)' A1 FT (<math>0 &lt; p &lt; 1</math>) 0.041 at least</p>	<b>3</b>

<p><b>(iii)</b></p> <p>(A) Probability = <math>1 - 0.79^5</math>  <math>= 1 - 0.3077</math>  <math>= 0.6923</math> (accept awrt 0.69)</p> <p>see additional notes for alternative solution</p> <p>(B) <math>1 - 0.79^n &gt; 0.9</math></p> <p><b>EITHER:</b>  <math>1 - 0.79^n &gt; 0.9</math> or <math>0.79^n &lt; 0.1</math>        (condone = and <math>\geq</math> throughout) but not reverse inequality</p> <p><math>n &gt; \frac{\log 0.1}{\log 0.79}</math>, so <math>n &gt; 9.768\dots</math></p> <p>Minimum <math>n = 10</math> Accept <math>n \geq 10</math></p> <hr/> <p><b>OR</b> (using trial and improvement):        Trial with <math>0.79^9</math> or <math>0.79^{10}</math></p> <p><math>1 - 0.79^9 = 0.8801</math> (<math>&lt; 0.9</math>) or <math>0.79^9 = 0.1198</math> (<math>&gt; 0.1</math>)</p> <p><math>1 - 0.79^{10} = 0.9053</math> (<math>&gt; 0.9</math>) or <math>0.79^{10} = 0.09468</math> (<math>&lt; 0.1</math>)</p> <p>Minimum <math>n = 10</math> Accept <math>n \geq 10</math></p> <hr/> <p>NOTE: <math>n = 10</math> unsupported scores SC1 only</p>	<p>M1 for <math>0.79^5</math> or <math>0.3077\dots</math>        M1 for <math>1 - 0.79^5</math> dep        A1 CAO</p> <p>M1 for equation/inequality in <math>n</math>        (accept either statement opposite)</p> <p>M1(indep) for process of using logs i.e. <math>\frac{\log a}{\log b}</math></p> <p>A1 CAO</p> <hr/> <p>M1(indep) for sight of <math>0.8801</math> or <math>0.1198</math></p> <p>M1( indep) for sight of <math>0.9053</math> or <math>0.09468</math></p> <p>A1 dep on both M's cao</p> <hr/>	<p>3</p> <p>3</p>	<p>16</p>
	<b>TOTAL</b>	<b>16</b>	



(v)	<p>Any two suitable comments such as:</p> <p>Outer London has a greater proportion (or %) of people under 20 (or almost equal proportion)</p> <p>The modal group in Inner London is 20-30 but in Outer London it is 30-40</p> <p>Outer London has a greater proportion (14%) of aged 65+</p> <p><b>All</b> populations in <b>each</b> age group are higher in Outer London</p> <p>Outer London has a more evenly spread distribution or balanced distribution (ages) o.e.</p>	<p>E1</p> <p>E1</p>	<p>2</p>
(vi)	<p>Mean increase ↑</p> <p>median unchanged (-)</p> <p>midrange increase ↑</p> <p>standard deviation increase ↑</p> <p>interquartile range unchanged. (-)</p>	<p>Any one correct B1</p> <p>Any two correct B2</p> <p>Any three correct B3</p> <p>All <b>five</b> correct B4</p>	<p>4</p>
		<b>TOTAL</b>	<b>20</b>

## 4767 Statistics 2

### Question 1

(i)	<p><b>EITHER:</b></p> $S_{xy} = \sum xy - \frac{1}{n} \sum x \sum y = 880.1 - \frac{1}{48} \times 781.3 \times 57.8$ $= -60.72$ $S_{xx} = \sum x^2 - \frac{1}{n} (\sum x)^2 = 14055 - \frac{1}{48} \times 781.3^2 = 1337.7$ $S_{yy} = \sum y^2 - \frac{1}{n} (\sum y)^2 = 106.3 - \frac{1}{48} \times 57.8^2 = 36.70$ $r = \frac{S_{xy}}{\sqrt{S_{xx} S_{yy}}} = \frac{-60.72}{\sqrt{1337.7 \times 36.70}} = -0.274$ <p><b>OR:</b></p> $\text{cov}(x,y) = \frac{\sum xy}{n} - \bar{x}\bar{y} = 880.1/48 - 16.28 \times 1.204$ $= -1.265$ $\text{rmsd}(x) = \sqrt{\frac{S_{xx}}{n}} = \sqrt{(1337.7/48)} = \sqrt{27.87} = 5.279$ $\text{rmsd}(y) = \sqrt{\frac{S_{yy}}{n}} = \sqrt{(36.70/48)} = \sqrt{0.7646} = 0.8744$ $r = \frac{\text{cov}(x,y)}{\text{rmsd}(x)\text{rmsd}(y)} = \frac{-1.265}{5.279 \times 0.8744} = -0.274$	<p>M1 for method for <math>S_{xy}</math></p> <p>M1 for method for at least one of <math>S_{xx}</math> or <math>S_{yy}</math></p> <p>A1 for at least one of <math>S_{xy}</math>, <math>S_{xx}</math>, <math>S_{yy}</math>. correct</p> <p>M1 for structure of <math>r</math> A1 CAO (-0.27 to -0.28)</p> <p>M1 for method for cov (x,y)</p> <p>M1 for method for at least one msd A1 for at least one of cov/msd correct M1 for structure of <math>r</math> A1 CAO (-0.27 to -0.28)</p>	5
(ii)	<p><math>H_0: \rho = 0</math> <math>H_1: \rho &lt; 0</math> (one-tailed test)</p> <p>where <math>\rho</math> is the population correlation coefficient</p> <p>For <math>n = 48</math>, 5% critical value = 0.2403</p> <p>Since <math> -0.274  &gt; 0.2403</math> we can reject <math>H_0</math>:</p> <p>There is sufficient evidence at the 5% level to suggest that there is negative correlation between education spending and population growth.</p>	<p>B1 for <math>H_0, H_1</math> in symbols</p> <p>B1 for defining <math>\rho</math></p> <p>B1FT for critical value</p> <p>M1 for sensible comparison leading to a conclusion A1 for result (FT <math>r &lt; 0</math>) E1 FT for conclusion in words</p>	6
(iii)	<p>Underlying distribution must be bivariate Normal. If the distribution is bivariate Normal then the scatter diagram will have an elliptical shape.</p>	<p>B1 CAO for bivariate Normal B1 indep for elliptical shape</p>	2
(iv)	<ul style="list-style-type: none"> <li>Correlation does not imply causation</li> <li>There could be a third factor</li> <li>increased growth could cause lower spending.</li> </ul> <p>Allow any sensible alternatives, including example of a possible third factor.</p>	<p>E1 E1 E1</p>	3
(v)	<p>Advantage – less effort or cost Disadvantage – the test is less sensitive (ie is less likely to detect any correlation which may exist)</p>	<p>E1 E1</p>	2
			18

## Question 2

(i)	<p>(A) <math>P(X = 2) = e^{-0.37} \frac{0.37^2}{2!} = 0.0473</math></p> <p>(B) <math>P(X &gt; 2)</math></p> $= 1 - \left( e^{-0.37} \frac{0.37^2}{2!} + e^{-0.37} \frac{0.37^1}{1!} + e^{-0.37} \frac{0.37^0}{0!} \right)$ $= 1 - (0.0473 + 0.2556 + 0.6907) = 0.0064$	<p>M1 A1 (2 s.f.)</p> <p>M1 for <math>P(X = 1)</math> and <math>P(X = 0)</math> M1 for complete method A1 <b>NB Answer given</b></p>	<b>5</b>
(ii)	<p>P(At most one day more than 2)</p> $= \binom{30}{1} \times 0.9936^{29} \times 0.0064 + 0.9936^{30} =$ $= 0.1594 + 0.8248 = 0.9842$	<p>M1 for coefficient M1 for <math>0.9936^{29} \times 0.0064</math> M1 for <math>0.9936^{30}</math> A1 CAO (min 2sf)</p>	<b>4</b>
(iii)	<p><math>\lambda = 0.37 \times 10 = 3.7</math></p> <p><math>P(X &gt; 8) = 1 - 0.9863</math></p> <p><math>= 0.0137</math></p>	<p>B1 for mean (SOI) M1 for probability A1 CAO</p>	<b>3</b>
(iv)	<p>Mean no. per 1000ml = <math>200 \times 0.37 = 74</math></p> <p>Using Normal approx. to the Poisson, <math>X \sim N(74, 74)</math></p> $P(X > 90) = P\left(Z > \frac{90.5 - 74}{\sqrt{74}}\right)$ $= P(Z > 1.918) = 1 - \Phi(1.918)$ $= 1 - 0.9724 = 0.0276$	<p>B1 for Normal approx. with correct parameters (SOI)</p> <p>B1 for continuity corr.</p> <p>M1 for probability using correct tail A1 CAO (min 2 s.f.), (but FT wrong or omitted CC)</p>	<b>4</b>
(v)	<p><math>P(\text{questionable}) = 0.0064 \times 0.0137 \times 0.0276</math></p> $= 2.42 \times 10^{-6}$	<p>M1 A1 CAO</p>	<b>2</b>
			<b>18</b>

## Question 3

<b>(i)</b>	$X \sim N(27500, 4000^2)$ $P(X > 25000) = P\left(Z > \frac{25000 - 27500}{4000}\right)$ $= P(Z > -0.625)$ $= \Phi(0.625) = 0.7340 \text{ (3 s.f.)}$	M1 for standardising  A1 for -0.625 M1 <i>dep</i> for correct tail A1CAO (must include use of differences)	<b>4</b>
<b>(ii)</b>	$P(7 \text{ of } 10 \text{ last more than } 25000)$ $= \binom{10}{7} \times 0.7340^7 \times 0.2660^3 = 0.2592$	M1 for coefficient M1 for $0.7340^7 \times 0.2660^3$ A1 FT (min 2sf)	<b>3</b>
<b>(iii)</b>	From tables $\Phi^{-1}(0.99) = 2.326$ $\frac{k - 27500}{4000} = -2.326$ $x = 27500 - 2.326 \times 4000 = 18200$	B1 for 2.326 seen M1 for equation in $k$ and negative $z$ -value  A1 CAO for awrt 18200	<b>3</b>
<b>(iv)</b>	$H_0: \mu = 27500; \quad H_1: \mu > 27500$ Where $\mu$ denotes the mean lifetime of the new tyres.	B1 for use of 27500 B1 for both correct B1 for definition of $\mu$	<b>3</b>
<b>(v)</b>	$\text{Test statistic} = \frac{28630 - 27500}{4000/\sqrt{15}} = \frac{1130}{1032.8}$ $= 1.094$ <p>5% level 1 tailed critical value of <math>z = 1.645</math>  <math>1.094 &lt; 1.645</math> so not significant.            There is not sufficient evidence to reject <math>H_0</math></p> <p>There is insufficient evidence to conclude that the new tyres last longer.</p>	M1 must include $\sqrt{15}$  A1 FT  B1 for 1.645 M1 <i>dep</i> for a sensible comparison leading to a conclusion  A1 for conclusion in words in context	<b>5</b>
			<b>18</b>



## Question 4

(i)	$H_0$ : no association between location and species. $H_1$ : some association between location and species.	B1 for both	1
(ii)	Expected frequency = $38/160 \times 42 = 9.975$  Contribution = $(3 - 9.975)^2 / 9.975$ = 4.8773	M1 A1 M1 for valid attempt at $(O-E)^2/E$ A1 <b>NB Answer given</b>	4
(iii)	Refer to $\chi^2_4$ Critical value at 5% level = 9.488  Test statistic $X^2 = 32.85$  Result is significant  There appears to be some association between location and species  NB if $H_0$ $H_1$ reversed, or 'correlation' mentioned, do not award first B1 or final E1	B1 for 4 deg of f (seen) B1 CAO for cv  M1 Sensible comparison, using 32.85, leading to a conclusion A1 for correct conclusion (FT their c.v.)  E1 conclusion in context	5
(iv)	<ul style="list-style-type: none"> <li>Limpets appear to be distributed as expected throughout all locations.</li> <li>Mussels are much more frequent in exposed locations and much less in pools than expected.</li> <li>Other shellfish are less frequent in exposed locations and more frequent in pools than expected.</li> </ul>	E1  E1, E1  E1, E1	5
(v)	$\frac{24}{53} \times \frac{32}{65} \times \frac{16}{42} = 0.0849$	M1 for one fraction M1 for product of all 3 A1 CAO	3
			<b>18</b>

## 4768 Statistics 3

Q1	$f(x) = k(20 - x) \quad 0 \leq x \leq 20$			
(a) (i)	$\int_0^{20} k(20 - x)dx = \left[ k \left( 20x - \frac{x^2}{2} \right) \right]_0^{20} = k \times 200 = 1$ $\therefore k = \frac{1}{200}$ <p>Straight line graph with negative gradient, in the first quadrant. Intercept correctly labelled (20, 0), with nothing extending beyond these points.</p> <p>Sarah is more likely to have only a short time to wait for the bus.</p>	M1 A1 G1 G1 E1	Integral of $f(x)$ , including limits (which may appear later), set equal to 1. Accept a geometrical approach using the area of a triangle. C.a.o.	5
(ii)	<p>Cdf <math>F(x) = \int_0^x f(t)dt</math></p> $= \frac{1}{200} \left( 20x - \frac{x^2}{2} \right)$ $= \frac{x}{10} - \frac{x^2}{400}$ <p><math>P(X &gt; 10) = 1 - F(10)</math> <math>= 1 - (1 - \frac{1}{4}) = \frac{1}{4}</math></p>	M1 A1 M1 A1	<p>Definition of cdf, including limits (or use of "+c" and attempt to evaluate it), possibly implied later. Some valid method must be seen.</p> <p>Or equivalent expression; condone absence of domain <math>[0, 20]</math>.</p> <p>Correct use of c's cdf. f.t. c's cdf. Accept geometrical method, e.g area = <math>\frac{1}{2}(20 - 10)f(10)</math>, or similarity.</p>	4
(iii)	<p>Median time, <math>m</math>, is given by <math>F(m) = \frac{1}{2}</math>.</p> $\therefore \frac{m}{10} - \frac{m^2}{400} = \frac{1}{2}$ $\therefore m^2 - 40m + 200 = 0$ $\therefore m = 5.86$	M1 M1 A1	<p>Definition of median used, leading to the formation of a quadratic equation.</p> <p>Rearrange and attempt to solve the quadratic equation.</p> <p>Other solution is 34.14; no explicit reference to/rejection of it is required.</p>	3

(b) (i)	A simple random sample is one where every sample of the required size has an equal chance of being chosen.	E2	S.C. Allow E1 for "Every member of the population has an equal chance of being chosen independently of every other member".	2
(ii)	Identify clusters which are capable of representing the population as a whole. Choose a random sample of clusters. Randomly sample or enumerate within the chosen clusters.	E1 E1 E1		3
(iii)	A random sample of the school population might involve having to interview single or small numbers of pupils from a large number of schools across the entire country. Therefore it would be more practical to use a cluster sample.	E1 E1	For "practical" accept e.g. convenient / efficient / economical.	2
				19

Q2	$A \sim N(100, \sigma = 1.9)$ $B \sim N(50, \sigma = 1.3)$		When a candidate's answers suggest that (s)he appears to have neglected to use the difference columns of the Normal distribution tables penalise the first occurrence only.	
(i)	$P(A < 103) = P\left(Z < \frac{103 - 100}{1.9} = 1.5789\right)$ $= 0.9429$	M1 A1 A1	For standardising. Award once, here or elsewhere. c.a.o.	3
(ii)	$A_1 + A_2 + A_3 \sim N(300,$ $\sigma^2 = 1.9^2 + 1.9^2 + 1.9^2 = 10.83)$ $P(\text{this} > 306) =$ $P\left(Z > \frac{306 - 300}{3 \cdot 291} = 1.823\right) = 1 - 0.9658 = 0.0342$	B1 B1 A1	Mean. Variance. Accept sd (= 3.291). c.a.o.	3
(iii)	$A + B \sim N(150,$ $\sigma^2 = 1.9^2 + 1.3^2 = 5.3)$ $P(\text{this} > 147) = P\left(Z > \frac{147 - 150}{2 \cdot 302} = -1.303\right)$ $= 0.9037$	B1 B1 A1	Mean. Variance. Accept sd (= 2.302). c.a.o.	3
(iv)	$B_1 + B_2 - A \sim N(0,$ $1.3^2 + 1.3^2 + 1.9^2 = 6.99)$ $P(-3 < \text{this} < 3)$ $= P\left(\frac{-3 - 0}{2.644} < Z < \frac{3 - 0}{2.644}\right) = P(-1.135 < Z < 1.135)$ $= 2 \times 0.8718 - 1 = 0.7436$	B1 B1 M1 A1 A1	Mean. Or $A - (B_1 + B_2)$ . Variance. Accept sd (= 2.644). Formulation of requirement ... ... two sided. c.a.o.	5
(v)	Given $\bar{x} = 302.3$ $s_{n-1} = 3.7$ CI is given by $302.3 \pm 1.96 \times \frac{3.7}{\sqrt{100}}$ $= 302.3 \pm 0.7252 = (301.57(48),$ $303.02(52))$ The batch appears not to be as specified since 300 is outside the confidence interval.	M1 B1 A1 E1	Correct use of 302.3 and $3.7/\sqrt{100}$ . For 1.96 c.a.o. Must be expressed as an interval.	4
				18

Q3												
(a)	$H_0: \mu_D = 0$ (or $\mu_I = \mu_{II}$ ) (i) $H_1: \mu_D \neq 0$ (or $\mu_{II} \neq \mu_I$ ) where $\mu_D$ is "mean for II – mean for I"  Normality of <u>differences</u> is required.	B1  B1  B1	Both. Hypotheses in words only must include "population". For adequate verbal definition. Allow absence of "population" if correct notation $\mu$ is used, but do NOT allow " $\bar{X}_I = \bar{X}_{II}$ " or similar unless $\bar{X}$ is clearly and explicitly stated to be a <u>population</u> mean.	3								
(ii)	<p><b>MUST</b> be PAIRED COMPARISON <math>t</math> test.          Differences are:</p> <table border="1" data-bbox="405 719 1195 757"> <tr> <td>10.0</td> <td>26.8</td> <td>42.7</td> <td>2.4</td> <td>-14.9</td> <td>-2.0</td> <td>16.3</td> <td>11.5</td> </tr> </table> <p><math>\bar{d} = 11.6</math>      <math>s_{n-1} = 17.707</math></p> <p>Test statistic is <math>\frac{11.6 - 0}{\frac{17.707}{\sqrt{8}}}</math></p> <p style="text-align: center;">= 1.852(92).</p> <p>Refer to <math>t_7</math>.          Double-tailed 5% point is 2.365.          Not significant.          Seems there is no difference between the mean yields of the two types of plant.</p>	10.0	26.8	42.7	2.4	-14.9	-2.0	16.3	11.5	B1  M1  A1  M1 A1 A1 A1	<p><math>s_n = 16.563</math> but do <b>NOT</b> allow this here or in construction of test statistic, but FT from there.</p> <p>Allow c's <math>\bar{d}</math> and/or <math>s_{n-1}</math>.          Allow alternative: <math>0 + (c's\ 2.365) \times \frac{17.707}{\sqrt{8}}</math> (= 14.806) for subsequent comparison with <math>\bar{d}</math>.          (Or <math>\bar{d} - (c's\ 2.365) \times \frac{17.707}{\sqrt{8}}</math> (= -3.206) for comparison with 0.)          c.a.o. but ft from here in any case if wrong.          Use of <math>0 - \bar{d}</math> scores M1A0, but ft.</p> <p>No ft from here if wrong.          No ft from here if wrong.          ft only c's test statistic.          ft only c's test statistic.          Special case: (<math>t_8</math> and 2.306) can score 1 of these last 2 marks if either form of conclusion is given.</p>	7
10.0	26.8	42.7	2.4	-14.9	-2.0	16.3	11.5					

(b)	Diff	-5	4	-14	-3	6	1	-11	-8	-7	-9	
	Rank of  diff	4	3	10	2	5	1	9	7	6	8	
	<p><math>W_+ = 1 + 3 + 5 = 9</math> (or <math>W_- = 2 + 4 + 6 + 7 + 8 + 9 + 10 = 46</math>)</p> <p>Refer to tables of Wilcoxon paired (/single sample) statistic for <math>n = 10</math>. Lower (or upper if 46 used) double-tailed 5% point is 8 (or 47 if 46 used). Result is not significant. No evidence to suggest the tasters differ on the whole.</p>	M1	For differences. ZERO in this section if differences not used.	M1	For ranks.	A1	FT from here if ranks wrong	B1				
		M1	No ft from here if wrong.	A1	i.e. a 2-tail test. No ft from here if wrong.	A1	ft only c's test statistic.	A1	ft only c's test statistic.			8
												18

Q4																																		
(a)	$\bar{x} = \frac{310}{100} = 3.1$ $s^2 = \frac{1288 - 100 \times 3.1^2}{99} = \frac{327}{99} = 3.303$ <p>Evidence could support Poisson since the variance is fairly close to the mean.</p>	B1 B1 E1	3																															
(ii)	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td><math>f_o</math></td> <td>6</td> <td>16</td> <td>19</td> <td>18</td> <td>17</td> <td>14</td> <td>6</td> <td>4</td> <td>0</td> </tr> <tr> <td><math>f_e</math></td> <td>4.50</td> <td>13.97</td> <td>21.65</td> <td>22.37</td> <td>17.33</td> <td>10.75</td> <td>5.55</td> <td>2.46</td> <td>1.42</td> </tr> <tr> <td>Merged</td> <td colspan="2">22 18.47</td> <td></td> <td></td> <td></td> <td></td> <td colspan="2">10 9.43</td> <td></td> </tr> </table> <p> <math display="block">X^2 = 0.6747 + 0.3244 + 0.8537 + 0.0063 + 0.9826 + 0.0345 = 2.876(2)</math> </p> <p>Refer to <math>\chi_4^2</math>. e.g. Upper 10% point is 7.779.</p> <p>Not significant. Suggests Poisson model does fit ... ... at any reasonable level of significance.</p>	$f_o$	6	16	19	18	17	14	6	4	0	$f_e$	4.50	13.97	21.65	22.37	17.33	10.75	5.55	2.46	1.42	Merged	22 18.47						10 9.43			M1 A1 A1  M1  M1 A1  M1  A1 A1 A1	<p>Calculation of expected frequencies. Last cell correct. All others correct, but ft if wrong.</p> <p>Combining cells. (Condone if not combined as fully as shown above, but require top two cells combined as a minimum.)</p> <p>Calculation of <math>X^2</math>.</p> <p>(Condone wrong last cell.) Depends on both of the preceding M marks.</p> <p>Allow correct df (= cells – 2) from wrongly grouped or ungrouped table, and FT. Otherwise, no FT if wrong.</p> <p>ft only c's test statistic. ft only c's test statistic. Or other sensible comment.</p>	10
$f_o$	6	16	19	18	17	14	6	4	0																									
$f_e$	4.50	13.97	21.65	22.37	17.33	10.75	5.55	2.46	1.42																									
Merged	22 18.47						10 9.43																											
(b)	<p>CI is given by</p> $1.465 \pm 2.262 \times \frac{0.3288}{\sqrt{10}}$ $= 1.465 \pm 0.2352 = (1.2298, 1.7002)$	M1 B1 B1  A1	<p>If <u>both</u> 1.465 and <math>0.3288/\sqrt{10}</math> are correct.</p> <p><b>If <math>t_9</math> used.</b> 95% 2-tail point for c's <math>t</math> distribution (Independent of previous mark).</p> <p>c.a.o. Must be expressed as an interval.</p>	4																														
			17																															

## 4769 Statistics 4

Q1				
(i)	$L = \frac{e^{-\theta} \theta^{x_1}}{x_1!} \dots \frac{e^{-\theta} \theta^{x_n}}{x_n!} \left[ = \frac{e^{-n\theta} \theta^{\sum x_i}}{x_1! x_2! \dots x_n!} \right]$ <p> <math>\ln L = \text{const} - n\theta + \sum x_i \ln \theta</math> </p> $\frac{d \ln L}{d\theta} = -n + \frac{\sum x_i}{\theta} = 0$ $\Rightarrow \hat{\theta} = \frac{\sum x_i}{n} (= \bar{x})$ <p>Check this is a maximum</p> <p>e.g. <math>\frac{d^2 \ln L}{d\theta^2} = -\frac{\sum x_i}{\theta^2} &lt; 0</math></p>	<p>M1 product form</p> <p>A1 fully correct</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1 CAO</p> <p>M1</p> <p>A1</p>		9
(ii)	$\lambda = P(X=0) = e^{-\theta}$	B1		1
(iii)	<p>We have <math>R \sim B(n, e^{-\theta})</math>,</p> <p>so <math>E(R) = ne^{-\theta}</math></p> <p><math>\text{Var}(R) = ne^{-\theta}(1 - e^{-\theta})</math></p> $\tilde{\lambda} = \frac{R}{n}$ <p><math>\therefore E(\tilde{\lambda}) = e^{-\theta}</math></p> <p>i.e. unbiased</p> $\text{Var}(\tilde{\lambda}) = \frac{e^{-\theta}(1 - e^{-\theta})}{n}$	<p>M1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p>	BEWARE PRINTED ANSWER	7



(iv)	<p>Relative efficiency of <math>\tilde{\lambda}</math> wrt ML est</p> $= \frac{\text{Var(ML Est)}}{\text{Var}(\tilde{\lambda})}$ $= \frac{\theta e^{-2\theta}}{n} \cdot \frac{n}{e^{-\theta}(1-e^{-\theta})} = \frac{\theta}{e^{\theta}-1}$ <p>Eg:- Expression is <math>\frac{\theta}{\theta + \frac{\theta^2}{2!} + \dots}</math></p> <p>always <math>&lt; 1</math></p> <p>and this is <math>\approx 1</math> if <math>\theta</math> is small  <math>\approx 0</math> if <math>\theta</math> is large</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>E1</p> <p>E1</p> <p>E1</p>	<p>any attempt to compare variances</p> <p>if correct</p> <p><b>BEWARE PRINTED ANSWER</b></p> <p>Allow statement that <math>\frac{\theta}{e^{\theta}-1} \rightarrow 0</math> as <math>\theta \rightarrow \infty</math></p>	7
------	--	---	--	---

Q2			
(i)	$P(X = x) = q^{x-1} p$ $\text{Pgf } G(t) = E(t^X) = \sum_{x=1}^{\infty} p t^x q^{x-1}$ $= pt(1 + qt + q^2 t^2 + \dots)$ $= \underline{\underline{pt(1 - qt)^{-1}}}$ $\mu = G'(1) \quad \sigma^2 = G''(1) + \mu - \mu^2$ $G'(t) = pt(-1)(1 - qt)^{-2}(-q) + p(1 - qt)^{-1}$ $= pqt(1 - qt)^{-2} + p(1 - qt)^{-1}$ $\therefore G'(1) = pq(1 - q)^{-2} + p(1 - q)^{-1} = \frac{q}{p} + 1 = \underline{\underline{\frac{1}{p}}}$ $G''(t) = pqt(-2)(1 - qt)^{-3}(-q) + pq(1 - qt)^{-2} + p(-1)(1 - qt)^{-2}(-q)$ $\therefore G''(1) = 2pq^2(1 - q)^{-3} + pq(1 - q)^{-2} + pq(1 - q)^{-2}$ $= \frac{2q^2}{p^2} + \frac{2q}{p}$ $\therefore \sigma^2 = \frac{2q^2}{p^2} + \frac{2q}{p} + \frac{1}{p} - \frac{1}{p^2} = \frac{2q^2 + 2pq + p - 1}{p^2}$ $= \frac{q}{p^2}(2q + 2p - 1) = \underline{\underline{\frac{q}{p^2}}}$	B1 M1 A1 A1 M1 A1 A1 A1 M1 A1	FT into pgf only  BEWARE PRINTED ANSWER [consideration of $ qt  < 1$ not required] for attempt to find $G'(t)$ and/or $G''(t)$  BEWARE PRINTED ANSWER  For inserting their values  BEWARE PRINTED ANSWER
			11

(ii)	$X_1 = \text{number of trials to first success}$ $X_2 = \text{ " " " " " next "}$ $\cdot$ $\cdot$ $\cdot$ $X_n = \text{ " " " " " nth "}$	$\therefore Y = X_1 + X_2 + \dots + X_n$ $= \text{total no of trials to the } n\text{th success}$	E1 E1		
	$\therefore \text{pgf of } Y = (\text{pgf of } X)^n = \underline{\underline{p^n t^n (1-qt)^{-n}}}$		1		
	$\underline{\underline{\mu_Y = n\mu_X = \frac{n}{p}}}$		1		
	$\underline{\underline{\sigma_Y^2 = n\sigma_X^2 = \frac{nq}{p^2}}}$		1		5
(iii)	N(candidate's $\mu_Y$ , candidate's $\sigma_Y^2$ )		1		1
(iv)	$Y = \text{no of tickets to be sold} \sim \text{random variable as in (ii) with } n = 140 \text{ and } p = 0.8$ $\sim \text{Approx } N\left(\frac{140}{0.8} = 175, \frac{140 \times 0.2}{(0.8)^2} = 43.75\right)$ $P(Y \geq 160) \approx P(N(175, 43.75) > 159 \frac{1}{2})$ $= P(N(0, 1) > -2.343)$ $= 0.9905$ For any sensible discussion <u>in context</u> (eg groups of passengers $\Rightarrow$ not indep.)		E1 1 M1 A1 A1 E1 E1	Do not award if cty corr absent or wrong, but FT if 160 used $\rightarrow$ -2.268, 0.9884  CAO	7
Q3	$X = \text{amount of salt} \sim N(\mu [750], \sigma^2 [20^2])$ Sample of $n=9$				
(i)	Type I error: rejecting null hypothesis ... ... when it is true.  Type II error: accepting null hypothesis ... ... when it is false.  OC: P (accepting null hypothesis ... ... as a function of the parameter under investigation)		B1 B1  B1 B1  B1 B1	Allow B1 for P(rej $H_0$ when true)  Allow B1 for P(acc $H_0$ when false)  [ P(type II error   the true value of the parameter) scores B1+B1]	6
(ii)	Reject if $\bar{x} < 735$ or $\bar{x} > 765$ $\alpha = P(\bar{X} < 735 \text{ or } \bar{X} > 765   \bar{X} \sim N(750, \frac{20^2}{9}))$ $= P(Z < \frac{(735-750)3}{20} = -2.25$ or $Z > \frac{(765-750)3}{20} = 2.25)$  $= 2(1-0.9878) = 2 \times 0.0122 = 0.0244$  This is the probability of rejecting good output and unnecessarily re-calibrating the machine – seems small [but not very small?]		M1  A1 A1  A1  E1 E1	Might be implicit    CAO  Accept any sensible comments	6


(iii)	Accept if $735 < \bar{x} < 765$ , and now $\mu = 725$ . $\beta = P(735 < \bar{X} < 765 \mid \bar{X} \sim N(725, 20^2/9))$ $= P(1.5 < Z < 6)$ $= 1 - 0.9332 = \underline{\underline{0.0668}}$ This is the probability of accepting output and carrying on when in fact $\mu$ has slipped to 725 – small[-ish?]	M1  A1 A1 A1  E1 E1	might be implicit  CAO If upper limit 765 not considered, maximum 2 of these 4 marks. If $\Phi(6)$ not considered, maximum 3 out of 4. accept sensible comments	6
(iv)	$OC = P(735 < \bar{X} < 765 \mid \bar{X} \sim N(\mu, 20^2/9))$ $= \Phi\left(\frac{(765 - \mu)3}{20}\right) - \Phi\left(\frac{(735 - \mu)3}{20}\right)$ <p style="text-align: center;">" <math>\Phi - \Phi</math> "</p> $\mu=720: \Phi(6.75) - \Phi(2.25) = 1 - 0.9878 = 0.0122$ $730: 5.25 \quad 0.75 = 1 - 0.7734 = 0.2266$ $740: 3.75 \quad -0.75 = 1 - (1 - 0.7734) = 0.7734$ $750: \text{similarly or by write-down from part (ii)}$ [ FT ] : 0.9756 $760, 770, 780 \text{ by symmetry}$ [FT]: 0.7734, 0.2266, 0.0122	M1  M1 A1  1  1  1	both correct  if any two correct	6
Q4				
(i)	$x_{ij} = \mu + \alpha_i + e_{ij}$ $\mu = \text{population ...}$ .. grand mean for whole experiment $\alpha_i = \text{population ...}$ .. mean by which $i$ th treatment differs from $\mu$ $e_{ij}$ are experimental errors... ~ ind N $(0, \sigma^2)$	1 1 1 1 1 1 1 3	Allow "uncorrelated" 1 for ind N; 1 for 0; 1 for $\sigma^2$ .	9
(ii)	Totals are 240, 246, 254, 264, 196 each from sample of size 5 Grand total 936  "Correction factor" $CF = \frac{936^2}{20} = 43804.8$  Total SS = 44544 - CF = 739.2			

	<p>Between contractors SS = <math>\frac{240^2}{5} + \dots + \frac{196^2}{5} - CF = 44209.6 - CF = 404.8</math></p> <p>Residual SS ( by subtraction) = <math>739.2 - 404.8 = 334.4</math></p> <table border="1" data-bbox="225 627 898 1041"> <thead> <tr> <th>Source of Variation</th> <th>SS</th> <th>df</th> <th>MS</th> <th>MS ratio</th> </tr> </thead> <tbody> <tr> <td>Between Contractors</td> <td>404.8</td> <td>3</td> <td>134.93</td> <td>6.456</td> </tr> <tr> <td>Residual</td> <td>334.4</td> <td>16</td> <td>20.9</td> <td></td> </tr> <tr> <td>Total</td> <td>739.2</td> <td>19</td> <td></td> <td></td> </tr> </tbody> </table> <p>Refer to <math>F_{3,16}</math></p> <p>Upper 5% point is 3.24</p> <p>Significant</p> <p>Seems performances of contractors are not all the same</p>	Source of Variation	SS	df	MS	MS ratio	Between Contractors	404.8	3	134.93	6.456	Residual	334.4	16	20.9		Total	739.2	19			<p>M1 M1 A1</p> <p>M1 M1</p> <p>1 A1</p> <p>1</p> <p>1 1 1 1</p>	<p>For correct methods for any two, if each calculated SS is correct.</p> <p>CAO</p> <p>NO FT IF WRONG</p> <p>NO FT IF WRONG</p>	<p>12</p>
Source of Variation	SS	df	MS	MS ratio																				
Between Contractors	404.8	3	134.93	6.456																				
Residual	334.4	16	20.9																					
Total	739.2	19																						
(iii)	<p>Randomised blocks</p> <p>Description</p>	<p>B1 E1 E1</p>	<p>Take the subject areas as "blocks", ensure each contractor is used at least once in each block</p>	<p>3</p>																				

# 4771 Decision Mathematics 1

## Solutions

1.



Objective has maximum value of 24 at (20,4)

M1 A1 third line  
 B1 shading  
 B1 (0,20) and (22,0)  
 B1 (9,14)  
 B1 (20,4)

M1 A1 solution

or M1 A1 B1,  
 B1 scale (implied OK),  
 B1 profit line, B1 (20,4)  
 M1 A1 (20,4) A1 (24)

## 2.

(i)	<table border="1"> <thead> <tr> <th></th> <th>X</th> <th>Y</th> </tr> </thead> <tbody> <tr> <td>5, 14, 153, 6, 24, 2, 14, 15</td> <td>5, 14, 153</td> <td>5, 2</td> </tr> <tr> <td>5, 14, 6, 24, 14, 15</td> <td>5, 14, 24</td> <td>5</td> </tr> <tr> <td>14, 6, 14, 15,</td> <td>14, 15</td> <td>14, 6</td> </tr> <tr> <td>14, 14</td> <td></td> <td></td> </tr> </tbody> </table> <p>Answer = 14 Comparisons = 30</p>		X	Y	5, 14, 153, 6, 24, 2, 14, 15	5, 14, 153	5, 2	5, 14, 6, 24, 14, 15	5, 14, 24	5	14, 6, 14, 15,	14, 15	14, 6	14, 14			M1 A1 A1
	X	Y															
5, 14, 153, 6, 24, 2, 14, 15	5, 14, 153	5, 2															
5, 14, 6, 24, 14, 15	5, 14, 24	5															
14, 6, 14, 15,	14, 15	14, 6															
14, 14																	
(ii)	<table border="1"> <thead> <tr> <th></th> <th>X</th> <th>Y</th> </tr> </thead> <tbody> <tr> <td>5, 14, 153, 6, 24, 2, 14</td> <td>5, 14, 153</td> <td>5, 2</td> </tr> <tr> <td>5, 14, 6, 24, 14</td> <td>5, 14, 24</td> <td>5</td> </tr> <tr> <td>14, 6, 14</td> <td>14</td> <td>14, 6</td> </tr> <tr> <td>14</td> <td></td> <td></td> </tr> </tbody> </table> <p>Answer = 14 Comparisons = 24</p>		X	Y	5, 14, 153, 6, 24, 2, 14	5, 14, 153	5, 2	5, 14, 6, 24, 14	5, 14, 24	5	14, 6, 14	14	14, 6	14			M1 A1 A1
	X	Y															
5, 14, 153, 6, 24, 2, 14	5, 14, 153	5, 2															
5, 14, 6, 24, 14	5, 14, 24	5															
14, 6, 14	14	14, 6															
14																	
(iii)	Median	B1															
(iv)	Time taken approximately proportional to square of length of list (or twice length takes four times the time, or equivalent).	B1															

## 3.

(i)	$T_1 \rightarrow T_2$ $T_1 \rightarrow T_3 \rightarrow T_2$ $T_1 \rightarrow T_3$ $T_1 \rightarrow T_2 \rightarrow T_3$ $T_1 \rightarrow T_2 \rightarrow T_3 \rightarrow T_4$ $T_1 \rightarrow T_3 \rightarrow T_4$	M1 A1
(ii)	$T_4 \rightarrow T_3 \rightarrow T_2 \rightarrow T_1$ $T_4 \rightarrow T_3 \rightarrow T_1$ $T_4 \rightarrow T_3 \rightarrow T_1 \rightarrow T_2$ $T_4 \rightarrow T_3 \rightarrow T_2$ $T_4 \rightarrow T_3$	M1 A1
(iii)	22	M1   allow for 23 A1
(iv)	11	M1   halving (not 11.5) A1

4.

(i) e.g. 00–09→1  
 10–39→2  
 40–79→3  
 80–89→4  
 90–99→5

M1  
 A1 proportions OK  
 A1 efficient

(ii) e.g. 00–15→1  
 16–47→2  
 48–55→3  
 56–79→4  
 80–87→5  
 88–95→6  
 96, 97, 98, 99 reject

M1 some rejected  
 A2 proportions OK  
 (–1 each error)  
 A1 efficient

(iii) &amp; (iv)

Sim. no.	Cars arriving after Joe – time interval   number of passengers										Time to 15 passengers (minutes)
1	3	2	2	1	1	2	2	2	3	1	6
2	3	1	2	2	1	4	1	2	5	1	6
3	5	1	2	2	2	1	3	4	2	2	12
4	4	6	3	2	4	1	1	2	2	3	4
5	5	1	4	1	3	2	5	4	2	2	17
6	4	4	4	2	5	3	1	4	1	4	8
7	4	1	4	2	3	1	5	4	1	3	16
8	2	2	2	2	2	4	3	5	1	2	6
9	1	1	1	1	1	1	1	1	1	2	5
10	2	4	3	2	2	6	2	5	2	1	5

M1  
 A2 (–1 each error)

M1 simulation  
 A1 time intervals  
 A1 passengers  
 A1 time to wait

(v) 0.8  
 more runs

B1  
 B1




5.

(a)(i) Activity D.  
Depends on A and B in project 1, but on A, B and C in project 2.

(ii) Project 1: Duration is 5 for  $x < 3$ , thence  $x + 2$ .  
Project 2: Duration is 5 for  $x < 2$ , thence  $x + 3$

(b) (i) & (ii)



Project duration – 7  
Critical activities – T, X

- M1
- A1
- A1
- B1 "5"
- B1 B1 beyond 5
- M1 activity-on-arc
- A1 single start and single end
- A2 precedences (-1 each error)
- M1 A1 forward pass
- M1 A1 backward pass
- B1
- B1

6.


(i)

Order of inclusion	1	3	6	4	5	2
	A	B	C	D	E	F
A	-	10	7	-	9	5
B	10	-	-	1	-	4
C	7	-	-	-	3	-
D	-	1	-	-	2	-
E	9	-	3	2	-	-
F	5	4	-	-	-	-

Arcs: AF, FB, BD, DE, EC

Length: 15

(ii) & (iii)



Arcs: AF, FB, BD, AC, AE

Length: 26

(iv) Cubic

n applications of Dijkstra, which is quadratic

M1  
A1 select  
A1 delete  
A1 order

B1

B1

B1 arcs  
B1 lengths

M1 Dijkstra  
A1 working values  
A1 order of labelling  
A1 labels

M1  
A1

B1

B1


# 4772 Decision Mathematics 2

1.

(a)(i) "not fail" → "succeed"  
"disagree less" → "agree more"

(ii) e.g. "I don't entirely agree with you".

(b) e.g.



(c)

$(a \wedge b) \vee (\sim a \wedge \sim b)$	$\Leftrightarrow$	$((\sim a \vee b) \wedge (a \vee \sim b))$
1 1 1 1 0 0 0 1	0 1 1 1 1 1 0	0 1 1 1 1 1 0
1 0 0 0 0 0 1 1	0 0 0 0 0 0 1	0 0 0 0 1 1 1
0 0 1 0 1 0 0 1	1 1 1 1 0 0 0	0 0 0 0 0 0 0
0 0 0 1 1 1 1 1	1 1 1 1 1 1 1	1 1 0 1 0 1 1


B1  
B1  
M1 same meaning  
A1 simpler

M1 2 switches + light in a circuit  
A4 one for each correct setting

M1 4 lines  
A1 a's and b's  
A1 negations  
A1 level 1 and's  
A1 level 1 or's  
A1 level 2  
A1 result


2.

(i) e.g. (Decisions could be in other order.)



Drive down for 2 lots of 4 weeks

(ii)



Jane could save money if she spent less than £10000 on a car

(iii) EMV – expected monetary value – probabilistically weighted cash values  
 Utility measure is an alternative.

- M1  $4*2/2*4$
- M1 rent/drive
- A1
- M1 costs
- A1
- B1 advice
- B1 buy/don't buy
- M1 don't buy analysis
- A1 costings
- M1 chance node
- A1 buy analysis
- M1 buy costings
- A1
- B1
- B1
- B1

3.

(a) (i)

	1	2	3	4
1	$\infty$	14	11	24
2	14	$\infty$	15	$\infty$
3	11	15	$\infty$	12
4	24	$\infty$	12	$\infty$

	1	2	3	4
1	1	2	3	4
2	1	2	3	4
3	1	2	3	4
4	1	2	3	4

	1	2	3	4
1	$\infty$	14	11	24
2	14	28	15	38
3	11	15	22	12
4	24	38	12	48

	1	2	3	4
1	1	2	3	4
2	1	1	3	1
3	1	2	1	4
4	1	1	3	1

	1	2	3	4
1	28	14	11	24
2	14	28	15	38
3	11	15	22	12
4	24	38	12	48


	1	2	3	4
1	2	2	3	4
2	1	1	3	1
3	1	2	1	4
4	1	1	3	1

	1	2	3	4
1	22	14	11	23
2	14	28	15	27
3	11	15	22	12
4	23	27	12	24

	1	2	3	4
1	3	2	3	3
2	1	1	3	3
3	1	2	1	4
4	3	3	3	3

	1	2	3	4
1	22	14	11	23
2	14	28	15	27
3	11	15	22	12
4	23	27	12	24

	1	2	3	4
1	3	2	3	3
2	1	1	3	3
3	1	2	1	4
4	3	3	3	3



(ii) 1 3 4 2 1  
64  
⇒ 1 3 4 3 2 1

M1 sca Floyd  
A1 distance  
A1 route

A1

A1

A1

B1 loops  
B1 rest

M1 A1  
B1  
B1

(iii) $27 + 11 + 14 = 52$ TSP solution has length between 52 and 64	M1 A1 M1 A1
(b) e.g. <b>1 3 1 2 3 4 1</b> length = 87 One repeated arc $\rightarrow$ Eulerian	M1 A1 A1 B1

4.

(i) Let a be the number of tonnes of A produced ...

$$\begin{aligned} \text{Max} & \quad a+b+c \\ \text{st} & \quad 3a+2b+5c < 60 \\ & \quad 5a+6b+2c < 50 \end{aligned}$$

M1 A1

B1  
B1  
B1

(ii) e.g.

P	a	b	c	s <sub>1</sub>	s <sub>2</sub>	RHS
1	-1	-1	-1	0	0	0
0	3	2	5	1	0	60
0	5	6	2	0	1	50
1	-0.4	-0.6	0	0.2	0	12
0	0.6	0.4	1	0.2	0	12
0	3.8	5.2	0	-0.4	1	26
1	>0	0	0	>0	>0	15
0		0	1			10
0	19/26	1	0	-2/26	5/26	5

M1 A1 initial tableau

M1 A1 pivot

M1 A1

Make 5 tonnes of B and 10 tonnes of C

B1 interpretation

(iii) & (iv) e.g.

A	P	a	b	c	s <sub>1</sub>	s <sub>2</sub>	s <sub>3</sub>	art	RHS
1	0	1	0	0	0	0	-1	0	8
0	1	-1	-1	-1	0	0	0	0	0
0	0	3	2	5	1	0	0	0	60
0	0	5	6	2	0	1	0	0	50
0	0	1	0	0	0	0	-1	1	8
1	0	0	0	0	0	0	0	-1	0
0	1	0	-1	-1	0	0	-1	1	8
0	0	0	2	5	1	0	3	-3	36
0	0	0	6	2	0	1	5	-5	10
0	0	1	0	0	0	0	-1	1	8
1	0	0	0	0	0	0	0	-1	0
	1	0	2	0	0	0.5	1.5		13
	0	0	-13	0	1	-2.5	-4.5		11
	0	0	3	1	0	0.5	2.5		5
	0	1	0	0	0	0	-1		8

B1 new constraint  
M1 surplus + artificial  
A1  
B1 new objective

M1 A1

B1

Make 8 tonnes of A and 5 tonnes of C

B1 interpretation

# 4773 Decision Mathematics Computation

1.

(i) $XA + XB + XE + XF \geq 1$  Indicator variables correspond to matrix column A (or row A) entries which are less than or equal to 5. Ensures that at least one such indicator is 1.	M1 A1 ">" OK  B1 indicator vars B1 $\leq 5$ B1
(ii) Min $XA+XB+XC+XD+XE+XF$ st $XA+XB+XE+XF \geq 1$ $XA+XB+XE+XF \geq 1$ $XC+XF \geq 1$ $XD+XE \geq 1$ $XA+XB+XD+XE+XF \geq 1$ $XA+XB+XC+XE+XF \geq 1$	B1  M1 A3 (–1 each error/ omission) allow (correct) reduced set of inequalities
(iii) 2 centres, at F&D or E&C or E&F	M1 A1 A1
(iv) e.g. add $XF=0$ to force solution E and C (v) Three solutions are F & D, E & C, E & F.	M1 A1 B1
(vi) Problem is unimodular (or convincing argument). In the interests of efficiency (and parsimony).	B1 B1



2.

<p>(i) e.g. (candidates should show formulae)</p> <table style="border-collapse: collapse; width: 100%;"> <tr> <td style="padding-right: 10px;"><math>\alpha =</math></td> <td style="padding-right: 10px;">0.01</td> <td style="padding-right: 10px;">10</td> <td style="padding-right: 10px;">P(birth)</td> <td style="padding-right: 10px;">P(death)</td> <td style="padding-right: 10px;">rand</td> <td style="padding-right: 10px;">rand</td> <td style="padding-right: 10px;">birth</td> <td style="padding-right: 10px;">death</td> </tr> <tr> <td style="padding-right: 10px;"><math>\beta =</math></td> <td style="padding-right: 10px;">0.04</td> <td style="padding-right: 10px;">9</td> <td style="padding-right: 10px;">0.1</td> <td style="padding-right: 10px;">0.4</td> <td style="padding-right: 10px;">0.4261</td> <td style="padding-right: 10px;">0.3537</td> <td style="padding-right: 10px;">0</td> <td style="padding-right: 10px;">1</td> </tr> <tr> <td></td> <td></td> <td style="padding-right: 10px;">8</td> <td style="padding-right: 10px;">0.09</td> <td style="padding-right: 10px;">0.36</td> <td style="padding-right: 10px;">0.257</td> <td style="padding-right: 10px;">0.1405</td> <td style="padding-right: 10px;">0</td> <td style="padding-right: 10px;">1</td> </tr> <tr> <td></td> <td></td> <td style="padding-right: 10px;">8</td> <td style="padding-right: 10px;">0.08</td> <td style="padding-right: 10px;">0.32</td> <td style="padding-right: 10px;">0.8854</td> <td style="padding-right: 10px;">0.8632</td> <td style="padding-right: 10px;">0</td> <td style="padding-right: 10px;">0</td> </tr> </table>	$\alpha =$	0.01	10	P(birth)	P(death)	rand	rand	birth	death	$\beta =$	0.04	9	0.1	0.4	0.4261	0.3537	0	1			8	0.09	0.36	0.257	0.1405	0	1			8	0.08	0.32	0.8854	0.8632	0	0	<p>B1 handling parameters</p> <p>B1 births</p> <p>B1 deaths</p> <p>B1 use of "rand"</p> <p>B1 use of "if"</p> <p>B1 updating population</p>
$\alpha =$	0.01	10	P(birth)	P(death)	rand	rand	birth	death																													
$\beta =$	0.04	9	0.1	0.4	0.4261	0.3537	0	1																													
		8	0.09	0.36	0.257	0.1405	0	1																													
		8	0.08	0.32	0.8854	0.8632	0	0																													
<p>(ii) e.g. 1 1 3 2 0 2 2 3 3 0 – 0.2</p>	<p>M1 A1 B1</p>																																				
<p>(iii) e.g.</p> <table style="border-collapse: collapse; width: 100%;"> <tr> <td style="padding-right: 10px;"><math>\beta</math></td> <td style="padding-right: 10px;"></td> <td style="padding-right: 10px;">0.01</td> <td style="padding-right: 10px;">0.02</td> <td style="padding-right: 10px;">0.03</td> <td style="padding-right: 10px;">0.04</td> <td style="padding-right: 10px;">0.05</td> <td style="padding-right: 10px;">0.06</td> <td style="padding-right: 10px;">0.07</td> </tr> <tr> <td style="padding-right: 10px;">prob extinction</td> <td style="padding-right: 10px;">0</td> <td style="padding-right: 10px;">0</td> <td style="padding-right: 10px;">0.1</td> <td style="padding-right: 10px;">0.2</td> <td style="padding-right: 10px;">0.5</td> <td style="padding-right: 10px;">0.7</td> <td style="padding-right: 10px;">0.8</td> <td></td> </tr> </table>	$\beta$		0.01	0.02	0.03	0.04	0.05	0.06	0.07	prob extinction	0	0	0.1	0.2	0.5	0.7	0.8		<p>M1</p> <p>A1 decent range</p> <p>A1 reasonable outcomes</p>																		
$\beta$		0.01	0.02	0.03	0.04	0.05	0.06	0.07																													
prob extinction	0	0	0.1	0.2	0.5	0.7	0.8																														
<p>(iv) Addition of another rand + another if + extra add-on</p>	<p>B1</p> <p>M1 A1</p> <p>B1</p>																																				
<p>(v) e.g.</p> <table style="border-collapse: collapse; width: 100%;"> <tr> <td style="padding-right: 10px;"><math>\beta</math></td> <td style="padding-right: 10px;"></td> <td style="padding-right: 10px;">0.01</td> <td style="padding-right: 10px;">0.02</td> <td style="padding-right: 10px;">0.03</td> <td style="padding-right: 10px;">0.04</td> <td style="padding-right: 10px;">0.05</td> <td style="padding-right: 10px;">0.06</td> <td style="padding-right: 10px;">0.07</td> </tr> <tr> <td style="padding-right: 10px;">prob. extinction</td> <td style="padding-right: 10px;">0</td> <td style="padding-right: 10px;">0</td> <td style="padding-right: 10px;">0</td> <td style="padding-right: 10px;">0</td> <td style="padding-right: 10px;">0</td> <td style="padding-right: 10px;">0.2</td> <td style="padding-right: 10px;">0.3</td> <td></td> </tr> </table>	$\beta$		0.01	0.02	0.03	0.04	0.05	0.06	0.07	prob. extinction	0	0	0	0	0	0.2	0.3		<p>M1</p> <p>A1</p>																		
$\beta$		0.01	0.02	0.03	0.04	0.05	0.06	0.07																													
prob. extinction	0	0	0	0	0	0.2	0.3																														

3.

<p>(a) max AB+AE            st AB-BC-BD+CB+DB=0            BC+DC-CB-CD-CG=0            BD+ED+CD-DB-DC-DE=0            AE-ED-EF+DE+FE=0            EF-FE-FG=0            AB&lt;8            AE&lt;3            BC&lt;5            CB&lt;5            BD&lt;7            DB&lt;7            CD&lt;1            DC&lt;1            DE&lt;2            ED&lt;2            EF&lt;4            FE&lt;4            CG&lt;8            FG&lt;6</p> <p>END</p> <p>OBJECTIVE FUNCTION VALUE</p> <p>1) 10.00000</p> <table border="1"> <thead> <tr> <th>VARIABLE</th> <th>VALUE</th> <th>REDUCED COST</th> </tr> </thead> <tbody> <tr><td>AB</td><td>7.000000</td><td>0.000000</td></tr> <tr><td>AE</td><td>3.000000</td><td>0.000000</td></tr> <tr><td>BC</td><td>5.000000</td><td>0.000000</td></tr> <tr><td>BD</td><td>2.000000</td><td>0.000000</td></tr> <tr><td>CB</td><td>0.000000</td><td>1.000000</td></tr> <tr><td>DB</td><td>0.000000</td><td>0.000000</td></tr> <tr><td>DC</td><td>1.000000</td><td>0.000000</td></tr> <tr><td>CD</td><td>0.000000</td><td>1.000000</td></tr> <tr><td>CG</td><td>6.000000</td><td>0.000000</td></tr> <tr><td>ED</td><td>0.000000</td><td>0.000000</td></tr> <tr><td>DE</td><td>1.000000</td><td>0.000000</td></tr> <tr><td>EF</td><td>4.000000</td><td>0.000000</td></tr> <tr><td>FE</td><td>0.000000</td><td>1.000000</td></tr> <tr><td>FG</td><td>4.000000</td><td>0.000000</td></tr> </tbody> </table> <p>Max flow of 10 with flows of 7 from A to B, ... etc.</p>	VARIABLE	VALUE	REDUCED COST	AB	7.000000	0.000000	AE	3.000000	0.000000	BC	5.000000	0.000000	BD	2.000000	0.000000	CB	0.000000	1.000000	DB	0.000000	0.000000	DC	1.000000	0.000000	CD	0.000000	1.000000	CG	6.000000	0.000000	ED	0.000000	0.000000	DE	1.000000	0.000000	EF	4.000000	0.000000	FE	0.000000	1.000000	FG	4.000000	0.000000	<p>B1 objective</p> <p>M1 flow balance A2 constraints</p> <p>M1 capacity A1 constraints</p> <p>M1 run</p> <p>A1 results</p> <p>B1 interpretation</p>
VARIABLE	VALUE	REDUCED COST																																												
AB	7.000000	0.000000																																												
AE	3.000000	0.000000																																												
BC	5.000000	0.000000																																												
BD	2.000000	0.000000																																												
CB	0.000000	1.000000																																												
DB	0.000000	0.000000																																												
DC	1.000000	0.000000																																												
CD	0.000000	1.000000																																												
CG	6.000000	0.000000																																												
ED	0.000000	0.000000																																												
DE	1.000000	0.000000																																												
EF	4.000000	0.000000																																												
FE	0.000000	1.000000																																												
FG	4.000000	0.000000																																												
<p>(b)</p> <p>min <math>8x_{12}+3x_{15}+8x_{21}+5x_{23}+7x_{24}+5x_{32}+x_{34}+8x_{37}+7x_{42}+x_{43}+2x_{45}+3x_{51}+2x_{54}+4x_{56}+4x_{65}+6x_{67}+8x_{73}+6x_{76}</math></p> <p>st <math>x_{12}+x_{15}=10</math>  <math>x_{21}+x_{23}+x_{24}=10</math>  <math>x_{32}+x_{34}+x_{37}=10</math>  <math>x_{42}+x_{43}+x_{45}=10</math></p>	<p>M1 objective A1</p> <p>M1 supply A1 constraints</p>																																													

$x_{51} + x_{54} + x_{56} = 10$ $x_{65} + x_{67} = 10$ $x_{73} + x_{76} = 10$ $x_{21} + x_{51} = 10$ $x_{12} + x_{32} + x_{42} = 10$ $x_{23} + x_{43} + x_{73} = 10$ $x_{24} + x_{34} + x_{54} = 10$ $x_{15} + x_{45} + x_{65} = 10$ $x_{56} + x_{76} = 10$ $x_{37} + x_{67} = 10$ END	M1 demand A1 constraints																																																									
OBJECTIVE FUNCTION VALUE 1) 310.0000	M1 run																																																									
<table border="1"> <thead> <tr> <th>VARIABLE</th> <th>VALUE</th> <th>REDUCED COST</th> </tr> </thead> <tbody> <tr><td>X12</td><td>0.000000</td><td>0.000000</td></tr> <tr><td>X15</td><td>10.000000</td><td>0.000000</td></tr> <tr><td>X21</td><td>0.000000</td><td>0.000000</td></tr> <tr><td>X23</td><td>10.000000</td><td>0.000000</td></tr> <tr><td>X24</td><td>0.000000</td><td>0.000000</td></tr> <tr><td>X32</td><td>0.000000</td><td>0.000000</td></tr> <tr><td>X34</td><td>10.000000</td><td>0.000000</td></tr> <tr><td>X37</td><td>0.000000</td><td>6.000000</td></tr> <tr><td>X42</td><td>10.000000</td><td>0.000000</td></tr> <tr><td>X43</td><td>0.000000</td><td>0.000000</td></tr> <tr><td>X45</td><td>0.000000</td><td>0.000000</td></tr> <tr><td>X51</td><td>10.000000</td><td>0.000000</td></tr> <tr><td>X54</td><td>0.000000</td><td>0.000000</td></tr> <tr><td>X56</td><td>0.000000</td><td>6.000000</td></tr> <tr><td>X65</td><td>0.000000</td><td>0.000000</td></tr> <tr><td>X67</td><td>10.000000</td><td>0.000000</td></tr> <tr><td>X73</td><td>0.000000</td><td>0.000000</td></tr> <tr><td>X76</td><td>10.000000</td><td>0.000000</td></tr> </tbody> </table>	VARIABLE	VALUE	REDUCED COST	X12	0.000000	0.000000	X15	10.000000	0.000000	X21	0.000000	0.000000	X23	10.000000	0.000000	X24	0.000000	0.000000	X32	0.000000	0.000000	X34	10.000000	0.000000	X37	0.000000	6.000000	X42	10.000000	0.000000	X43	0.000000	0.000000	X45	0.000000	0.000000	X51	10.000000	0.000000	X54	0.000000	0.000000	X56	0.000000	6.000000	X65	0.000000	0.000000	X67	10.000000	0.000000	X73	0.000000	0.000000	X76	10.000000	0.000000	A1 results
VARIABLE	VALUE	REDUCED COST																																																								
X12	0.000000	0.000000																																																								
X15	10.000000	0.000000																																																								
X21	0.000000	0.000000																																																								
X23	10.000000	0.000000																																																								
X24	0.000000	0.000000																																																								
X32	0.000000	0.000000																																																								
X34	10.000000	0.000000																																																								
X37	0.000000	6.000000																																																								
X42	10.000000	0.000000																																																								
X43	0.000000	0.000000																																																								
X45	0.000000	0.000000																																																								
X51	10.000000	0.000000																																																								
X54	0.000000	0.000000																																																								
X56	0.000000	6.000000																																																								
X65	0.000000	0.000000																																																								
X67	10.000000	0.000000																																																								
X73	0.000000	0.000000																																																								
X76	10.000000	0.000000																																																								
Cost = 310 by sending 10 from W1 to S5, ... etc.	B1 interpretation																																																									

4.

<p>(a) Auxiliary equation:  <math>2\lambda^2 - 3\lambda + 1 = 0</math>  <math>(2\lambda - 1)(\lambda - 1) = 0</math>  <math>\lambda = 1</math> or <math>\frac{1}{2}</math></p> <p><math>u_n = A + B(\frac{1}{2})^n</math></p> <p><math>5 = A + B</math>  <math>3 = A + \frac{1}{2}B</math></p> <p><math>u_n = 1 + 4(\frac{1}{2})^n</math></p> <p><math>u_2 = 2, u_3 = 1.5, u_{10} = 1.003906</math>  <math>u_{1000000} \approx 1</math></p>	<p>M1 A1  M1  A1</p> <p>B1 B1</p> <p>B1  B1</p> <p>M1 A1</p> <p>B1 B1  B1</p>																																										
<p>(b)(i) &amp; (ii)</p> <table border="0"> <tbody> <tr><td>0</td><td>5</td></tr> <tr><td>1</td><td>3</td></tr> <tr><td>2</td><td>4.5</td></tr> <tr><td>3</td><td>8.75</td></tr> <tr><td>4</td><td>13.625</td></tr> <tr><td>5</td><td>16.6875</td></tr> <tr><td>6</td><td>16.40625</td></tr> <tr><td>7</td><td>12.92188</td></tr> <tr><td>8</td><td>7.976563</td></tr> <tr><td>9</td><td>4.042969</td></tr> <tr><td>10</td><td>3.087891</td></tr> <tr><td>11</td><td>5.588867</td></tr> <tr><td>12</td><td>10.29541</td></tr> <tr><td>13</td><td>14.85425</td></tr> <tr><td>14</td><td>16.98596</td></tr> <tr><td>15</td><td>15.62469</td></tr> <tr><td>16</td><td>11.45108</td></tr> <tr><td>17</td><td>6.551926</td></tr> <tr><td>18</td><td>3.376808</td></tr> <tr><td>19</td><td>3.513287</td></tr> <tr><td>20</td><td>6.893122</td></tr> </tbody> </table>	0	5	1	3	2	4.5	3	8.75	4	13.625	5	16.6875	6	16.40625	7	12.92188	8	7.976563	9	4.042969	10	3.087891	11	5.588867	12	10.29541	13	14.85425	14	16.98596	15	15.62469	16	11.45108	17	6.551926	18	3.376808	19	3.513287	20	6.893122	<p>M1  A1</p> <p>A1 3.087891</p> <p>A1 6.893122</p>
0	5																																										
1	3																																										
2	4.5																																										
3	8.75																																										
4	13.625																																										
5	16.6875																																										
6	16.40625																																										
7	12.92188																																										
8	7.976563																																										
9	4.042969																																										
10	3.087891																																										
11	5.588867																																										
12	10.29541																																										
13	14.85425																																										
14	16.98596																																										
15	15.62469																																										
16	11.45108																																										
17	6.551926																																										
18	3.376808																																										
19	3.513287																																										
20	6.893122																																										
<p>(iii) Limited wrt to (very) long-term</p>	<p>B1</p>																																										

## 4776 Numerical Methods

1	x	3	3.5	root = $(3 \times (-0.8) - 3.5 \times 0.5) / (-0.8 - 0.5)$	[M1A1A1]
	f(x)	0.5	-0.8	= .192308 (3.192, 3.19)	[A1]

(-) mpe is  $3.5 - 3.192308 = 0.307692$  (0.308, 0.31) [M1A1]

[TOTAL 6]

2	1	2				
	3	1	-1			
	5	5	4	5		
	7	k	k-5	k-9	k-14	
	9	2	2-k	7-2k	16-3k	[M1A1A1A1]

$16-3k = k-14$  hence  $k = 7.5$  [M1A1]

[TOTAL 6]

3	h	f(2+h)	f(2-h)	f'(2)		derivatives
	0.2	.494507	.867869	.566594		[M1A1A1A1]
	0.1	.323418	.010586	.564163	-0.00243	differences
	0.05	.241636	.085281	.563555	-0.00061	[M1A1]

differences reducing by a factor 4 so next estimate about 1.56340. [M1]

1.563 secure to 3 dp. [B1]

[TOTAL 8]

4	$f(x) = x^3 - 25$	$f'(x) = 3x^2$			[M1A1A1]
	$x_{r+1} = x_r - (x_r^3 - 25) / 3x_r^2$	(a.g.)			

	r	0	1	2	3	
	$x_r$	4	3.1875	.945197	2.92417	[M1A1]
	diffs		-0.8125	-0.2423	-0.02103	[B1]
	ratios			.298219	.086783	[B1]

differences reducing at an increasing rate (*hence faster than first order*) [E1]

[TOTAL 8]

5 (i)	0.001 369 352	(accept 0.001 369 4)	[B1]
-------	---------------	----------------------	------

(ii)	$\sin 86^\circ = 0.997$	$\sin 85^\circ = 0.996$	[B1B1]
	564	195	

$\sin 86^\circ - \sin 86^\circ = 0.001 369$  [A1]

(iii)	$2 \times 0.0784591 \times 0.008 726 54$	[M1]
	$= 0.00136935$	[A1]

(iv)	Rounding has different effects in the two expressions ( <i>may be implied</i> )	[E1]
	First method involves subtraction of nearly equal numbers and so loses accuracy	[E1]

[TOTAL 8]

4776

Mark Scheme

6 (i)	h	M	T		
	2	2.763547	<b>2.425240</b>		
	1	<b>2.677635</b>	<b>2.594393</b>	mid-point:	[M1A1A1]
	0.5	<b>2.656743</b>	<b>2.636014</b>	trapezium:	[M1A1A1A1] [subtotal 7]

(ii)	M:	2.763547	diffs	
		2.677635	-0.08591	
		2.656743	-0.02089	reducing by a factor 4 ( <i>may be implied</i> )
	Differences in T reduce by a factor 4, too			[B1] [subtotal 4]

(iii)		M	T	S		
		2.763547	2.425240	2.650778		[M1]
		2.677635	2.594393	2.649888	-0.00089033	S values: [A1A1]
		2.656743	2.636014	2.649833	-0.000054333	diffs [A1]
	Differences in S reducing fast e.g by a factor of (about) 16				[E1]	
	How this leads to an answer, e.g:					
	Next difference about -0.0000034 and/or next answer about 2.649830				[E1]	
	Accept 2.6498 or 2.64983				[A1] [subtotal 7]	
					[TOTAL 18]	

7 (i)	Eg: graph of $x^2$ and $4 + 1/x$ for $x > 0$ showing single intersection	[G2]
	Change of sign to find interval (2,3) - i.e. a = 2	[B1]

r	0	1	2	3	4	5	
$x_r$	2.5	2.097618	2.115829	2.114859	2.11491	2.114907	[M1A1A1]
	2.1149 secure to 4 dp						[A1] [subtotal 7]

(ii)	The iteration gives positive values only.	[E1]
------	---	------

r	0	1	2	3	4	5	
$x_r$	-2	-1.87083	-1.86158	-1.86087	-1.86081	-1.86081	[M1A1A1]
	-1.8608 secure to 4 dp						[A1] [subtotal 5]

4776

Mark Scheme

June 200



(iii)	Eg	r	0	1	2	3	4
		$x_r$	-0.5	-1.41421	-1.81463	-1.85713	-1.86052

not converging to required root (converging to previous root)

[M1A1]

Eg  $x_{r+1} = 1 / (x_r^2 - 4)$

[M1]

r	0	1	2	3	4	5
$x_r$	-0.5	-0.26667	-0.25452	-0.25412	-0.2541	-0.2541

[M1A1]

-0.2541 secure to 4 dp

[A1]

[subtotal 6]

[TOTAL 18]

# 4777 Numerical Computation

1 Eg:  $e_{r+1}$  is approximately  $ke_r$  [E2]

(i) Uses  $y_0 = \alpha + e_0$ ,  $y_1 = \alpha + ke_0$ ,  $y_2 = \alpha + k^2e_0$  or equivalent [M1A1]

Convincing algebra to eliminate  $k$  hence given result [A1A1]  
[subtotal 6]


(ii) Convincing re-arrangement [A1]

$y_0$	$y_1$	$y_2$	extrap (new $y_0$ )	new $y_1$	new $y_2$	extrap	once	[M1A1]
1	0.908662	0.917409	0.916644	0.916648	0.916647	0.916647	twice	[M1A1]
4 or 5 sf looks secure [A1]								
[subtotal 6]								

(iii) [A1]

$x$	$y_0$	$y_1$	$y_2$	extrap (new $y_0$ )	new $y_1$	new $y_2$	extrap	
1.1	1	0.908662	0.917409	0.916644	0.916648	0.916647	0.916647	
1.2	0.916647	0.845937	0.858962	0.856936	0.85695	0.856947	0.856948	set up
1.3	0.856948	0.799744	0.815042	0.811814	0.81184	0.811833	0.811835	SS
1.4	0.811835	0.763904	0.780556	0.776263	0.776302	0.776288	0.776292	[M2A2]
1.5	0.776292	0.734953	0.752555	0.747298	0.747351	0.747329	0.747335	
1.6	0.747335	0.7108	0.729213	0.723043	0.72311	0.723076	0.723087	values
1.7	0.723087	0.690112	0.70934	0.702258	0.70234	0.702292	0.70231	[A3]
1.8	0.70231	0.671996	0.692131	0.684095	0.684194	0.684128	0.684155	
1.9	0.684155	0.655831	0.677026	0.667954	0.668075	0.667985	0.668023	
2	0.668023	0.641175	0.663627	0.653402	0.65355	0.653427	0.653483	[A1]
3 or 4 sf looks secure								

$x$	$y$
0.653483	2
0.668023	1.9
0.684155	1.8
0.70231	1.7
0.723087	1.6
0.747335	1.5
0.776292	1.4
0.811835	1.3
0.856948	1.2
0.916647	1.1
1	1
1.1	0.916647
1.2	0.856948
1.3	0.811835
1.4	0.776292
1.5	0.747335
1.6	0.723087
1.7	0.70231
1.8	0.684155
1.9	0.668023
2	0.653483



organise

data

[M1A1]

graph

G2

Sub Total 12

TOTAL 24




2  $T_n - I = A_2h^2 + A_4h^4 + A_6h^6 + \dots$

- (i)  $T_{2n} - I = A_2(h/2)^2 + A_4(h/2)^4 + A_6(h/2)^6 + \dots$  [M1A1]
  - $4(T_{2n} - I) - (T_n - I) = b_4h^4 + b_6h^6 + \dots$  [M1]
  - $4T_{2n} - T_n - 3I = b_4h^4 + b_6h^6 + \dots$  [A1]
  - $(4T_{2n} - T_n)/3 - I = B_4h^4 + B_6h^6 + \dots$  [A1]
  - $(T_n^* = (4T_{2n} - T_n)/3$  has error of order  $h^4$  as given)
  - $T_n^{**} = (16T_{2n}^* - T_n^*)/15$  has error of order  $h^6$  [B1]
- [subtotal 6]

(ii)	x	f(x)	T	T*	T**	(T***)		
	0	0						
	2	3.523188	3.523188					
	1	0.731059	2.492653	2.149141				
	0.5	0.155615					f:	[A1]
	1.5	1.839543	2.243905	2.160989	2.161779		T:	[M1A2]
	0.25	0.035136					T*:	[M1A1]
	0.75	0.382038					T**:	[M1A1]
	1.25	1.214531						
	1.75	2.609105	2.182155	2.161572	2.161611	2.161608		
	0.125	0.0083					answer:	[A1]
	0.375	0.083344						
	0.625	0.254435						
	0.875	0.540367						
	1.125	0.955439						
	1.375	1.509072						
	1.625	2.206199						
	1.875	3.048173	2.166744	2.161606	2.161609	2.161609		

[subtotal 9]

(iii)	k	I
	0	0
	0.25	0.002847
	0.5	0.024686
	0.75	0.089495
	1	0.225935
	1.25	0.466242
	1.5	0.845007
	1.75	1.398068
	2	2.161609



- modify SS [M2]
- values of I [A2]
- graph [G2]

[subtotal 6]


(iv)	k	I			
	1.57	0.980739	accept 1.57	evidence of t&e:	[M2]
	1.58	1.001291	or 1.58	result:	[A1]
	1.579	0.999223	(or in between)		

[subtotal 3]  
[TOTAL 24]

3 (i)	h	x	y	k 1	k 2	k 3	k 4
	0.2	0	0	0.2	0.110557	0.121189	0.086653
	0.2	0.2	0.125024	0.085978	0.063177	0.064854	0.046393
	0.2	0.4	0.189763	0.046408	0.031125	0.032033	0.018694
	0.2	0.6	0.221666	0.018708	0.007021	0.007628	-0.00291
	0.2	0.8	0.229182	-0.0029	-0.01239	-0.01194	-0.02066
	0.2	1	0.217146	-0.02065	-0.02863	-0.02828	-0.0357
	0.2	1.2	0.188783	-0.03569	-0.04256	-0.04228	-0.04872
	0.2	1.4	0.146433	-0.04871	-0.05472	-0.05449	-0.06015
	0.2	1.6	0.091887	-0.06015	-0.06547	-0.06527	-0.07031
	0.2	1.8	0.026567	-0.0703	-0.07506	-0.07488	-0.07941
	0.2	2	-0.04836	-0.0794	-0.08369	-0.08353	-0.08762
	0.2	2.2	-0.13194	-0.08761	-0.0915	-0.09136	-0.09507
	0.2	2.4	-0.22334	-0.09507	-0.0986	-0.09849	-0.10187
	0.2	2.6	-0.32186	-0.10187	-0.1051	-0.105	-0.1081
	0.2	2.8	-0.42689	-0.1081	-0.11107	-0.11097	-0.11382
	0.2	3	-0.53789	-0.11382	-0.11656	-0.11647	-0.1191

setup  
[M3]

values  
[A3]



[G2]

Maximum about (0.8, 0.23) root about 1.8

[A1A1A1]  
[subtotal11]

- (ii) Eg:  
 $h = 0.01$  gives  $(p, q)$  as  $(0.77, 0.22743)$  hence  $(0.77, 0.23)$   
 $h = 0.01$  gives root as between 1.87 and 1.88 accept either

[M2]  
[A1A1]  
[A1]  
[subtotal5]

(iii) Eg:

s	h	x	y	k 1	k 2	k 3	k 4
1	0.01	0	0	0.01	0.009	0.009025	0.008621
1	0.01	0.01	0.009112	0.008618	0.008314	0.008319	0.008065
1	0.01	0.02	0.017437	0.008065	0.007844	0.007847	0.007649
1	0.01	0.03	0.025286	0.007649	0.007468	0.00747	0.007303
1	0.01	0.04	0.032757	0.007303	0.007147	0.007148	0.007002

Mods  
[M3]  
t & e  
[M3]

$s = 0.715, h = 0.01$  gives root closest to  $x = 1$  accept 0.71 to 0.72

[A2]

[subtotal8]  
[TOTAL 24]

4  $Q = \sum (y - a - bx - cx^2)^2$  [M1]

(i)  $dQ/da = 0$   $\sum y = na + b \sum x + c \sum x^2$  as given [M1A1]  
gives


other equations:  $\sum xy = a \sum x + b \sum x^2 + c \sum x^3$  [B1]

$\sum x^2y = a \sum x^2 + b \sum x^3 + c \sum x^4$  [B1]

[subtotal 5]

(ii)

X	Y
0	1.02
0.5	2.08
1	2.73
1.5	3.14
2	2.87
2.5	2.22
3	1.43



[G2]

roughly parabolic (quadratic) in shape

[E1]

[subtotal 3]

(iii)

x	y	xy	x <sup>2</sup> y	x <sup>2</sup>	x <sup>3</sup>	x <sup>4</sup>
0	1.02	0	0	0	0	0
0.5	2.08	1.04	0.52	0.25	0.125	0.0625
1	2.73	2.73	2.73	1	1	1
1.5	3.14	4.71	7.065	2.25	3.375	5.0625
2	2.87	5.74	11.48	4	8	16
2.5	2.22	5.55	13.875	6.25	15.625	39.0625
3	1.43	4.29	12.87	9	27	81
<b>10.5</b>	<b>15.49</b>	<b>24.06</b>	<b>48.54</b>	<b>22.75</b>	<b>55.125</b>	<b>142.1875</b>

[M2]

[A2]

normal equations:

7	10.5	22.75	15.49	
10.5	22.75	55.125	24.06	
22.75	55.125	142.1875	48.54	a= 1.017619
	-6.46154	-21	0.554615	b= 2.562143
	-2.69231	-10.5	1.656923	
		-1.75	1.425833	c= -0.81476

form equations

[M1A1]

solution

[M2A2]

x	y	y fitted	residual	res <sup>2</sup>
0	1.02	1.017619	0.002381	5.67E-06
0.5	2.08	2.095	-0.015	0.000225
1	2.73	2.765	-0.035	0.001225
1.5	3.14	3.027619	0.112381	0.012629
2	2.87	2.882857	-0.01286	0.000165
2.5	2.22	2.330714	-0.11071	0.012258
3	1.43	1.37119	0.05881	0.003459
			<b>-3.6E-15</b>	<b>0.029967</b>

y fitted

[M1A1]

residuals

[M1A1]

residual sum is zero (except for rounding errors) as it should be

[E1]

residual sum of squares is 0.029967

[A1]

[subtotal 16]

[TOTAL 24]

# Grade Thresholds

Advanced GCE MEI Mathematics 3895 7895  
June 2008 Examination Series

## Unit Threshold Marks

Unit		Maximum Mark	A	B	C	D	E	U
All units	UMS	100	80	70	60	50	40	0
4751	Raw	72	61	53	45	37	30	0
4752	Raw	72	55	48	41	34	28	0
4753	Raw	72	59	52	46	40	33	0
4753/02	Raw	18	15	13	11	9	8	0
4754	Raw	90	75	67	59	51	43	0
4755	Raw	72	60	51	42	34	26	0
4756	Raw	72	57	51	45	39	33	0
4757	Raw	72	50	44	38	33	28	0
4758	Raw	72	58	50	42	34	26	0
4758/02	Raw	18	15	13	11	9	8	0
4761	Raw	72	57	48	39	30	22	0
4762	Raw	72	56	48	40	33	26	0
4763	Raw	72	53	45	37	29	21	0
4764	Raw	72	55	47	40	33	26	0
4766	Raw	72	53	45	38	31	24	0
4767	Raw	72	57	49	41	33	26	0
4768	Raw	72	56	49	42	35	28	0
4769	Raw	72	57	49	41	33	25	0
4771	Raw	72	58	51	44	37	31	0
4772	Raw	72	51	44	37	31	25	0
4773	Raw	72	51	44	37	30	24	0
4776	Raw	72	57	49	41	34	26	0
4776/02	Raw	18	14	12	10	8	7	0
4777	Raw	72	54	46	39	32	25	0

### Specification Aggregation Results

Overall threshold marks in UMS (ie after conversion of raw marks to uniform marks)

	Maximum Mark	A	B	C	D	E	U
<b>7895-7898</b>	600	480	420	360	300	240	0
<b>3895-3898</b>	300	240	210	180	150	120	0

The cumulative percentage of candidates awarded each grade was as follows:

	A	B	C	D	E	U	Total Number of Candidates
<b>7895</b>	42.5	63.7	79.2	90.7	97.5	100	9600
<b>7896</b>	58.0	78.2	89.2	95.3	98.7	100	1539
<b>7897</b>	73.5	85.3	88.2	100	100	100	34
<b>7898</b>	27.8	52.8	61.1	77.8	91.7	100	36
<b>3895</b>	30.5	46.0	60.6	73.6	83.7	100	12767
<b>3896</b>	49.7	68.6	81.4	90.0	95.2	100	2039
<b>3897</b>	82.1	88.5	92.3	97.4	100	100	78
<b>3898</b>	47.8	52.2	69.6	87.0	95.7	100	23

For a description of how UMS marks are calculated see:  
[http://www.ocr.org.uk/learners/ums\\_results.html](http://www.ocr.org.uk/learners/ums_results.html)

Statistics are correct at the time of publication.

**OCR (Oxford Cambridge and RSA Examinations)**  
1 Hills Road  
Cambridge  
CB1 2EU

**OCR Customer Contact Centre**

**14 – 19 Qualifications (General)**  
Telephone: 01223 553998  
Facsimile: 01223 552627  
Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

**[www.ocr.org.uk](http://www.ocr.org.uk)**

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

**Oxford Cambridge and RSA Examinations**  
is a Company Limited by Guarantee  
Registered in England  
Registered Office; 1 Hills Road, Cambridge, CB1 2EU  
Registered Company Number: 3484466  
OCR is an exempt Charity



**OCR (Oxford Cambridge and RSA Examinations)**  
Head office  
Telephone: 01223 552552  
Facsimile: 01223 552553