


1)  $x^2 - 12x + 1 \equiv (x-6)^2 - 36 + 1 \equiv (x-6)^2 - 35$  Core

2) (i)

(ii) A translation through 1 unit to the right parallel to the x-axis.

(2)



3)

$y = x^3 - 4x^2 + 7$

$\frac{dy}{dx} = 3x^2 - 8x = -4$  when  $x=2$

The product of the gradients of perpendicular lines is -1

$\therefore$  Gradient of normal =  $-\frac{1}{-4} = \frac{1}{4}$

$y - y_1 = m(x - x_1)$

$\therefore$  equation of normal is at (2, 1)

$y - 1 = \frac{1}{4}(x - 2)$

$4y + 4 = x - 2$

$-x + 4y + 6 = 0$

(7)

4) (i)  $3^m = 81 = 3^4 \therefore m = 4$

(ii)  $(36p^4)^{\frac{1}{2}} = 24 \therefore 36^{\frac{1}{2}} p^2 = 24$

$\therefore m = 4$

$\therefore 36^{\frac{1}{2}} p^2 = 24$

$\therefore 6p^2 = 24$

$\therefore p^2 = 4$

$\therefore p = \pm 2$  (3)

(iii)  $5^n \times 5^{n+4} = 25$

$\therefore 5^{2n+4} = 5^2$

$\therefore 2n+4 = 2$

$\therefore 2n = -2$

$\therefore n = -1$  (3)

5)

$x - 8\sqrt{x} + 13 = 0$  Let  $u = \sqrt{x} \therefore u^2 - 8u + 13 = 0$

$x = \frac{-(-8) \pm \sqrt{(-8)^2 - 4 \times 1 \times 13}}{2 \times 1} = \frac{8 \pm \sqrt{12}}{2} = \frac{8 \pm 2\sqrt{3}}{2} = 4 \pm \sqrt{3}$

$\therefore \sqrt{x} = 4 + \sqrt{3}$

or  $\sqrt{x} = 4 - \sqrt{3}$

$x = 16 + 8\sqrt{3} + 3$

or  $x = 16 - 8\sqrt{3} + 3$

$x = 19 \pm 8\sqrt{3}$

(7)

6) (i)

$y = x^2 + 5$

$\therefore \frac{dy}{dx} = 2x$

at A(1, 6)

gradient = 2

(2)

(ii)

Gradient of AB =  $\frac{a^2 + 5 - 6}{a - 1} = \frac{a^2 - 1}{a - 1}$

=  $\frac{(a+1)(a-1)}{a-1} = a+1 = 2.3$

$\therefore a = 1.3$

(4)

(iii)

Gradients of AC is a number between 2 and 2.3 e.g. 2.1

(1)

7) (i)

(a)  $y = (3-x)^2$  corresponds to Fig. 3

(b)  $y = x^2 + 9$  corresponds to Fig. 1

(c)  $y = (3-x)(x+3)$  corresponds to Fig. 4

(1)  
(1)  
(1)

(ii)

The equation of the curve is  $y = -(x-3)^2$

(2)

8) (i)

$x^2 + y^2 + 6x - 4y - 4 = 0$

$(x+3)^2 - 9 + (y-2)^2 - 4 - 4 = 0$

$(x+3)^2 + (y-2)^2 = 17$

$\therefore$  centre is (-3, 2) radius =  $\sqrt{17}$

(3)

(ii)

To find the co-ordinates of the points of intersection it is necessary to solve  $(x+3)^2 + (y-2)^2 = 17$  and  $y = 3x+4$  as simultaneous equations.

Substituting for y

$(x+3)^2 + (3x+4-2)^2 = 17$

$(x+3)^2 + (3x+2)^2 = 17$

$x^2 + 6x + 9$

$+ 9x^2 + 12x + 4 = 17$

8 cont.)

$$10x^2 + 18x + 13 = 17$$

$$10x^2 + 18x - 4 = 0$$

$$5x^2 + 9x - 2 = 0$$

$$(5x - 1)(x + 2) = 0$$

$$x = \frac{1}{5} \quad \text{or} \quad x = -2$$

when  $x = \frac{1}{5}$ ,  $y = 3 \times \frac{1}{5} + 4 = 4\frac{3}{5}$  ; when  $x = -2$ ,  $y = 3 \times -2 + 4 = -2$

∴ the circle meets the line  $y = 3x + 4$  at  $(\frac{1}{5}, 4\frac{3}{5})$  and  $(-2, -2)$ . (6)

9)  $f(x) = \frac{1}{x} - \sqrt{x} + 3 = x^{-1} - x^{\frac{1}{2}} + 3$

(i) ∴  $f'(x) = -x^{-2} - \frac{1}{2}x^{-\frac{1}{2}} = -\frac{1}{x^2} - \frac{1}{2\sqrt{x}}$  (3)

(ii)  $f''(x) = 2x^{-3} + \frac{1}{4}x^{-\frac{3}{2}} = \frac{2}{x^3} + \frac{1}{4(\sqrt{x})^3}$


when  $x = 4$ ,  $f''(4) = \frac{2}{4^3} + \frac{1}{4(\sqrt{4})^3} = \frac{2}{64} + \frac{1}{32} = \frac{2}{32} = \frac{1}{16}$  (5)

10) If the quadratic  $kx^2 - 30x + 25k = 0$  has equal roots then the

discriminant  $b^2 - 4ac = 0$  i.e.  $(-30)^2 - 4 \times k \times 25k = 0$  (4)

$$900 - 100k^2 = 0$$

$$900 = 100k^2 \quad \therefore k^2 = 9 \quad \therefore k = \pm 3$$



Using Pythagorean Theorem

$$k^2 = (3x)^2 + (4x)^2 = 9x^2 + 16x^2 = 25x^2$$

$$\therefore k = 5x$$

∴ perimeter  $P = (2+5x) + 5x + (2+x) + 3x = 4 + 14x$  (2)

(ii) Area of a trapezium =  $\frac{1}{2}(a+b)h = \frac{1}{2}[2+x+2+5x] \times 3x = \frac{1}{2}[4+6x]3x = 6x + 9x^2$  (2)

$P \geq 39 \text{ m}$        $A < 99 \text{ m}^2$

∴  $4 + 14x \geq 39 \text{ m}$       ∴  $14x \geq 35 \text{ m}$        $x \geq \frac{35}{14}$  ,  $x \geq \frac{5}{2} \text{ m}$

$$9x^2 + 6x < 99$$

$$9x^2 + 6x - 99 < 0$$

$$3(3x^2 + 2x - 33) < 0$$



$$3(3x + 11)(x - 3) < 0$$

It can be seen from the graph that

$$-\frac{11}{3} < x < 3$$

To satisfy both conditions

$$\frac{5}{2} \leq x < 3$$



(7)