

Mathematics

Advanced GCE A2 7890 - 2

Advanced Subsidiary GCE AS 3890 - 2

Mark Schemes for the Units

June 2007

3890-2/7890-2/MS/R/07

OCR (Oxford, Cambridge and RSA Examinations) is a unitary awarding body, established by the University of Cambridge Local Examinations Syndicate and the RSA Examinations Board in January 1998. OCR provides a full range of GCSE, A level, GNVQ, Key Skills and other qualifications for schools and colleges in the United Kingdom, including those previously provided by MEG and OCEAC. It is also responsible for developing new syllabuses to meet national requirements and the needs of students and teachers.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2007

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 870 6622
Facsimile: 0870 870 6621
E-mail: publications@ocr.org.uk

CONTENTS

Advanced GCE Mathematics (7890)
 Advanced GCE Pure Mathematics (7891)
 Advanced GCE Further Mathematics (7892)



Advanced Subsidiary GCE Mathematics (3890)
 Advanced Subsidiary GCE Pure Mathematics (3891)
 Advanced Subsidiary GCE Further Mathematics (3892)

MARK SCHEME ON THE UNITS

Unit	Content	Page
4721	Core Mathematics 1	1
4722	Core Mathematics 2	7
4723	Core Mathematics 3	11
4724	Core Mathematics 4	15
4725	Further Pure Mathematics 1	19
4726	Further Pure Mathematics 2	25
4727	Further Pure Mathematics 3	31
4728	Mechanics 1	37
4729	Mechanics 2	43
4730	Mechanics 3	47
4731	Mechanics 4	53
4732	Probability & Statistics 1	59
4733	Probability & Statistics 2	65
4734	Probability & Statistics 3	69
4735	Probability & Statistics 4	75
4736	Decision Mathematics 1	79
4737	Decision Mathematics 2	85
*	Grade Thresholds	91


Mark Scheme 4721

June 2007

1	$(4x^2 + 20x + 25) - (x^2 - 6x + 9)$ $= 3x^2 + 26x + 16$ <p><u>Alternative method using difference of two squares:</u></p> $(2x + 5 + (x - 3))(2x + 5 - (x - 3))$ $= (3x + 2)(x + 8)$ $= 3x^2 + 26x + 16$	M1 A1 A1 3 3	Square one bracket to give an expression of the form $ax^2 + bx + c$ ($a \neq 0, b \neq 0, c \neq 0$) One squared bracket fully correct All 3 terms of final answer correct M1 2 brackets with same terms but different signs A1 One bracket correctly simplified A1 All 3 terms of final answer correct
2 (a)(i)		B1 B1 2 B1 1	Excellent curve for $\frac{1}{x}$ in either quadrant Excellent curve for $\frac{1}{x}$ in other quadrant SR B1 Reasonably correct curves in 1 st and 3 rd quadrants Correct graph, minimum point at origin, symmetrical
(ii)		B1 B1 2 5	
(b)	Stretch Scale factor 8 in y direction or scale factor $\frac{1}{2}$ in x direction	B1 B1 2 5	
3 (i)	$3\sqrt{20} \text{ or } 3\sqrt{2} \sqrt{5} \times \sqrt{2} \text{ or } \sqrt{180}$ $\text{or } \sqrt{90} \times \sqrt{2}$ $= 6\sqrt{5}$	M1 A1 2	Correctly simplified answer
(ii)	$10\sqrt{5} + 5\sqrt{5}$ $= 15\sqrt{5}$	M1 B1 A1 3 5	Attempt to change both surds to $\sqrt{5}$ One part correct and fully simplified cao

4 (i)	$(-4)^2 - 4 \times k \times k$ $= 16 - 4k^2$	M1 A1 2	Uses $b^2 - 4ac$ (involving k) $16 - 4k^2$
(ii)	$16 - 4k^2 = 0$ $k^2 = 4$ $k = 2$ or $k = -2$	M1 B1 B1 3 5	Attempts $b^2 - 4ac = 0$ (involving k) or attempts to complete square (involving k)
5 (i)	Length = $20 - 2x$ Area = $x(20 - 2x)$ $= 20x - 2x^2$	M1 A1 2	Expression for length of enclosure in terms of x Correctly shows that area = $20x - 2x^2$ AG
(ii)	$\frac{dA}{dx} = 20 - 4x$ For max, $20 - 4x = 0$ $x = 5$ only Area = 50	M1 M1 A1 A1 4 6	Differentiates area expression Uses $\frac{dy}{dx} = 0$
6	Let $y = (x + 2)^2$ $y^2 + 5y - 6 = 0$ $(y + 6)(y - 1) = 0$ $y = -6$ or $y = 1$ $(x + 2)^2 = 1$ $x = -1$ or $x = -3$	B1 M1 A1 M1 A1 A1 6 6	Substitute for $(x + 2)^2$ to get $y^2 + 5y - 6 (= 0)$ Correct method to find roots Both values for y correct Attempt to work out x One correct value Second correct value and no extra real values
7 (a)	$f(x) = x + 3x^{-1}$ $f'(x) = 1 - 3x^{-2}$	M1 A1 A1 A1 4	Attempt to differentiate First term correct x^{-2} soi www Fully correct answer
(b)	$\frac{dy}{dx} = \frac{5}{2}x^{\frac{3}{2}}$ When $x = 4$, $\frac{dy}{dx} = \frac{5}{2}\sqrt{4^3}$ $= 20$	M1 B1 B1 M1 A1 5 9	Use of differentiation to find gradient $\frac{5}{2}x^c$ $kx^{\frac{3}{2}}$ $\sqrt{4^3}$ soi SR If 0 scored for first 3 marks, award B1 if $\sqrt{4^n}$ correctly evaluated.

8 (i)	$(x + 4)^2 - 16 + 15$ $= (x + 4)^2 - 1$	B1 M1 A1 3	a = 4 15 – their a^2 cao in required form
(ii)	(-4, -1)	B1 ft B1 ft 2	Correct x coordinate Correct y coordinate
(iii)	$x^2 + 8x + 15 > 0$ $(x + 5)(x + 3) > 0$ $x < -5, x > -3$	M1 A1 M1 A1 4	Correct method to find roots -5, -3 Correct method to solve quadratic inequality eg +ve quadratic graph $x < -5, x > -3$ (not wrapped, strict inequalities, no 'and')
9 (i)	$(x - 3)^2 - 9 + y^2 - k = 0$ $(x - 3)^2 + y^2 = 9 + k$ Centre (3, 0) $9 + k = 4^2$ $k = 7$	B1 B1 M1 A1 4	$(x - 3)^2$ soi Correct centre Correct value for k (may be embedded) <u>Alternative method using expanded form:</u> Centre $(-g, -f)$ M1 Centre (3, 0) A1 $4 = \sqrt{f^2 + g^2} - (-k)$ M1 $k = 7$ A1
(ii)	$(3 - 3)^2 + y^2 = 16$ $y^2 = 16$ $y = 4$ Length of AB = $\sqrt{(-1 - 3)^2 + (0 - 4)^2}$ $= \sqrt{32}$ $= 4\sqrt{2}$	M1 A1 M1 A1 ft A1 5	Attempt to substitute $x = 3$ into original equation or their equation $y = 4$ (do not allow ± 4) Correct method to find line length using Pythagoras' theorem $\sqrt{32}$ or $\sqrt{16 + a^2}$ cao
(iii)	Gradient of AB = 1 or $\frac{a}{4}$ $y - 0 = m(x + 1)$ or $y - 4 = m(x - 3)$ $y = x + 1$	B1 ft M1 A1 3	Attempts equation of straight line through their A or B with their gradient Correct equation in any form with simplified constants
		12	

10 (i)	$(3x + 1)(x - 5) = 0$ $x = \frac{-1}{3}$ or $x = 5$	M1 A1 A1 3	Correct method to find roots Correct brackets or formula Both values correct
(ii)		B1 B1 B1 ft 3	SR B1 for $x = 5$ spotted www Positive quadratic (must be reasonably symmetrical) y intercept correct both x intercepts correct
(iii)	$\frac{dy}{dx} = 6x - 14$ $6x - 14 = 4$ $x = 3$ On curve, when $x = 3$, $y = -20$ $-20 = (4 \times 3) + c$ $c = -32$ <u>Alternative method:</u> $3x^2 - 14x - 5 = 4x + c$ $3x^2 - 18x - 5 - c = 0$ has one solution $b^2 - 4ac = 0$ $(-18)^2 - (4 \times 3 \times (-5 - c)) = 0$ $c = -32$	M1* M1* A1 A1 ft M1dep A1 6 M1 B1 M1 M1 A1 A1 12	Use of differentiation to find gradient of curve Equating their gradient expression to 4 Finding y co ordinate for their x value N.B. dependent on both previous M marks Equate curve and line (or substitute for x) Statement that only one solution for a tangent (may be implied by next line) Use of discriminant = 0 Attempt to use a, b, c from their equation Correct equation $c = -32$

Mark Scheme 4722


June 2007



<p>1 (i) $u_2 = 12$ $u_3 = 9.6, u_4 = 7.68$ (or any exact equivalents)</p> <p>(ii) $S_{20} = \frac{15(1-0.8^{20})}{1-0.8}$ $= 74.1$</p> <p style="text-align: center;">OR</p>	<p>B1 B1✓ 2</p> <p>M1 A1 A1 3</p> <p>M1 A2</p> <p style="text-align: center;">5</p>	<p>State $u_2 = 12$ Correct u_3 and u_4 from their u_2</p> <p>Attempt use of $S_n = \frac{a(1-r^n)}{1-r}$, with $n = 20$ or 19 Obtain correct unsimplified expression Obtain 74.1 or better</p> <p>List all 20 terms of GP Obtain 74.1</p>
<p>2 $(x + \frac{2}{x})^4 = x^4 + 4x^3(\frac{2}{x}) + 6x^2(\frac{2}{x})^2 + 4x(\frac{2}{x})^3 + (\frac{2}{x})^4$</p> <p style="text-align: center;">OR</p> <p>$= x^4 + 8x^2 + 24 + \frac{32}{x^2} + \frac{16}{x^4}$ (or equiv)</p> <p style="text-align: center;">OR</p>	<p>M1*</p> <p>M1* A1dep* A1 A1 5</p> <p>M1* M1*</p> <p>A1dep* A1 A1</p> <p style="text-align: center;">5</p>	<p>Attempt expansion, using powers of x and $\frac{2}{x}$ (or the two terms in their bracket), to get at least 4 terms Use binomial coefficients of 1, 4, 6, 4, 1 Obtain two correct, simplified, terms Obtain a further one correct, simplified, term Obtain a fully correct, simplified, expansion</p> <p>Attempt expansion using all four brackets Obtain expansion containing the correct 5 powers only (could be unsimplified powers eg $x^3 \cdot x^{-1}$)</p> <p>Obtain two correct, simplified, terms Obtain a further one correct, simplified, term Obtain a fully correct, simplified, expansion</p>
<p>3 $\log 3^{(2x+1)} = \log 5^{200}$ $(2x+1)\log 3 = 200\log 5$</p> <p style="text-align: center;">OR</p> <p>$2x + 1 = \frac{200\log 5}{\log 3}$ $x = 146$</p> <p>$(2x + 1) = \log_3 5^{200}$ $2x + 1 = 200\log_3 5$</p>	<p>M1 M1 A1 M1 A1 5</p> <p>M1 M1 A1 M1 A1</p> <p style="text-align: center;">5</p>	<p>Introduce logarithms throughout Drop power on at least one side Obtain correct linear equation (now containing no powers) Attempt solution of linear equation Obtain $x = 146$, or better</p> <p>Introduce \log_3 on right-hand side Drop power of 200 Obtain correct equation Attempt solution of linear equation Obtain $x = 146$, or better</p>
<p>4 (i) $\text{area} \approx \frac{1}{2} \times \frac{1}{2} \times \{ \sqrt{5} + 2(\sqrt{7} + \sqrt{9} + \sqrt{11}) + \sqrt{13} \}$</p> <p style="text-align: center;">OR</p> <p>$\approx 0.25 \times 23.766\dots$ ≈ 5.94</p> <p>(ii) This is an underestimate..... ...as the tops of the trapezia are below the curve</p>	<p>M1 M1 A1 A1 4</p> <p>*B1 B1dep*B1 2</p> <p style="text-align: center;">6</p>	<p>Attempt y-values for at least 4 of the $x = 1, 1.5, 2, 2.5, 3$ only Attempt to use correct trapezium rule Obtain $\frac{1}{2} \times \frac{1}{2} \times \{ \sqrt{5} + 2(\sqrt{7} + \sqrt{9} + \sqrt{11}) + \sqrt{13} \}$, or decimal equiv Obtain 5.94 or better (answer only is 0/4)</p> <p>State underestimate Correct statement or sketch</p>



<p>5 (i) $3(1 - \sin^2 \theta) = \sin \theta + 1$ $3 - 3 \sin^2 \theta = \sin \theta + 1$ $3 \sin^2 \theta + \sin \theta - 2 = 0$</p> <p>(ii) $(3 \sin \theta - 2)(\sin \theta + 1) = 0$ $\sin \theta = \frac{2}{3}$ or -1 $\theta = 42^\circ, 138^\circ, 270^\circ$</p>	<p>M1 A1 2 M1 A1 A1 A1 A1 A1 A1 A1 A1</p> <p>5</p>	<p>Use $\cos^2 \theta = 1 - \sin^2 \theta$</p> <p>Show given equation correctly</p> <p>Attempt to solve quadratic equation in $\sin \theta$</p> <p>Both values of $\sin \theta$ correct</p> <p>Correct answer of 270°</p> <p>Correct answer of 42°</p> <p>For correct non-principal value answer, following their first value of θ in the required range (any extra values for θ in required range is max 4/5) (radians is max 4/5)</p> <p>SR: answer only (or no supporting method) is B1 for 42°, B1 for 138°, B1 for 270°</p> <p style="text-align: center;">7</p>
<p>6 (a) (i) $\int x^3 - 4x = \frac{1}{4}x^4 - 2x^2 + c$</p> <p>(ii) $\left[\frac{1}{4}x^4 - 2x^2\right]_1^6$ $= (324 - 72) - (\frac{1}{4} - 2)$ $= 253\frac{3}{4}$</p> <p>(b) $\int 6x^{-3} dx = -3x^{-2} + c$</p>	<p>M1 A1 B1 3 M1 A1 2 B1 M1 A1 3</p> <p>8</p>	<p>Expand and attempt integration</p> <p>Obtain $\frac{1}{4}x^4 - 2x^2$ (A0 if \int or dx still present) + c (mark can be given in (b) if not gained here)</p> <p>Use limits correctly in integration attempt (ie F(6) - F(1))</p> <p>Obtain $253\frac{3}{4}$ (answer only is M0A0)</p> <p>Use of $\frac{1}{x^3} = x^{-3}$</p> <p>Obtain integral of the form kx^2</p> <p>Obtain correct $-3x^{-2} (+c)$ (A0 if \int or dx still present, but only penalise once in question)</p> <p style="text-align: center;">8</p>
<p>7 (a) $S_{70} = \frac{70}{2}\{(2 \times 12) + (70 - 1)d\}$ $35(24 + 69d) = 12915$ $d = 5$</p> <p>OR</p> <p>$\frac{70}{2}\{12 + l\} = 12915$ $l = 357$ $12 + 69d = 357$ $d = 5$</p> <p>(b) $ar = -4$ $\frac{a}{1-r} = 9$ $\frac{4}{r} = 9 - 9r$ or $a = 9 - (9 \times \frac{4}{a})$ $9r^2 - 9r - 4 = 0$ $a^2 - 9a - 36 = 0$ $(3r - 4)(3r + 1) = 0$ $(a + 3)(a - 12) = 0$ $r = \frac{4}{3}, r = -\frac{1}{3}$ $a = -3, a = 12$ Hence $r = -\frac{1}{3}$</p>	<p>M1 A1 M1 A1 4 M1 A1 M1 A1 B1 B1 M1 A1 M1 A1 A1</p> <p>7</p> <p style="text-align: center;">11</p>	<p>Attempt S_{70}</p> <p>Obtain correct unsimplified expression</p> <p>Equate attempt at S_{70} to 12915, and attempt to find d</p> <p>Obtain $d = 5$</p> <p>Attempt to find d by first equating $^n/2 (a + l)$ to 12915</p> <p>Obtain $l = 357$</p> <p>Equate u_{70} to l</p> <p>Obtain $d = 5$</p> <p>Correct statement for second term</p> <p>Correct statement for sum to infinity</p> <p>Attempt to eliminate either a or r</p> <p>Obtain correct equation (no algebraic denominators/brackets)</p> <p>Attempt solution of three term quadratic equation</p> <p>Obtain at least $r = -\frac{1}{3}$ (from correct working only)</p> <p>Obtain $r = -\frac{1}{3}$ only (from correct working only)</p> <p>SR: answer only / T&I is B2 only</p>



<p>8 (i) $\frac{1}{2} \times AB^2 \times 0.9 = 16.2$ $AB^2 = 36 \Rightarrow AB = 6$</p> <p>(ii) $\frac{1}{2} \times 6 \times AC \times \sin 0.9 = 32.4$ $AC = 13.8$ cm</p> <p>(iii) $BC^2 = 6^2 + 13.8^2 - 2 \times 6 \times 13.8 \times \cos 0.9$ Hence $BC = 11.1$ cm</p> <p>$BD = 6 \times 0.9 = 5.4$ cm Hence perimeter = $11.1 + 5.4 + (13.8 - 6)$ = 24.3 cm</p>	<p>M1 A1 2 16.2)</p> <p>M1* M1 dep* A1 3</p> <p>M1 A1√ A1</p> <p>B1 M1 A1 6</p> <p>11</p>	<p>Use $(\frac{1}{2})r^2\theta = 16.2$ Confirm $AB = 6$ cm (or verify $\frac{1}{2} \times 6^2 \times 0.9 = 16.2$)</p> <p>Use $\Delta = \frac{1}{2} bc \sin A$, or equiv Equate attempt at area to 32.4 Obtain $AC = 13.8$ cm, or better</p> <p>Attempt use of correct cosine formula in ΔABC Correct unsimplified equation, from their AC Obtain $BC = 11.1$ cm, or anything that rounds to this</p> <p>State $BD = 5.4$ cm (seen anywhere in question) Attempt perimeter of region BCD Obtain 24.3 cm, or anything that rounds to this</p>
<p>9 (i) (a) $f(-1) = -1 + 6 - 1 - 4 = 0$</p> <p>(b) $x = -1$ $f(x) = (x+1)(x^2 + 5x - 4)$</p> $x = \frac{-5 \pm \sqrt{25+16}}{2}$ $x = \frac{1}{2}(-5 \pm \sqrt{41})$ <p>(ii) (a) $\log_2(x+3)^2 + \log_2 x - \log_2(4x+2) = 1$</p> $\log_2\left(\frac{(x+3)^2 x}{4x+2}\right) = 1$ $\frac{(x+3)^2 x}{4x+2} = 2$ $(x^2 + 6x + 9)x = 8x + 4$ $x^3 + 6x^2 + x - 4 = 0$ <p>(b) $x > 0$, otherwise $\log_2 x$ is undefined $x = \frac{1}{2}(-5 + \sqrt{41})$</p>	<p>B1 1</p> <p>B1 M1 A1 A1 M1</p> <p>A1 6</p> <p>B1 M1</p> <p>A1</p> <p>B1</p> <p>A1 5</p> <p>B1* B1√dep* 2</p> <p>14</p>	<p>Confirm $f(-1) = 0$, through any method</p> <p>State $x = -1$ at any point Attempt complete division by $(x + 1)$, or equiv Obtain $x^2 + 5x + k$ Obtain completely correct quotient</p> <p>Attempt use of quadratic formula, or equiv, find roots Obtain $\frac{1}{2}(-5 \pm \sqrt{41})$</p> <p>State or imply that $2\log(x+3) = \log(x+3)^2$ Add or subtract two, or more, of their algebraic logs correctly</p> <p>Obtain correct equation (or any equivalent, with single term on each side)</p> <p>Use $\log_2 a = 1 \Rightarrow a = 2$ at any point</p> <p>Confirm given equation correctly</p> <p>State or imply that $\log x$ only defined for $x > 0$ State $x = \frac{1}{2}(-5 + \sqrt{41})$ (or $x = 0.7$) only, following their single positive root in (i)(b)</p>

**Mark Scheme 4723
June 2007**

1 (i)	Attempt use of product rule	M1	
	Obtain $3x^2(x+1)^5 + 5x^3(x+1)^4$	A1	2 or equiv
	[Or: (following complete expansion and differentiation term by term)		
	Obtain $8x^7 + 35x^6 + 60x^5 + 50x^4 + 20x^3 + 3x^2$	B2	allow B1 if one term incorrect]
(ii)	Obtain derivative of form $kx^3(3x^4 + 1)^n$	M1	any constants k and n
	Obtain derivative of form $kx^3(3x^4 + 1)^{-\frac{1}{2}}$	M1	
	Obtain correct $6x^3(3x^4 + 1)^{\frac{1}{2}}$	A1	3 or (unsimplified) equiv
<hr/>			
2	Identify critical value $x = 2$	B1	
	Attempt process for determining both critical values	M1	
	Obtain $\frac{1}{3}$ and 2	A1	
	Attempt process for solving inequality	M1	table, sketch ...; implied by plausible answer
	Obtain $\frac{1}{3} < x < 2$	A1	5
<hr/>			
3 (i)	Attempt correct process for composition	M1	numerical or algebraic
	Obtain (16 and hence) 7	A1	2
	(ii) Attempt correct process for finding inverse	M1	maybe in terms of y so far
	Obtain $(x - 3)^2$	A1	2 or equiv; in terms of x , not y
(iii)	Sketch (more or less) correct $y = f(x)$	B1	with 3 indicated or clearly implied on y -axis, correct curvature, no maximum point
	Sketch (more or less) correct $y = f^{-1}(x)$	B1	right hand half of parabola only
	State reflection in line $y = x$	B1	3 or (explicit) equiv; independent of earlier marks
<hr/>			
4 (i)	Obtain integral of form $k(2x + 1)^{\frac{4}{3}}$	M1	or equiv using substitution; any constant k
	Obtain correct $\frac{3}{8}(2x + 1)^{\frac{4}{3}}$	A1	or equiv
	Substitute limits in expression of form $(2x + 1)^n$ and subtract the correct way round	M1	using adjusted limits if subn used
	Obtain 30	A1	4
	(ii) Attempt evaluation of $k(y_0 + 4y_1 + y_2)$	M1	any constant k
Identify k as $\frac{1}{3} \times 6.5$	A1		
Obtain 29.6	A1	3 or greater accuracy (29.554566...)	
[SR: (using Simpson's rule with 4 strips)			
Obtain $\frac{1}{3} \times 3.25(1 + 4 \times \sqrt[3]{7.5} + 2 \times \sqrt[3]{14} + 4 \times \sqrt[3]{20.5} + 3)$ and hence 29.9	B1	or greater accuracy (29.897...)]	

5 (i)	State $e^{-0.04t} = 0.5$	B1	or equiv
	Attempt solution of equation of form $e^{-0.04t} = k$	M1	using sound process; maybe implied
	Obtain 17	A1	3 or greater accuracy (17.328...)
(ii)	Differentiate to obtain form $k e^{-0.04t}$	*M1	constant k different from 240
	Obtain $(\pm) 9.6e^{-0.04t}$	A1	or (unsimplified) equiv
	Equate attempt at first derivative to $(\pm) 2.1$ and attempt solution	M1	dep *M; method maybe implied
	Obtain 38	A1	4 or greater accuracy (37.9956...)
<hr/>			
6 (i)	Obtain integral of form $k_1 e^{2x} + k_2 x^2$	M1	any non-zero constants k_1, k_2
	Obtain correct $3e^{2x} + \frac{1}{2}x^2$	A1	
	Obtain $3e^{2a} + \frac{1}{2}a^2 - 3$	A1	
	Equate definite integral to 42 and attempt rearrangement	M1	using sound processes
	Confirm $a = \frac{1}{2}\ln(15 - \frac{1}{6}a^2)$	A1	5 AG; necessary detail required
(ii)	Obtain correct first iterate 1.348...	B1	
	Attempt correct process to find at least 2 iterates	M1	
	Obtain at least 3 correct iterates	A1	
	Obtain 1.344	A1	4 answer required to exactly 3 d.p.; allow recovery after error
[1 → 1.34844 → 1.34382 → 1.34389]			
<hr/>			
7 (i)	Show correct general shape (alternating above and below x -axis)	M1	with no branch reaching x -axis
	Draw (more or less) correct sketch	A1	2 with at least one of 1 and -1 indicated or clearly implied
(ii)	Attempt solution of $\cos x = \frac{1}{3}$	M1	maybe implied; or equiv
	Obtain 1.23 or 0.392π	A1	or greater accuracy
	Obtain 5.05 or 1.61π	A1	3 or greater accuracy and no others within $0 \leq x \leq 2\pi$; penalise answer(s) to 2sf only once
(iii)	<u>Either</u> : Obtain equation of form $\tan \theta = k$	M1	any constant k ; maybe implied
	Obtain $\tan \theta = 5$	A1	
	Obtain two values only of form $\theta, \theta + \pi$	M1	within $0 < x < 2\pi$; allow degrees at this stage
	Obtain 1.37 and 4.51 (or 0.437π and 1.44π)	A1	4 allow ± 1 in third sig fig; or greater accuracy
	<u>Or</u> : (for methods which involve squaring, etc.)		
Attempt to obtain eqn in one trig ratio	M1		
Obtain correct value	A1	$\tan^2 \theta = 25, \cos^2 \theta = \frac{1}{26}, \dots$	
Attempt solution at least to find one value in first quadrant and one value in third	M1		
Obtain 1.37 and 4.51 (or equivs as above)	A1	ignoring values in second and fourth quadrants	

8 (i)	Attempt use of quotient rule	M1	allow for numerator ‘wrong way round’; or equiv
	Obtain $\frac{(4 \ln x + 3)\frac{4}{x} - (4 \ln x - 3)\frac{4}{x}}{(4 \ln x + 3)^2}$	A1	or equiv
	Confirm $\frac{24}{x(4 \ln x + 3)^2}$	A1	3 AG; necessary detail required
(ii)	Identify $\ln x = \frac{3}{4}$	B1	or equiv
	State or imply $x = e^{\frac{3}{4}}$	B1	
	Substitute e^k completely in expression for derivative	M1	and deal with $\ln e^k$ term
	Obtain $\frac{2}{3} e^{-\frac{3}{4}}$	A1	4 or exact (single term) equiv
(iii)	State or imply $\int \frac{4\pi}{x(4 \ln x + 3)^2} dx$	B1	
	Obtain integral of form $k \frac{4 \ln x - 3}{4 \ln x + 3}$		
	or $k(4 \ln x + 3)^{-1}$	*M1	any constant k
	Substitute both limits and subtract right way round	M1	dep *M
	Obtain $\frac{4}{21}\pi$	A1	4 or exact equiv
<hr/>			
9 (i)	Attempt use of either of $\tan(A \pm B)$ identities	M1	
	Substitute $\tan 60^\circ = \sqrt{3}$ or $\tan^2 60^\circ = 3$	B1	
	Obtain $\frac{\tan \theta + \sqrt{3}}{1 - \sqrt{3} \tan \theta} \times \frac{\tan \theta - \sqrt{3}}{1 + \sqrt{3} \tan \theta}$	A1	or equiv (perhaps with $\tan 60^\circ$ still involved)
	Obtain $\frac{\tan^2 \theta - 3}{1 - 3 \tan^2 \theta}$	A1	4 AG
(ii)	Use $\sec^2 \theta = 1 + \tan^2 \theta$	B1	
	Attempt rearrangement and simplification of equation involving $\tan^2 \theta$	M1	or equiv involving $\sec \theta$
	Obtain $\tan^4 \theta = \frac{1}{3}$	A1	or equiv $\sec^2 \theta = 1.57735\dots$
	Obtain 37.2	A1	or greater accuracy
	Obtain 142.8	A1	5 or greater accuracy; and no others between 0 and 180
(iii)	Attempt rearrangement of $\frac{\tan^2 \theta - 3}{1 - 3 \tan^2 \theta} = k^2$ to form		
	$\tan^2 \theta = \frac{f(k)}{g(k)}$	M1	
	Obtain $\tan^2 \theta = \frac{k^2 + 3}{1 + 3k^2}$	A1	
	Observe that RHS is positive for all k , giving one value in each quadrant	A1	3 or convincing equiv

Mark Scheme 4724

June 2007

4724

Mark Scheme

June 2007

1	<p>(i) Correct format $\frac{A}{x+2} + \frac{B}{x-3}$ $A = 1$ and $B = 2$ (ii) $-A(x+2)^{-2} - B(x-3)^{-2}$ f.t. Convincing statement that each denom > 0 State whole exp < 0 AG</p>	<p>M1 A1 2 $\sqrt{A1}$ B1 B1 3</p>	<p>s.o.i. in answer for both accept ≥ 0. Do not accept $x^2 > 0$. <u>Dep on previous 4 marks.</u></p>	5
2	<p>Use parts with $u = x^2, dv = e^x$ Obtain $x^2 e^x - \int 2x e^x (dx)$ Attempt parts again with $u = (-)(2)x, dv = e^x$ Final = $(x^2 - 2x + 2)e^x$ AEF incl brackets Use limits correctly throughout $e^{(1)} - 2$ ISW Exact answer only</p>	<p>*M1 A1 M1 A1 dep*M1 A1 6</p>	<p>obtaining a result $f(x) + / - \int g(x)(dx)$ s.o.i. eg $e + (-2x + 2)e^x$ Tolerate (their value for $x = 1$) (-0) Allow 0.718 \rightarrow M1</p>	6
3	<p>Volume = $(k) \int_0^{\pi} \sin^2 x (dx)$ Suitable method for integrating $\sin^2 x$ $\int \sin^2 x (dx) = \frac{1}{2} \int 1 - \cos 2x (dx)$ $\int \cos 2x (dx) = \frac{1}{2} \sin 2x$ Use limits correctly Volume = $\frac{1}{2} \pi^2$ WWW Exact answer</p>	<p>B1 *M1 A1 A1 dep*M1 A1 6</p>	<p>where $k = \pi, 2\pi$ or 1; limits necessary eg $\int + / - 1 + / - \cos 2x (dx)$ or single integ by parts & connect to $\int \sin^2 x (dx)$ or $-\sin x \cos x + \int \cos^2 x (dx)$ or $-\sin x \cos x + \int 1 - \sin^2 x (dx)$ Beware: wrong working leading to $\frac{1}{2} \pi^2$</p>	6
4	<p>(i) $(1 + \frac{x}{2})^{-2}$ $= 1 + (-2)(\frac{x}{2}) + \frac{-2 \cdot -3}{2} (\frac{x}{2})^2 + \frac{-2 \cdot -3 \cdot -4}{3!} (\frac{x}{2})^3$ $= 1 - x$ $+ \frac{3}{4} x^2 - \frac{1}{2} x^3$ $(2 + x)^{-2} = \frac{1}{4} (\text{their exp of } (1 + ax)^{-2})$ mult out $x < 2$ or $-2 < x < 2$ (but not $\frac{1}{2}x < 1$) (ii) If (i) is $a + bx + cx^2 + dx^3$ evaluate $b + d$ $-\frac{3}{8} (x^3)$</p>	<p>M1 B1 A1 $\sqrt{B1}$ B1 5 M1 $\sqrt{A1}$ 2</p>	<p>Clear indication of method of ≥ 3 terms First two terms, not dependent on M1 For both third and fourth terms Correct: $\frac{1}{4} - \frac{1}{4}x + \frac{3}{16}x^2 - \frac{1}{8}x^3$ Follow-through from $b + d$</p>	7

<p>5(i) $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$ $= \frac{-4 \sin 2t}{-\sin t}$ $= 8 \cos t$ ≤ 8 AG</p> <p>(ii) Use $\cos 2t = 2 \cos^2 t + / - 1$ or $1 - 2 \cos^2 t$ Use correct version $\cos 2t = 2 \cos^2 t - 1$ Produce WWW $y = 4x^2 + 1$ AG</p> <p>(iii) U-shaped parabola above x-axis, sym abt y-axis Portion between $(-1, 5)$ and $(1, 5)$</p> <p>N.B. If (ii) answered or quoted before (i) attempted, allow in part</p>	<p>M1 A1 A1 A1 M1 A1 A1</p>	<p>Accept $\frac{4 \sin 2t}{\sin t}$ WWW 4 with brief explanation eg $\cos t \leq 1$ If <u>starting with</u> $y = 4x^2 + 1$, then Subst $x = \cos t$, $y = 3 + 2 \cos 2t$ M1 3 <u>Either</u> substitute <u>a</u> formula for $\cos 2t$ M1 Obtain $0=0$ or $4\cos^2 t + 1 = 4\cos^2 t + 1$ A1 <u>Or</u> Manip to give formula for $\cos 2t$ M1 Obtain corr formula & say it's correct A1 Any labelling must be correct 2 either $x = \pm 1$ or $y = 5$ must be marked (i) B2 for $\frac{dy}{dx} = 8x$ +B1,B1 if earned. 9</p>
<p>6 (i) $\frac{d}{dx}(y^2) = 2y \frac{dy}{dx}$ Using $d(uv) = u dv + v du$ for the $(3)xy$ term $\frac{d}{dx}(x^2 + 3xy + 4y^2) = 2x + 3x \frac{dy}{dx} + 3y + 8y \frac{dy}{dx}$ Solve for $\frac{dy}{dx}$ & subst $(x, y) = (2, 3)$</p> <p>$\frac{dy}{dx} = -\frac{13}{30}$ Grad normal = $\frac{30}{13}$ follow-through Find equ <u>any</u> line thro $(2, 3)$ with <u>any</u> num grad $30x - 13y - 21 = 0$ AEF</p>	<p>B1 M1 A1 M1 A1 √B1 M1 A1</p>	<p>or v.v. Subst now or at normal eqn stage; (M1 dep on either/both B1 M1 earned) Implied if grad normal = $\frac{30}{13}$ This f.t. mark awarded only if numerical 8 No fractions in final answer 8</p>
<p>7 (i) Leading term in quotient = $2x$ <u>Suff evidence</u> of division or identity process Quotient = $2x + 3$ Remainder = x</p> <p>(ii) their quotient + $\frac{\text{their remainder}}{x^2 + 4}$</p> <p>(iii) <u>Working with their expression in part (ii)</u> their $Ax + B$ integrated as $\frac{1}{2}Ax^2 + Bx$ their $\frac{Cx}{x^2 + 4}$ integrated as $k \ln(x^2 + 4)$ $k = \frac{1}{2}C$ Limits used correctly throughout $14 + \frac{1}{2} \ln \frac{13}{5}$</p>	<p>B1 M1 A1 A1 √B1 √B1 M1 √A1 M1 A1</p>	<p>Stated or in relevant position in division 4 Accept $\frac{x}{x^2 + 4}$ as remainder 1 $2x + 3 + \frac{x}{x^2 + 4}$ Ignore any integration of $\frac{D}{x^2 + 4}$ 5 logs need not be combined. 10</p>

<p>8</p> <p>(i) Sep variables eg $\int \frac{1}{6-h} (dh) = \int \frac{1}{20} (dt)$</p> <p>LHS = $-\ln(6-h)$</p> <p>RHS = $\frac{1}{20}t$ (+c)</p> <p>Subst $t = 0, h = 1$ into equation containing 'c'</p> <p>Correct value of their c = $-(20)\ln 5$ WWW</p> <p>Produce $t = 20 \ln \frac{5}{6-h}$ WWW AG</p> <p>(ii) When $h = 2, t = 20 \ln \frac{5}{4} = 4.46(2871)$</p> <p>(iii) Solve $10 = 20 \ln \frac{5}{6-h}$ to $\frac{5}{6-h} = e^{0.5}$</p> <p>$h = 2.97(2.9673467\dots)$</p> <p>[In (ii),(iii) accept non-decimal (exact) answers but -1 once.]</p> <p>Accept truncated values in (ii),(iii).</p> <p>(iv) Any indication of (approximately) 6 (m)</p>	<p>*M1</p> <p>A1</p> <p>A1</p> <p>dep*M1</p> <p>A1</p> <p>A1</p> <p>6</p> <p>B1</p> <p>1</p> <p>M1</p> <p>A1</p> <p>2</p> <p>B1</p> <p>1</p>	<p>s.o.i. Or $\frac{dt}{dh} = \frac{20}{6-h} \rightarrow M1$</p> <p>& then $t = -20 \ln(6-h)$ (+c) $\rightarrow A1+A1$</p> <p>or $(20)\ln 5$ if on LHS</p> <p>Must see $\ln 5 - \ln(6-h)$</p> <p>Accept 4.5, $4\frac{1}{2}$</p> <p>or $\frac{6-h}{5} = e^{-0.5}$ or suitable $\frac{1}{2}$-way stage</p> <p>$6 - 5e^{-0.5}$ or $6 - e^{1.109}$</p> <p style="text-align: right;">10</p>
<p>9</p> <p>(i) Use $-6\mathbf{i} + 8\mathbf{j} - 2\mathbf{k}$ and $\mathbf{i} + 3\mathbf{j} + 2\mathbf{k}$ only</p> <p>Correct method for scalar product</p> <p>Correct method for magnitude</p> <p>68 or 68.5 (68.47546); 1.2(0) (1.1951222) rad</p> <p>[N.B. 61 (60.562) will probably have been generated by $5\mathbf{i} - \mathbf{j} - 2\mathbf{k}$ and $3\mathbf{i} - 8\mathbf{j}$]</p> <p>(ii) Indication that relevant vectors are parallel</p> <p>$c = -4$</p> <p>(iii) Produce 2/3 equations containing t, u (& c)</p> <p>Solve the 2 equations not containing 'c'</p> <p>$t = 2, u = 1$</p> <p>Subst their (t, u) into equation containing c</p> <p>$c = -3$</p> <p>Alternative method for final 4 marks</p> <p>Solve two equations, one with 'c', for t and u in terms of c, and substitute into third equation</p> <p>$c = -3$</p>	<p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>4</p> <p>M1</p> <p>A1</p> <p>2</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>5</p> <p>(M2)</p> <p>(A2)</p>	<p>of <u>any</u> two vectors $(-6 + 24 - 4 = 14)$</p> <p>of <u>any</u> vector $(\sqrt{36 + 64 + 4} = \sqrt{104}$ or $\sqrt{1 + 9 + 4} = \sqrt{14})$</p> <p>$-6\mathbf{i} + 8\mathbf{j} - 2\mathbf{k}$ & $3\mathbf{i} + c\mathbf{j} + \mathbf{k}$ with some indic of method of attack</p> <p>eg $-6\mathbf{i} + 8\mathbf{j} - 2\mathbf{k} = \lambda(3\mathbf{i} + c\mathbf{j} + \mathbf{k})$</p> <p>$c = -4$ WW \rightarrow B2</p> <p>eg $3 + t = 2 + 3u, -8 + 3t = 1 + cu$ and $2t = 3 + u$</p> <p style="text-align: right;">11</p>

**Mark Scheme 4725
June 2007**

1	<p><i>EITHER</i> $a = 2$</p> <p>$b = 2\sqrt{3},$ <i>OR</i></p> <p>$a = 2 \quad b = 2\sqrt{3}$</p>	<p>M1 A1 M1 A1 M1 M1 A1 A1</p>	<p>4</p> <p>4</p>	<p>Use trig to find an expression for a (or b) Obtain correct answer Attempt to find other value Obtain correct answer a.e.f. (Allow 3.46) State 2 equations for a and b</p> <p>Attempt to solve these equations Obtain correct answers a.e.f. SR \pm scores A1 only</p>
2	<p>$(1^3 =) \frac{1}{4} \times 1^2 \times 2^2$</p> <p>$\frac{1}{4}n^2(n+1)^2 + (n+1)^3$</p> <p>$\frac{1}{4}(n+1)^2(n+2)^2$</p>	<p>B1 M1 M1(indep) A1 A1</p>	<p>5</p> <p>5</p>	<p>Show result true for $n = 1$</p> <p>Add next term to given sum formula Attempt to factorise and simplify Correct expression obtained convincingly</p> <p>Specific statement of induction conclusion</p>
3	<p>$3\Sigma r^2 - 3\Sigma r + \Sigma 1$</p> <p>$3\Sigma r^2 = \frac{1}{2}n(n+1)(2n+1)$</p> <p>$3\Sigma r = \frac{3}{2}n(n+1)$</p> <p>$\Sigma 1 = n$ n^3</p>	<p>M1 A1 A1 A1 M1 A1</p>	<p>6</p> <p>6</p>	<p>Consider the sum of three separate terms</p> <p>Correct formula stated</p> <p>Correct formula stated</p> <p>Correct term seen Attempt to simplify Obtain given answer correctly</p>
4	<p>(i) $\frac{1}{2} \begin{pmatrix} 5 & -1 \\ -3 & 1 \end{pmatrix}$</p> <p>(ii)</p> <p>$\frac{1}{2} \begin{pmatrix} 2 & 0 \\ 23 & -5 \end{pmatrix}$</p>	<p>B1 B1 M1 M1(indep) A1ft A1ft</p>	<p>2</p> <p>4</p> <p>6</p>	<p>Transpose leading diagonal and negate other diagonal or solve sim. eqns. to get 1st column Divide by the determinant or solve 2nd pair to get 2nd column</p> <p>Attempt to use $B^{-1}A^{-1}$ or find B Attempt at matrix multiplication One element correct, a.e.f. All elements correct, a.e.f. NB ft consistent with their (i)</p>

5	<p>(i) $\frac{1}{r(r+1)}$</p> <p>(ii) $1 - \frac{1}{n+1}$</p> <p>(iii) $S_{\infty} = 1$ $\frac{1}{n+1}$</p>	<p>B1</p> <p>M1 M1 A1</p> <p>B1 ft M1 A1 c.a.o.</p>	<p>1 Show correct process to obtain given result</p> <p>3 Express terms as differences using (i) Show that terms cancel Obtain correct answer, must be n not any other letter</p> <p>3 State correct value of sum to infinity Ft their (ii) Use sum to infinity – their (ii)</p> <p>7 Obtain correct answer a.e.f.</p>
6	<p>(i) (a) $\alpha + \beta + \gamma = 3, \alpha\beta + \beta\gamma + \gamma\alpha = 2$</p> <p>(b)</p> <p>$\alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha)$ $= 9 - 4 = 5$</p> <p>$\frac{3}{u^3} - \frac{9}{u^2} + \frac{6}{u} + 2 = 0$</p> <p>(ii) (a) $2u^3 + 6u^2 - 9u + 3 = 0$</p> <p>$\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} = -3$</p> <p>(b)</p>	<p>B1 B1</p> <p>M1</p> <p>A1 ft</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1 ft</p>	<p>2 State correct values</p> <p>2 State or imply the result and use their values</p> <p>2 Obtain correct answer</p> <p>2 Use given substitution to obtain an equation</p> <p>2 Obtain correct answer</p> <p>8 Required expression is related to new cubic stated or implied -(their “b” / their “a”)</p>

7	(i) $a(a - 12) + 32$ (ii) $\det \mathbf{M} = 12$ non-singular (iii) <i>EITHER</i> <i>OR</i>	M1 M1 A1 M1 A1ft B1 M1 A1 M1 A1 A1	3 2 3 8	Show correct expansion process Show evaluation of a 2 x 2 determinant Obtain correct answer a.e.f. Substitute $a = 2$ in their determinant Obtain correct answer and state a consistent conclusion $\det \mathbf{M} = 0$ so non-unique solutions Attempt to solve and obtain 2 inconsistent equations Deduce that there are no solutions Substitute $a = 4$ and attempt to solve Obtain 2 correct inconsistent equations Deduce no solutions
8	(i) Circle, centre (3, 0), y -axis a tangent at origin Straight line, through (1, 0) with +ve slope In 1 st quadrant only (ii) Inside circle, below line, above x -axis	B1B1 B1 B1 B1 B1 B2ft	6 2 8	Sketch showing correct features N.B. treat 2 diagrams as MR Sketch showing correct region SR: B1ft for any 2 correct features

<p>9</p>	<p>(i) $\begin{pmatrix} \sqrt{2} & 0 \\ 0 & \sqrt{2} \end{pmatrix}$</p> <p>(ii) Rotation (centre O), 45°, clockwise</p> <p>(iii)</p> <p>(iv) $\begin{pmatrix} 0 \\ 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} \begin{pmatrix} 2 \\ 0 \end{pmatrix}$</p> <p>(v) $\det C = 2$ area of square has been doubled</p>	<p>B1</p> <p>B1B1B1</p> <p>B1</p> <p>M1 A1</p> <p>B1</p> <p>B1</p>	<p>1</p> <p>3</p> <p>1</p> <p>2</p> <p></p> <p>2</p> <p>9</p>	<p>Correct matrix</p> <p>Sensible alternatives OK, must be a single transformation</p> <p>Matrix multiplication or combination of transformations</p> <p>For at least two correct images For correct diagram</p> <p>State correct value</p> <p>State correct relation a.e.f.</p>
<p>10</p>	<p>(i)</p> <p>$x^2 - y^2 = 16$ and $xy = 15$</p> <p>$\pm(5 + 3i)$</p> <p>(ii)</p> <p>$z = 1 \pm \sqrt{16 + 30i}$</p> <p>$6 + 3i, -4 - 3i$</p>	<p>M1</p> <p>A1A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>M1*</p> <p>A1 *M1dep A1 A1ft</p>	<p></p> <p></p> <p></p> <p></p> <p>6</p> <p></p> <p>5</p> <p>11</p>	<p>Attempt to equate real and imaginary parts of $(x + iy)^2$ and $16 + 30i$</p> <p>Obtain each result</p> <p>Eliminate to obtain a quadratic in x^2 or y^2</p> <p>Solve to obtain $x = (\pm) 5$ or $y = (\pm) 3$</p> <p>Obtain correct answers as complex numbers</p> <p>Use quadratic formula or complete the square</p> <p>Simplify to this stage Use answers from (i) Obtain correct answers</p>

Mark Scheme 4726

June 2007

1	<p>Correct formula with correct r Rewrite as $a + b\cos 6\theta$ Integrate their expression correctly Get $\frac{1}{3}\pi$</p>	<p>M1 Allow $r^2 = 2 \sin^2 3\theta$ M1 $a, b \neq 0$ A1√ From $a + b\cos 6\theta$ A1 cao</p>
2	<p>(i) Expand to $\sin 2x \cos \frac{1}{4}\pi + \cos 2x \sin \frac{1}{4}\pi$ Clearly replace $\cos \frac{1}{4}\pi, \sin \frac{1}{4}\pi$ to A.G.</p> <p>(ii) Attempt to expand $\cos 2x$ Attempt to expand $\sin 2x$ Get $\frac{1}{2}\sqrt{2} (1 + 2x - 2x^2 - 4x^3/3)$</p>	<p>B1 B1 M1 Allow $1 - 2x^2/2$ M1 Allow $2x - 2x^3/3$ A1 Four correct unsimplified terms in any order; allow bracket; AEEF SR Reasonable attempt at $f^n(0)$ for $n=0$ to 3 M1 Attempt to replace their values in Maclaurin M1 Get correct answer only A1</p>
3	<p>(i) Express as $A/(x-1) + (Bx+C)/(x^2+9)$ Equate (x^2+9x) to $A(x^2+9) + (Bx+C)(x-1)$ Sub. for x or equate coeff. Get $A=1, B=0, C=9$</p> <p>(ii) Get $A \ln(x-1)$ Get $C/3 \tan^{-1}(x/3)$</p>	<p>M1 Allow $C=0$ here M1√ May imply above line; on their P.F. M1 Must lead to at least 3 coeff.; allow cover-up method for A A1 cao from correct method B1√ On their A B1√ On their C; condone no constant; ignore any $B \neq 0$</p>
4	<p>(i) Reasonable attempt at product rule Derive or quote diff. of $\cos^{-1}x$ Get $-x^2(1-x^2)^{-1/2} + (1-x^2)^{1/2} + (1-x^2)^{-1/2}$ Tidy to $2(1-x^2)^{1/2}$</p> <p>(ii) Write down integral from (i) Use limits correctly Tidy to $\frac{1}{2}\pi$</p>	<p>M1 Two terms seen M1 Allow + A1 A1 cao B1 On any $k\sqrt{1-x^2}$ M1 In any reasonable integral A1 SR Reasonable sub. B1 Replace for new variable and attempt to integrate (ignore limits) M1 Clearly get $\frac{1}{2}\pi$ A1</p>

5	(i)	Attempt at parts on $\int 1 (\ln x)^n dx$	M1	Two terms seen	
		Get $x (\ln x)^n - \int^n (\ln x)^{n-1} dx$	A1		
		Put in limits correctly in line above	M1		
		Clearly get A.G.	A1	$\ln e = 1, \ln 1 = 0$ seen or implied	
	(ii)	Attempt I_3 to I_2 as $I_3 = e - 3I_2$	M1		
		Continue sequence in terms of I_n	A1	$I_2 = e - 2I_1$ and/or $I_1 = e - I_0$	
		Attempt I_2 or I_1	M1	$(I_0 = e - 1, I_1 = 1)$	
		Get $6 - 2e$	A1	cao	
6	(i)	Area under graph ($= \int 1/x^2 dx, 1$ to $n+1$)	B1	Sum (total) seen or implied eg diagram; accept areas (of rectangles)	
		< Sum of rectangles (from 1 to n)			
		Area of each rectangle = Width x Height	B1	Some evidence of area worked out – seen or implied	
		$= 1 \times 1/x^2$			
	(ii)	Indication of new set of rectangles	B1		
		Similarly, area under graph from 1 to n	B1	Sum (total) seen or implied	
		> sum of areas of rectangles from 2 to n	B1	Diagram; use of left-shift of previous areas	
	(iii)	Show complete integrations of RHS, using correct, different limits	Correct answer, using limits, to one integral	M1	Reasonable attempt at $\int x^{-2} dx$
			Add 1 to their second integral to get complete series	A1	
			Clearly arrive at A.G.	M1	
				A1	
	(iv)	Get one limit	B1	Quotable	
Get both 1 and 2			B1	Quotable; limits only required	

7	(i)	Use correct definition of cosh or sinh x	B1	Seen anywhere in (i)
		Attempt to mult. their cosh/sinh	M1	
		Correctly mult. out and tidy	A1√	
		Clearly arrive at A.G.	A1	Accept e^{x-y} and e^{y-x}
	(ii)	Get $\cosh(x - y) = 1$	M1	
		Get or imply $(x - y) = 0$ to A.G.	A1	
	(iii)	Use $\cosh^2 x = 9$ or $\sinh^2 x = 8$	B1	
		Attempt to solve $\cosh x = 3$ (not -3)	M1	$x = \ln(3 + \sqrt{8})$ from formulae book
		or $\sinh x = \pm\sqrt{8}$ (allow $+\sqrt{8}$ or $-\sqrt{8}$ only)		or from basic cosh definition
		Get at least one x solution correct	A1	
		Get both solutions correct, x and y	A1	$x, y = \ln(3 \pm 2\sqrt{2})$; AEEF
			SR	Attempt $\tanh = \sinh/\cosh$ B1
				Get $\tanh x = \pm\sqrt{8}/3$ (+ or -) M1
				Get at least one sol. correct A1
				Get both solutions correct A1
			SR	Use exponential definition B1
				Get quadratic in e^x or e^{2x} M1
				Solve for one correct x A1
				Get both solutions, x and y A1
8	(i)	$x_2 = 0.1890$	B1	
		$x_3 = 0.2087$	B1√	From their x_1 (or any other correct)
		$x_4 = 0.2050$	B1√	Get at least two others correct,
		$x_5 = 0.2057$		all to a minimum of 4 d.p.
		$x_6 = 0.2055$		
		$x_7 (= x_8) = 0.2056$ (to x_7 minimum)		
		$\alpha = 0.2056$	B1	cao; answer may be retrieved despite some errors
	(ii)	Attempt to diff. $f(x)$	M1	$k/(2+x)^3$
		Use α to show $f'(\alpha) \neq 0$	A1√	Clearly seen, or explain $k/(2+x)^3 \neq 0$ as $k \neq 0$; allow ± 0.1864
			SR	Translate $y=1/x^2$ M1
				State/show $y=1/x^2$ has no TP A1
	(iii)	$\delta_3 = -0.0037$ (allow -0.004)	B1√	Allow \pm , from their x_4 and x_3
	(iv)	Develop from $\delta_{10} = f'(\alpha)$ δ_9 etc. to get δ_i		
		or quote $\delta_{10} = \delta f'(\alpha)^7$	M1	Or any δ_i eg use $\delta_9 = x_{10} - x_9$
		Use their δ_i and $f'(\alpha)$	M1	
		Get 0.000000028	A1	Or answer that rounds to ± 0.00000003

- 9 (i) Quote $x = a$ B1
 Attempt to divide out M1 Allow M1 for $y=x$ here; allow
 A1 $(x-a) + k/(x-a)$ seen or implied
 Get $y = x - a$ A1 Must be equations
- (ii) Attempt at quad. in x ($=0$) M1
 Use $b^2 - 4ac \geq 0$ for real x M1 Allow $>$
 Get $y^2 + 4a^2 \geq 0$ A1
 State/show their quad. is always >0 B1 Allow \geq
- (iii) B1√ Two asymptotes from (i) (need not
 be labelled)
 B1 Both crossing points
 B1√ Approaches – correct shape
 SR Attempt diff. by quotient/product
 rule M1
 Get quadratic in x for $dy/dx = 0$
 and note $b^2 - 4ac < 0$ A1
 Consider horizontal asymptotes B1
 Fully justify answer B1


Mark Scheme 4727

June 2007

1 (i) $zz^* = re^{i\theta} \cdot re^{-i\theta} = r^2 = z ^2$	B1 1	For verifying result AG
(ii) Circle Centre $0 (+0i)$ OR $(0, 0)$ OR O , radius 3	B1 B1 2 3	For stating circle For stating correct centre and radius
2 EITHER: $(\mathbf{r} \Rightarrow) [3+t, 1+4t, -2+2t]$ $8(3+t) - 7(1+4t) + 10(-2+2t) = 7$ $\Rightarrow (0t) + (-3) = 7 \Rightarrow$ contradiction l is parallel to Π , no intersection	M1 M1 A1 A1 B1 5	For parametric form of l seen or implied For substituting into plane equation For obtaining a contradiction For conclusion from correct working
OR: $[1, 4, 2] \cdot [8, -7, 10] = 0$ $\Rightarrow l$ is parallel to Π $(3, 1, -2)$ into Π $\Rightarrow 24 - 7 - 20 \neq 7$ l is parallel to Π , no intersection	M1 A1 M1 A1 B1	For finding scalar product of direction vectors For correct conclusion For substituting point into plane equation For obtaining a contradiction For conclusion from correct working
OR: Solve $\frac{x-3}{1} = \frac{y-1}{4} = \frac{z+2}{2}$ and $8x - 7y + 10z = 7$ eg $y - 2z = 3$, $2y - 2 = 4z + 8$ eg $4z + 4 = 4z + 8$ l is parallel to Π , no intersection	M1 A1 M1 A1 B1 5	For eliminating one variable For eliminating another variable For obtaining a contradiction For conclusion from correct working
3 Aux. equation $m^2 - 6m + 8 (= 0)$ $m = 2, 4$ CF $(y \Rightarrow) Ae^{2x} + Be^{4x}$ PI $(y \Rightarrow) Ce^{3x}$ $9C - 18C + 8C = 1 \Rightarrow C = -1$ GS $y = Ae^{2x} + Be^{4x} - e^{3x}$	M1 A1 A1√ M1 A1 B1√ 6 6	For auxiliary equation seen For correct roots For correct CF. f.t. from their m For stating and substituting PI of correct form For correct value of C For GS. f.t. from their CF + PI with 2 arbitrary constants in CF and none in PI

4 (i) $q(st) = qp = s$ $(qs)t = tt = s$	B1 B1 2	For obtaining s For obtaining s
(ii) METHOD 1 Closed: see table Identity = r Inverses: $p^{-1} = s, q^{-1} = t, (r^{-1} = r),$ $s^{-1} = p, t^{-1} = q$	B1 B1 M1 A1 4	For stating closure with reason For stating identity r For checking for inverses For stating inverses <i>OR</i> For giving sufficient explanation to justify each element has an inverse eg r occurs once in each row and/or column
METHOD 2 Identity = r eg $p^2 = t, p^3 = q, p^4 = s$ $\Rightarrow p^5 = r$, so p is a generator	B1 M1 A1 A1	For stating identity r For attempting to establish a generator $\neq r$ For showing powers of p (<i>OR</i> q, s or t) are different elements of the set For concluding p^5 (<i>OR</i> q^5, s^5 or t^5) = r
(iii) e, d, d^2, d^3, d^4	B2 2 8	For stating all elements AEF eg d^{-1}, d^{-2}, dd

5 (i) $(\cos 6\theta =) \operatorname{Re}(c + is)^6$ $(\cos 6\theta =) c^6 - 15c^4s^2 + 15c^2s^4 - s^6$ $(\cos 6\theta =)$ $c^6 - 15c^4(1 - c^2) + 15c^2(1 - c^2)^2 - (1 - c^2)^3$ $(\cos 6\theta =) 32c^6 - 48c^4 + 18c^2 - 1$	M1 A1 M1 A1 4	For expanding (real part of) $(c + is)^6$ at least 4 terms and 1 evaluated binomial coefficient needed For correct expansion For using $s^2 = 1 - c^2$ For correct result AG
(ii) $64x^6 - 96x^4 + 36x^2 - 3 = 0 \Rightarrow \cos 6\theta = \frac{1}{2}$ $\Rightarrow (\theta =) \frac{1}{18}\pi, \frac{5}{18}\pi, \frac{7}{18}\pi$ etc. $\cos 6\theta = \frac{1}{2}$ has multiple roots largest x requires smallest θ \Rightarrow largest positive root is $\cos \frac{1}{18}\pi$	M1 A1 M1 A1 4 8	For obtaining a numerical value of $\cos 6\theta$ For any correct solution of $\cos 6\theta = \frac{1}{2}$ For stating or implying at least 2 values of θ For identifying $\cos \frac{1}{18}\pi$ AEF as the largest positive root from a list of 3 positive roots <i>OR</i> from general solution <i>OR</i> from consideration of the cosine function

<p>6 (i) $\mathbf{n} = l_1 \times l_2$</p> $\mathbf{n} = [2, -1, 1] \times [4, 3, 2]$ $\mathbf{n} = k[-1, 0, 2]$ $[3, 4, -1] \cdot k[-1, 0, 2] = -5k$ $\mathbf{r} \cdot [-1, 0, 2] = -5$	B1 M1* A1 M1 (*dep) A1 5	For stating or implying in (i) or (ii) that \mathbf{n} is perpendicular to l_1 and l_2 For finding vector product of direction vectors For correct vector (any k) For substituting a point of l_1 into $\mathbf{r} \cdot \mathbf{n}$ For obtaining correct p . AEF in this form
<p>(ii) $[5, 1, 1] \cdot k[-1, 0, 2] = -3k$</p> $\mathbf{r} \cdot [-1, 0, 2] = -3$	M1 A1√ 2	For using same \mathbf{n} and substituting a point of l_2 For obtaining correct p . AEF in this form f.t. on incorrect \mathbf{n}
<p>(iii) $d = \frac{ -5+3 }{\sqrt{5}}$ OR $d = \frac{ [2, -3, 2] \cdot [-1, 0, 2] }{\sqrt{5}}$</p> $\text{OR } d \text{ from } (5, 1, 1) \text{ to } \Pi_1 = \frac{ 5(-1)+1(0)+1(2)+5 }{\sqrt{5}}$ $\text{OR } d \text{ from } (3, 4, -1) \text{ to } \Pi_2 = \frac{ 3(-1)+4(0)-1(2)+3 }{\sqrt{5}}$ $\text{OR } [3-t, 4, -1+2t] \cdot [-1, 0, 2] = -3 \Rightarrow t = \frac{2}{5}$ $\text{OR } [5-t, 1, 1+2t] \cdot [-1, 0, 2] = -5 \Rightarrow t = -\frac{2}{5}$ $d = \frac{2}{\sqrt{5}} = \frac{2\sqrt{5}}{5} = 0.894427\dots$	M1 A1√ 2	For using a distance formula from their equations Allow omission of $ $ OR For finding intersection of \mathbf{n}_1 and Π_2 or \mathbf{n}_2 and Π_1 For correct distance AEF f.t. on incorrect \mathbf{n}
<p>(iv) d is the shortest OR perpendicular distance between l_1 and l_2</p>	B1 1 10	For correct statement
<p>7 (i) $(z - e^{i\phi})(z - e^{-i\phi}) \equiv z^2 - (2z \frac{(e^{i\phi} + e^{-i\phi})}{2}) + 1$</p> $\equiv z^2 - (2 \cos \phi)z + 1$	B1 1	For correct justification AG
<p>(ii) $z = e^{\frac{2k\pi i}{7}}$</p> <p>for $k = 0, 1, 2, 3, 4, 5, 6$ OR $0, \pm 1, \pm 2, \pm 3$</p> 	B1 B1 B1 B1 4	For general form OR any one non-real root For other roots specified ($k=0$ may be seen in any form, eg $1, e^0, e^{2\pi i}$) For answers in form $\cos \theta + i \sin \theta$ allow maximum B1 B0 For any 7 points equally spaced round unit circle (circumference need not be shown) For 1 point on $+^{\text{ve}}$ real axis, and other points in correct quadrants
<p>(iii) $(z^7 - 1) = (z - 1)(z - e^{\frac{2\pi i}{7}})(z - e^{\frac{4\pi i}{7}})$</p> $(z - e^{\frac{6\pi i}{7}})(z - e^{\frac{2\pi i}{7}})(z - e^{\frac{4\pi i}{7}})(z - e^{\frac{6\pi i}{7}})$ $= (z - e^{\frac{2\pi i}{7}})(z - e^{\frac{2\pi i}{7}}) \times (z - e^{\frac{4\pi i}{7}})(z - e^{\frac{4\pi i}{7}})$ $(z - e^{\frac{6\pi i}{7}})(z - e^{\frac{6\pi i}{7}}) \times (z - 1)$ $= (z^2 - (2 \cos \frac{2}{7} \pi)z + 1) \times$ $(z^2 - (2 \cos \frac{4}{7} \pi)z + 1) \times (z^2 - (2 \cos \frac{6}{7} \pi)z + 1) \times$ $(z - 1)$	M1 M1 B1 A1 A1 5 10	For using linear factors from (ii), seen or implied For identifying at least one pair of complex conjugate factors For linear factor seen For any one quadratic factor seen For the other 2 quadratic factors and expression written as product of 4 factors

<p>8 (i) Integrating factor $e^{\int \tan x (dx)}$ $= e^{-\ln \cos x}$ $= (\cos x)^{-1}$ OR $\sec x$ $\Rightarrow \frac{d}{dx}(y(\cos x)^{-1}) = \cos^2 x$ $y(\cos x)^{-1} = \int \frac{1}{2}(1 + \cos 2x) (dx)$ $y(\cos x)^{-1} = \frac{1}{2}x + \frac{1}{4}\sin 2x (+c)$ $y = \left(\frac{1}{2}x + \frac{1}{4}\sin 2x + c\right)\cos x$</p>	<p>B1 M1 A1 B1√ M1 M1 A1 A1 8</p>	<p>For correct IF For integrating to ln form For correct simplified IF AEF For $\frac{d}{dx}(y \cdot \text{their IF}) = \cos^2 x \cdot \text{their IF}$ For integrating LHS For attempting to use $\cos 2x$ formula OR parts for $\int \cos^2 x dx$ For correct integration both sides AEF For correct general solution AEF</p>
<p>(ii) $2 = \left(\frac{1}{2}\pi + c\right) \cdot -1 \Rightarrow c = -2 - \frac{1}{2}\pi$ $y = \left(\frac{1}{2}x + \frac{1}{4}\sin 2x - 2 - \frac{1}{2}\pi\right)\cos x$</p>	<p>M1 A1 2 10</p>	<p>For substituting $(\pi, 2)$ into their GS and solve for c For correct solution AEF</p>
<p>9 (i) $3^n \times 3^m = 3^{n+m}$, $n + m \in \mathbb{Z}$ $(3^p \times 3^q) \times 3^r = (3^{p+q}) \times 3^r = 3^{p+q+r}$ $= 3^p \times (3^{q+r}) = 3^p \times (3^q \times 3^r) \Rightarrow$ associativity Identity is 3^0 Inverse is 3^{-n} $3^n \times 3^m = 3^{n+m} = 3^{m+n} = 3^m \times 3^n \Rightarrow$ commutativity</p>	<p>B1 M1 A1 B1 B1 B1 6</p>	<p>For showing closure For considering 3 distinct elements, seen bracketed 2+1 or 1+2 For correct justification of associativity For stating identity. Allow 1 For stating inverse For showing commutativity</p>
<p>(ii) (a) $3^{2n} \times 3^{2m} = 3^{2n+2m} (= 3^{2(n+m)})$ Identity, inverse OK</p>	<p>B1* B1 (*dep) 2</p>	<p>For showing closure For stating other two properties satisfied and hence a subgroup</p>
<p>(b) For 3^{-n}, $-n \notin$ subset</p>	<p>M1 A1 2</p>	<p>For considering inverse For justification of not being a subgroup 3^{-n} must be seen here or in (i)</p>
<p>(c) EITHER: eg $3^{1^2} \times 3^{2^2} = 3^5$ $\neq 3^{r^2} \Rightarrow$ not a subgroup OR: $3^{n^2} \times 3^{m^2} = 3^{n^2+m^2}$ $\neq 3^{r^2}$ eg $1^2 + 2^2 = 5 \Rightarrow$ not a subgroup</p>	<p>M1 A1 2 M1 A1 12</p>	<p>For attempting to find a specific counter-example of closure For a correct counter-example and statement that it is not a subgroup For considering closure in general For explaining why $n^2 + m^2 \neq r^2$ in general and statement that it is not a subgroup</p>

Mark Scheme 4728
June 2007

1(i)	$X = 5$ $Y = 12$	B1 B1 [2]	$X = -5$ B0. Both may be seen/implied in (ii) No evidence for which value is X or Y available from (ii) award B1 for the pair of values 5 and 12 irrespective of order
(ii)	$R^2 = 5^2 + 12^2$ Magnitude is 13 N $\tan \theta = 12/5$ Angle is 67.4°	M1 A1 M1 A1 [4]	For using $R^2 = X^2 + Y^2$ Allow 13 from $X = -5$ For using correct angle in a trig expression SR: $p = 14.9$ and $Q = 11.4$ giving $R = 13 \pm 0.1$ B2, Angle = 67.5 ± 0.5 B2
2(i)	$250 + \frac{1}{2}(290 - 250)$ $t = 270$	M1 A1 [2]	Use of the ratio 12:12 (may be implied), or $v = u + at$
(ii)	$\frac{1}{2} \times 40 \times 12 + 210 \times 12 + \frac{1}{2} \times 20 \times 12 -$ $\frac{1}{2} \times 20 \times 12$ or $\frac{1}{2} \times 40 \times 12 + 210 \times 12$ or $\frac{1}{2} \times (210 + 250) \times 12$ etc Displacement is 2760m	M1 M1 A1 [3]	The idea that area represents displacement Correct <u>structure</u> , ie triangle1 + rectangle2 + triangle3 - triangle4 with triangle3 = triangle4 , triangle1 + rectangle2, trapezium1&2, etc
(iii)	appropriate <u>structure</u> , ie triangle + rectangle + triangle + triangle , triangle + rectangle + 2triangle, etc Distance is 3000m	M1 A1 [2]	All terms positive Treat candidate doing (ii) in (iii) and (iii) in (ii) as a mis-read.
3(i)	$R + T \sin 72^\circ = 50g$	M1 A1 [2]	An equation with R, T and 50 in linear combination. $R + 0.951T = 50g$
(ii)	$T = 50g / \sin 72^\circ$ $T = 515$ (AG) $T = mg$ $m = 52.6$	M1 A1 B1 B1 [4]	Using $R = 0$ (may be implied) and $T \sin 72^\circ = 50(g)$ Or better Accept 52.5
(iii)	$X = T \cos 72^\circ$ $X = 159$	B1 B1 [2]	Implied by correct answer Or better
4(i)	<i>In Q4 right to left may be used as the positive sense throughout.</i> $0.18 \times 2 - 3m = 0$ $m = 0.12$	M1 A1 A1 [3]	For using Momentum 'before' is zero 3 marks possible if g included consistently
(iia)	Momentum after $= -0.18 \times 1.5 + 1.5m$ $0.18 \times 2 - 3m = -0.18 \times 1.5 + 1.5m$ $m = 0.14$	B1 M1 A1 [3]	For using conservation of momentum 3 marks possible if g included consistently
(iib)	$0.18 \times 2 - 3m$ $= (0.18 + m)1.5$ $m = 0.02$ $0.18 \times 2 - 3m = -(0.18 + m)1.5$ $m = 0.42$	B1ft B1 B1ft B1 [4]	ft wrong momentum 'before' 0 marks if g included

5(iii) OR (without finding exactly where or when)	M1	Using $v^2 = u^2 + 2as$ for P <i>and</i> for Q, $a = +/-g$, cv(5.6), different expressions for s.
cont $v_p^2 = 8.4^2 - 2g(s+/-2)$ and		Correct sign for g, cv(5.6), (s+/-2) used only once cao. Verbal explanation essential
$v_Q^2 = 5.6^2 - 2g[(s+/-2)]$ $v_p^2 = v_Q^2$ for all values of s so that the speeds are always the same at the same heights.	A1	Using $v = u+at$ for P <i>and</i> for Q, $a = +/-g$
$0 = 8.4 - gt$ and $0 = 5.6 - gt$	A1	Correct sign for g, correct choice for velocity of zero, cv(5.6)
$t_p = 6/7$ and $t_Q = 4/7$ means there is a time interval when Q has started to descend but P is still rising, and there will be a position where they have the same height but are moving in opposite directions.	M1	cao. Verbal explanation essential
	A1	
6(i) $v = 0.004t^3 - 0.12t^2 + 1.2t$ $v(10) = 4 - 12 + 12 = 4\text{ms}^{-1}$ (AG)	M1	For differentiating s
	A1	Condone the inclusion of +c
	A1	Correct formula for v (no +c) and t=10
	[3]	stated sufficient
6(ii) $v = 0.8t - 0.04t^2$ (+ C)	M1	For integrating a
$8 - 4 + C = 4$	A1	
$v = 0.8 \times 20 - 0.04 \times 20^2$ (+ C)	M1*	Only for using $v(10) = 4$ to find C
$v(20) = 16 - 16 = 0$ (AG)	M1	
	DA1	Dependant on M1*
	[5]	
6(iii) $S = 0.4t^2 - 0.04t^3/3$ (+K)	M1	For integrating v
$s(10) = 10 - 40 + 60 = 30$	A1	Accept $0.4t^2 - 0.013t^3$ (+ ct +K, must be linear)
$40 - 40/3 + K = 30 \rightarrow K = 10/3$	B1	
	M1	For using $S(10) = 30$ to find K
$S(20) = 160 - 320/3 + 10/3 = 56.7\text{m}$	A1	Not if S includes ct term
OR	A1	Accept 56.6 to 56.7, Adding 30 subsequently is not isw, hence B0
$s(10) = 10 - 40 + 60 = 30$	[6]	
$S = 0.4t^2 - 0.04t^3/3$	B1	
$S(20) - S(10) = 26.6, 26.7$	M1	For integrating v
displacement is 56.7m	A1	Accept $0.4t^2 - 0.013t^3$ (+ ct +K, must be linear)
	M1	Using limits of 10 and 20 (limits 0, 10 M0A0B0)
	A1	For 53.3 - 26.7 or better (Note $S(10) = 26.7$ is fortuitously correct M0A0B0)
	B1	Accept 56.6 to 56.7

7(i)	$R = 1.5g\cos 21^\circ$ Frictional force is 10.98N (AG)	B1 M1 A1 [3]	For using $F = \mu R$ Note $1.2g\cos 21^\circ = 10.98$ fortuitously, B0M0A0
(ii)	$T + 1.5g\sin 21^\circ - 10.98 = 1.5a$ $1.2g - T = 1.2a$	M1 A2 A2 [5]	For obtaining an N2L equation relating to the block in which F, T, m and a are in linear combination or For obtaining an N2L equation relating to the object in which T, m and a are in linear combination -A1 for each error to zero -A1 for each error to zero Error is a wrong/omitted term, failure to substitute a numerical value for a letter (excluding g), excess terms. Minimise error count.
(iii)	$T - 1.5a = 5.71$ and $1.2a + T = 11.76$ $a = 2.24$ (AG)	M1 A1 [2]	For solving the simultaneous equations in T and a for a. Evidence of solving needed
(iva)	$v^2 = 2 \times 2.24 \times 2$ Speed of the block is 2.99ms^{-1}	M1 A1 [2]	For using $v^2 = 2as$ with cv (a) or 2.24 Accept 3
(ivb)	$a = -3.81$ $v^2 = 2.99^2 + 2 \times (-3.81) \times 0.8$ Speed of the block is 1.69ms^{-1}	M1 A1 M1 A1 [4]	For using $T = 0$ to find a For using $v^2 = u^2 + 2as$ with cv(2.99) and $s = 2.8 - 2$ and any value for a Accept art 1.7 from correct work

Mark Scheme 4729

June 2007

1	$40 \cos 35^\circ$	B1	
	$WD = 40 \cos 35^\circ \times 100$	M1	
	3280 J	A1 3	ignore units 3
2	$0 = 12 \sin 27^\circ t - 4.9t^2$ any correct.	M1	or $R = u^2 \sin 2\theta / g$ (B2)
	$t = 1.11$method for total time	A1	correct formula only
	$R = 12 \cos 27^\circ \times t$	M1	$12^2 \times \sin 54^\circ / 9.8$ sub in values
	11.9	A1 4	11.9 4
3 (i)	$WD = \frac{1}{2} \times 250 \times 150^2 - \frac{1}{2} \times 250 \times 100^2$	M1	
	1 560 000	A1	1 562 500
	$450\,000 = 1\,560\,000/t$	M1	
	3.47	A1 4	
(ii)	$F = 450\,000/120$	M1	
	3750	A1	
	$3750 = 250a$	M1	
	15 ms^{-2}	A1 4	8
4 (i)	$x = 7t$	B1	
	$y = 21t - 4.9t^2$	M1	or $-g/2$
		A1	
	$y = 21 \cdot x/7 - 4.9 x^2/49$	M1	
	$y = 3x - x^2/10$	A1 5	AG
(ii)	$-25 = 3x - x^2/10$ (must be -25)	M1	or method for total time (5.26)
	solving quadratic	M1	or 7 x total time
	36.8 m	A1 3	8
5(i)	$\frac{1}{2} \cdot 70 \cdot 4^2$	M1	
	560 J	A1 2	
(ii)	$70 \times 9.8 \times 6$	M1	
	4120	A1 2	4116
(iii)	60d	B1	
	$8000 = 560 + 4120 + 60d$	M1	4 terms
		A1 ✓	✓ their KE and PE
	55.4 m	A1 4	8

6 (i)	$5\cos 30^\circ = 0.3 \times 9.8 + S\cos 60^\circ$	M1	res. vertically (3 parts with comps)
		A1	
	2.78 N	A1	3
(ii)	$r = 0.4\sin 30^\circ = 0.2$	B1	may be on diagram
	$5\sin 30^\circ + S\sin 60^\circ = 0.3 \times 0.2 \times \omega^2$	M1	res. horizontally (3 parts with comps)
	9.04 rads^{-1}	A1	3
(iii)	$v = 0.2 \times 9.04$	M1	or previous v via mv^2/r
	$\text{KE} = \frac{1}{2} \times 0.3 \times (0.2 \times 9.04)^2$	M1	
	0.491 J or 0.49	A1 ✓	3 ✓ their $\omega^2 \times 0.006$ 9

7 (i)	$1.8 = -0.3 + 3m$	M1	
	$m = 0.7$	A1	2 AG
(ii)	$e = 4/6$	M1	accept 2/6 for M1
	2/3	A1	2 accept 0.67
(iii)	$\pm 3f$	B1	
	$1/3 \odot f (\ominus 1)$	B1	2
(iv)	$I = 3f \times 0.7 - - 3 \times 0.7$	M1	ok for only one minus sign for M1
		A1	
	$I = 2.1(f + 1)$	A1	3 aef 2 marks only for $-2.1(f + 1)$
(v)	$0.3 + 6.3/4 = 0.3a + 0.7b$	M1	can be $-0.7b$
	$3a + 7b = 18.75$	A1	*
	$2/3 = (a - b)/5/4$	M1	allow $e=3/4$ or their e for M1
	$3a - 3b = 5/2$	A1	*
	solve	M1	
	$a = 2.5$	A1	(2.46) allow \pm (59/24)
	$b = 1.6$	A1	7 (1.625) allow \pm (13/8) 16

8 (i)	com of hemisphere 0.3 from O	B1	or 0.5 from base
	com of cylinder $h/2$ from O	B1	
	$0.6 \times 45 = 40 \times 0.5 + (0.8 + h/2) \times 5$ or	M1	or $40 \times 0.3 - 5xh/2 = 45 \times 0.2$
	$45(h + 0.2) = 5h/2 + 40(h + 0.3)$	A1	or $5(0.2 + h/2) = 40 \times 0.1$
	$27 = 20 + (0.8 + h/2) \times 5$	M1	solving
	$h = 1.2$	A1	6 AG
(ii)	1.2 T	B1	
	0.8 F	B1	
	$0.8F = 1.2T$	M1	
	$F = 3T/2$	A1	4 aef
(iii)	$F + T\cos 30^\circ$	B1	or $45 \times 0.8 \sin 30^\circ$
	$45\sin 30^\circ$ must be involved in res.	B1	$T \times (1.2 + 0.8\cos 30^\circ)$
	resolving parallel to the slope	M1	mom. about point of contact
	$F + T\cos 30^\circ = 45\sin 30^\circ$ aef	A1	$45 \times 0.8\sin 30^\circ = T(1.2 + 0.8\cos 30^\circ)$
	$T = 9.51$	A1	
	$F = 14.3$	A1	6 16
or	$T + F\cos 30^\circ = R\sin 30^\circ$	B1	res. horizontally
(iii)	$R\cos 30^\circ + F\sin 30^\circ = 45$	B1	res. vertically
	$\tan 30^\circ = (T + F\cos 30^\circ)/(45 - F\sin 30^\circ)$	M1	eliminating R

Mark Scheme 4730

June 2007

1	(i) $[\omega = 2\pi/6.1 = 1.03]$	M1	For using $T = 2\pi/\omega$
	Speed is 3.09ms^{-1}	M1 A1	For using $v_{\max} = a\omega$
	(ii)	M1	For using $v^2 = \omega^2(A^2 - x^2)$ or for using $v = A\omega \cos \omega t$ and $x = A \sin \omega t$
	$2.5^2 = 1.03^2(3^2 - x^2)$ or $x = 3\sin(1.03 \times 0.60996\dots)$ Distance is 1.76m	A1ft A1	ft incorrect ω
2	[Magnitudes 0.6, 0.057×7 , 0.057×10]	M1	For triangle with magnitudes shown
	For magnitudes of 2 sides correctly marked	A1	
	For magnitudes of all 3 sides correctly marked	A1	
		M1	For attempting to find angle (α) opposite to the side of magnitude 0.057×7
		M1	For correct use of the cosine rule or equivalent
	$0.399^2 = 0.57^2 + 0.6^2 - 2 \times 0.57 \times 0.6 \cos \alpha$ Angle is 140°	A1ft A1	$(180 - 39.8)^\circ$
2	ALTERNATIVE METHOD		
		M1	For using $I = \Delta mv$ parallel to the initial direction of motion or parallel to the impulse
	$-0.6 \cos \alpha = 0.057 \times 7 \cos \beta - 0.057 \times 10$ or $0.6 = 0.057 \times 10 \cos \alpha + 0.057 \times 7 \cos \gamma$	A1	
		M1	For using $I = \Delta mv$ perpendicular to the initial direction of motion or perpendicular to the impulse
	$0.6 \sin \alpha = 0.057 \times 7 \sin \beta$ or $0.057 \times 10 \sin \alpha = 0.057 \times 7 \sin \gamma$	A1	
	$0.399^2 = (0.57 - 0.6 \cos \alpha)^2 + (0.6 \sin \alpha)^2$ or $0.399^2 = (0.6 - 0.57 \cos \alpha)^2 + (0.057 \sin \alpha)^2$ Angle is 140°	M1 A1ft A1	For eliminating β *or γ

3	(i) $[0.2v \, dv/dx = -0.4v^2]$	M1		For using Newton's second law with $a = v \, dv/dx$
	$(1/v) \, dv/dx = -2$	A1	2	AG
	(ii) $[\int (1/v) \, dv = \int -2 \, dx]$	M1		For separating variables and attempting to integrate
	$\ln v = -2x \quad (+C)$	A1		
	$[\ln v = -2x + \ln u]$	M1		For using $v(0) = u$
	$v = ue^{-2x}$	A1	4	AG
	(iii) $[\int e^{2x} \, dx = \int u \, dt]$	M1		For using $v = dx/dt$ and separating variables
	$e^{2x}/2 = ut \quad (+C)$	A1		
	$[e^{2x}/2 = ut + 1/2]$	M1		For using $x(0) = 0$
	$u = 6.70$	A1	4	Accept $(e^4 - 1)/8$
ALTERNATIVE METHOD FOR PART (iii)				
	$[\int \frac{1}{v^2} \, dv = -2 \int dt \rightarrow -1/v = -2t + A, \text{ and}]$	M1		For using $a = dv/dt$, separating variables, attempting to integrate and using $v(0) = u$
	$A = -1/u]$	M1		For substituting $v = ue^{-2x}$
	$-e^{2x}/u = -2t - 1/u$	A1		
	$u = 6.70$	A1	4	Accept $(e^4 - 1)/8$
4	$y = 15 \sin \alpha \quad (=12)$	B1		
	$[4(15 \cos \alpha) - 3 \times 12 = 4a + 3b]$	M1		For using principle of conservation of momentum in the direction of l.o.c.
	Equation complete with not more than one error	A1		
	$4a + 3b = 0$	A1		
		M1		For using NEL in the direction of l.o.c.
	$0.5(15 \cos \alpha + 12) = b - a$	A1		
	$[a = -4.5, b = 6]$	M1		For solving for a and b
	$[\text{Speed} = \sqrt{(-4.5)^2 + 12^2},$	M1		For correct method for speed or direction of A
	$\text{Direction } \tan^{-1}(12/(-4.50))]$			
	Speed of A is 12.8ms^{-1} and direction is 111° anticlockwise from 'i' direction	A1		Direction may be stated in any form, including $\theta = 69^\circ$ with θ clearly and appropriately indicated
Speed of B is 6ms^{-1} to the right	A1	10	Depends on first three M marks	

5	(i)	M1	For taking moments of forces on BC about B
	$80 \times 0.7 \cos 60^\circ = 1.4T$	A1	
	Tension is 20N	A1	
	$[X = 20 \cos 30^\circ]$	M1	For resolving forces horizontally
	Horizontal component is 17.3N	A1ft	ft $X = T \cos 30^\circ$
	$[Y = 80 - 20 \sin 30^\circ]$	M1	For resolving forces vertically
	Vertical component is 70N	A1ft	ft $Y = 80 - T \sin 30^\circ$
	(ii)	M1	For taking moments of forces on AB, or on ABC, about A
	$17.3 \times 1.4 \sin \alpha = (80 \times 0.7 + 70 \times 1.4) \cos \alpha$ or	A1ft	
	$80 \times 0.7 \cos \alpha + 80(1.4 \cos \alpha + 0.7 \cos 60^\circ) =$		
	$20 \cos 60^\circ(1.4 \cos \alpha + 1.4 \cos 60^\circ) +$		
	$20 \sin 60^\circ(1.4 \sin \alpha + 1.4 \sin 60^\circ)$		
	$[\tan \alpha = (\frac{1}{2} 80 + 70)/17.3 = 11/\sqrt{3}]$	M1	For obtaining a numerical expression for $\tan \alpha$
	$\alpha = 81.1^\circ$	A1	4

ALTERNATIVE METHOD FOR PART (i)			
		M1	For taking moments of forces on BC about B
	$H \times 1.4 \sin 60^\circ + V \times 1.4 \cos 60^\circ = 80 \times 0.7 \cos 60^\circ$	A1	Where H and V are components of T
		M1	For using $H = V\sqrt{3}$ and solving simultaneous equations
	Tension is 20N	A1	
	Horizontal component is 17.3N	B1ft	ft value of H used to find T
	$[Y = 80 - V]$	M1	For resolving forces vertically
	Vertical component is 70N	A1ft	ft value of V used to find T

6	(i) [T = 2058x/5.25] 2058x/5.25 = 80 x 9.8 (x = 2) OP = 7.25m	M1 A1 A1	3	For using $T = \lambda x/L$ AG From 5.25 + 2
	(ii) Initial PE = (80 + 80)g(5) (= 7840) or (80 + 80)gX used in energy equation Initial KE = $\frac{1}{2}(80 + 80)3.5^2$ (= 980) [Initial EE = 2058x ² /(2x5.25) (= 784), Final EE = 2058x ⁷ /(2x5.25) (= 9604), or 2058(X + 2) ² /(2x5.25)] [Initial energy = 7840 + 980 + 784, final energy = 9604 or 1568X + 980 + 784 = 196(X ² + 4X + 4) → 196X ² - 784X - 980 = 0]	B1 B1 M1 M1		For using $EE = \lambda x^2/2L$ For attempting to verify compatibility with the principle of conservation of energy, or using the principle and solving for X
	Initial energy = final energy or X = 5 → P&Q just reach the net	A1	5	AG
	(iii) [PE gain = 80g(7.25 + 5)] PE gain = 9604 PE gain = EE at net level → P just reaches O	M1 A1 A1	3	For finding PE gain from net level to O AG
(iv) For any one of 'light rope', 'no air resistance', 'no energy lost in rope' For any other of the above	B1 B1	2		

FIRST ALTERNATIVE METHOD FOR PART (ii)				
[160g - 2058x/5.25 = 160v dv/dx]	M1			For using Newton's second law with $a = v dv/dx$, separating the variables and attempting to integrate
$v^2/2 = gx - 1.225x^2$ (+ C)	A1			Any correct form
C = -8.575	M1 A1			For using $v(2) = 3.5$
[$v(7)^2/2 = 68.6 - 60.025 - 8.575 = 0$ → P&Q just reach the net]	A1	5	AG	

SECOND ALTERNATIVE METHOD FOR PART (ii)				
$\ddot{x} = g - 2.45x$ (= -2.45(x - 4))	B1 M1			For using $n^2 = 2.45$ and $v^2 = n^2(A^2 - (x - 4)^2)$
$3.5^2 = 2.45(A^2 - (-2)^2)$ (A = 3)	A1			
[(4 - 2) + 3]	M1			For using 'distance travelled downwards by P and Q = distance to new equilibrium position + A
distance travelled downwards by P and Q = 5 → P&Q just reach the net	A1	5	AG	



7	(i) [a = 0.7 ² /0.4] For not more than one error in $T - 0.8g\cos 60^\circ = 0.8 \times 0.7^2 / 0.4$ Above equation complete and correct Tension is 4.9N	M1 A1 A1 A1	For using a = v ² /r 4
	(ii) $\frac{1}{2} 0.8v^2 =$ $\frac{1}{2} 0.8(0.7)^2 + 0.8g0.4 - 0.8g0.4 \cos 60^\circ$ $(2.1 - 0)/7 = 2u$ Q's initial speed is 0.15ms ⁻¹	M1 A1 M1 A1	For using the principle of conservation of energy (v = 2.1) For using NEL 4 AG
	(iii) $(m)0.4 \ddot{\theta} = -(m)g \sin \theta$ $[0.4 \ddot{\theta} \approx -g\theta]$ $[\frac{1}{2} m0.15^2 = mg0.4(1 - \cos \theta_{\max})]$ $\rightarrow \theta_{\max} = 4.34^\circ (0.0758\text{rad})]$	M1 A1 M1 M1	For using Newton's second law transversely *Allow m = 0.8 (or any other numerical value) For using $\sin \theta \approx \theta$ For using the principle of conservation of energy to find θ_{\max}
	θ_{\max} small justifies $0.4 \ddot{\theta} \approx -g\theta$, and this implies SHM	A1	5
	(iv) [T = 2π / √24.5 = 1.269..] [√24.5 t = π]	M1	For using T = 2π/n or for solving either sin nt = 0 (non-zero t) (considering displacement) or cos nt = -1 (considering velocity)
	Time interval is 0.635s	A1ft	2 From t = ½ T

Mark Scheme 4731

June 2007

1 (i)	Using $\theta = \omega_0 t + \frac{1}{2} \alpha t^2$, $56 = 0 + \frac{1}{2} \alpha \times 8^2$ $\alpha = 1.75 \text{ rad s}^{-2}$	M1 A1	2
	(ii) Using $\omega_1^2 = \omega_0^2 + 2\alpha\theta$, $36^2 = 20^2 + 2 \times 1.75\theta$ $\theta = 256 \text{ rad}$	M1 A1 ft	ft is $448 \div \alpha$ 2
2	Volume is $\int_0^a \pi(4a^2 - x^2) dx = \pi \left[4a^2 x - \frac{1}{3} x^3 \right]_0^a$ $= \frac{11}{3} \pi a^3$	M1 A1	π may be omitted throughout (Limits not required)
	$\int_0^a \pi x(4a^2 - x^2) dx$ $= \pi \left[2a^2 x^2 - \frac{1}{4} x^4 \right]_0^a$ $= \frac{7}{4} \pi a^4$ $\bar{x} = \frac{\frac{7}{4} \pi a^4}{\frac{11}{3} \pi a^3}$ $= \frac{21}{44} a$	M1 A1 A1 M1 A1	(Limits not required) for $\frac{\int x y^2 dx}{\int y^2 dx}$
3 (i)	$I = 6.2 + 2.8 = 9.0 \text{ kg m}^2$	B1	1
	(ii) WD against frictional couple is $L \times \frac{1}{2} \pi$ Loss of PE is $6 \times 9.8 \times 1.3$ (= 76.44) Gain of KE is $\frac{1}{2} \times 9.0 \times 2.4^2$ (= 25.92) By work-energy principle, $L \times \frac{1}{2} \pi = 76.44 - 25.92$ $L = 32.2 \text{ Nm}$	B1 B1 B1 ft M1 A1	Equation involving WD, KE and PE 5 Accept 32.1 to 32.2
	(iii) $6 \times 9.8 \times 0.8 - L = I \alpha$ $\alpha = 1.65 \text{ rad s}^{-2}$	M1 A1 ft A1	Moments equation 3

<p>4 (i)</p>	<p>MI of elemental disc about a diameter is</p> $\frac{1}{4} \left(\frac{M}{3a} \delta x \right) a^2$ <p>MI of elemental disc about AB is</p> $\frac{1}{4} \left(\frac{M}{3a} \delta x \right) a^2 + \left(\frac{M}{3a} \delta x \right) x^2$ $I = \frac{M}{3a} \int_0^{3a} \left(\frac{1}{4} a^2 + x^2 \right) dx$ $= \frac{M}{3a} \left[\frac{1}{4} a^2 x + \frac{1}{3} x^3 \right]_0^{3a}$ $= \frac{M}{3a} \left(\frac{3}{4} a^3 + 9a^3 \right)$ $= M \left(\frac{1}{4} a^2 + 3a^2 \right)$ $= \frac{13}{4} M a^2$	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1 (ag)</p> <p>7</p>	<p>$\frac{M}{3a}$ may be $\rho \pi a^2$ throughout (condone use of $\rho = 1$)</p> <p>Using parallel axes rule (can award A1 for $\frac{1}{4} m a^2 + m x^2$)</p> <p>Integrating MI of disc about AB Correct integral expression for I</p> <p>Obtaining an expression for I in terms of M and a Dependent on previous M1</p>
<p>(ii)</p>	<p>Period is $2\pi \sqrt{\frac{I}{Mgh}}$</p> $= 2\pi \sqrt{\frac{\frac{13}{4} M a^2}{Mg \frac{3}{2} a}}$ $= 2\pi \sqrt{\frac{13a}{6g}}$	<p>M1</p> <p>A1</p> <p>A1</p> <p>3</p>	<p>or $-Mgh \sin \theta = I \ddot{\theta}$</p>

<p>5 (i)</p>	 $\frac{\sin \theta}{12} = \frac{\sin 115}{16}$ $\theta = 42.8^\circ$ <p>Bearing of v_B is 007.2°</p> $\frac{u}{\sin 22.2} = \frac{16}{\sin 115}$ $u = 6.66$ <p>Time taken is $\frac{2400}{6.664} = 360$ s</p>	<p>M1 A1 M1 A1 M1 A1 M1*A1 ft 8</p>	<p>Relative velocity on bearing 050 Correct velocity diagram; or $\begin{pmatrix} u \sin 50 \\ u \cos 50 \end{pmatrix} = \begin{pmatrix} 16 \sin \alpha \\ 16 \cos \alpha \end{pmatrix} - \begin{pmatrix} 12 \sin 345 \\ 12 \cos 345 \end{pmatrix}$ or eliminating u (or α) or obtaining equation for u (or α) <i>For equations in α and t</i> <i>M1*M1A1 for equations</i> <i>M1 for eliminating t (or α)</i> <i>A1 for $\alpha = 7.2$</i> <i>M1A1 ft for equation for t (or α)</i> <i>A1 cao for $t = 360$</i></p>
<p>(ii)</p>	 $\cos \phi = \frac{10}{12}$ $\phi = 33.6^\circ$ <p>Bearing of v_B is 018.6°</p>	<p>M1 A1 M1 A1 4</p>	<p>Relative velocity perpendicular to v_B Correct velocity diagram For alternative methods: M2 for a completely correct method A2 for 018.6 (give A1 for a correct relevant angle)</p>

6 (i)	$I = \frac{1}{3}ma^2 + m\left(\frac{1}{3}a\right)^2$ $= \frac{4}{9}ma^2$ $mg\left(\frac{1}{3}a \cos \theta\right) = I \alpha$ $\alpha = \frac{\frac{1}{3}mga \cos \theta}{\frac{4}{9}ma^2} = \frac{3g \cos \theta}{4a}$	M1 A1 M1 A1 (ag)	Using parallel axes rule 4
(ii)	By conservation of energy, $\frac{1}{2}I \omega^2 = mg\left(\frac{1}{3}a \sin \theta\right)$ $\frac{2}{9}ma^2 \omega^2 = \frac{1}{3}mg a \sin \theta$ $\omega = \sqrt{\frac{3g \sin \theta}{2a}}$	M1 A1 ft A1	<i>Condone</i> $\omega^2 = \frac{3g \sin \theta}{2a}$ 3
OR	$\omega \frac{d\omega}{d\theta} = \frac{3g \cos \theta}{4a}$ $\frac{1}{2}\omega^2 = \int \frac{3g \cos \theta}{4a} d\theta$ $= \frac{3g \sin \theta}{4a} (+ C)$ $\omega = \sqrt{\frac{3g \sin \theta}{2a}}$	M1 A1 A1	
(iii)	Acceleration parallel to rod is $\left(\frac{1}{3}a\right)\omega^2$ $F - mg \sin \theta = m\left(\frac{1}{3}a\right)\omega^2$ $F - mg \sin \theta = \frac{1}{2}mg \sin \theta$ $F = \frac{3}{2}mg \sin \theta$	B1 M1 A1	Radial equation with 3 terms
	Acceleration perpendicular to rod is $\left(\frac{1}{3}a\right)\alpha$ $mg \cos \theta - R = m\left(\frac{1}{3}a\right)\alpha$ $mg \cos \theta - R = \frac{1}{4}mg \cos \theta$ $R = \frac{3}{4}mg \cos \theta$	B1 ft M1 A1	ft is $r \alpha$ with r the same as before Transverse equation with 3 terms 6
OR	$R\left(\frac{1}{3}a\right) = I_G \alpha$ $R\left(\frac{1}{3}a\right) = \left(\frac{1}{3}ma^2\right)\left(\frac{3g \cos \theta}{4a}\right)$ $R = \frac{3}{4}mg \cos \theta$	M1 A1 A1	Must use I_G
(iv)	On the point of slipping, $F = \mu R$ $\frac{3}{2}mg \sin \theta = \mu\left(\frac{3}{4}mg \cos \theta\right)$ $\tan \theta = \frac{1}{2}\mu$	M1 A1 (ag)	Correctly obtained 2 <i>Dependent on 6 marks earned in (iii)</i>

7 (i)	$\text{GPE} = (-) mg(2a \cos \theta) \cos \theta$ $\text{EPE} = \frac{1}{2} \frac{mg}{2a} (AR - a)^2$ $= \frac{1}{2} \frac{mg}{2a} (2a \cos \theta - a)^2$ $V = \frac{1}{4} mga(2 \cos \theta - 1)^2 - 2mga \cos^2 \theta$ $= mga(\cos^2 \theta - \cos \theta + \frac{1}{4} - 2 \cos^2 \theta)$ $= mga(\frac{1}{4} - \cos \theta - \cos^2 \theta)$	B1 M1 A1 A1 (ag)	or $(-) mg(a + a \cos 2\theta)$ 4
(ii)	$\frac{dV}{d\theta} = mga(\sin \theta + 2 \cos \theta \sin \theta)$ $= mga \sin \theta (1 + 2 \cos \theta)$ Equilibrium when $\frac{dV}{d\theta} = 0$ ie when $\theta = 0$	B1 M1 A1 (ag)	3
(iii)	KE is $\frac{1}{2} m(2a \dot{\theta})^2$ $2ma^2 \dot{\theta}^2 + V = \text{constant}$ Differentiating with respect to t , $4ma^2 \dot{\theta} \ddot{\theta} + \frac{dV}{d\theta} \dot{\theta} = 0$ $4ma^2 \dot{\theta} \ddot{\theta} + mga \sin \theta (1 + 2 \cos \theta) \dot{\theta} = 0$ $\ddot{\theta} = -\frac{g}{4a} \sin \theta (1 + 2 \cos \theta)$	B1 M1 M1 A1 ft A1 (ag)	(can award this M1 if no KE term) SR B2 (replacing the last 3 marks) for the given result correctly obtained by differentiating w.r.t. θ 5
(iv)	When θ is small, $\sin \theta \approx \theta$, $\cos \theta \approx 1$ $\ddot{\theta} \approx -\frac{g}{4a} \theta (1 + 2) = -\frac{3g}{4a} \theta$ Period is $2\pi \sqrt{\frac{4a}{3g}}$	M1 A1 A1	3

Mark Scheme 4732

June 2007


Note: "3 sfs" means an answer which is equal to, or rounds to, the given answer. If such an answer is seen and then later rounded, apply ISW.

1	$(0 \times 0.1) + 1 \times 0.2 + 2 \times 0.3 + 3 \times 0.4$ $= 2(.0)$ $(0^2 \times 0.1) + 1 \times 0.2 + 2^2 \times 0.3 + 3^2 \times 0.4 (= 5)$ $- 2^2$ $= 1$	M1 A1 M1 M1 A1 5	≥ 2 non-zero terms correct eg $\div 4$: M0 ≥ 2 non-zero terms correct $\div 4$: M0 Indep, ft their μ . Dep +ve result $(-2)^2 \times 0.1 + (-1)^2 \times 0.2 + 0^2 \times 0.3 + 1^2 \times 0.4$: M2 ≥ 2 non-0 correct: M1 $\div 4$: M0
Total		5	
2	UK Fr Ru Po Ca 1 2 3 4 5 or 5 4 3 2 1 4 3 1 5 2 2 3 5 1 4 Σd^2 $(= 24)$ $r_s = 1 - \frac{6 \times "24"}{5 \times (5^2 - 1)}$ $= -\frac{1}{5}$ or -0.2	M1 A1 M1 M1 A1 5	Consistent attempt rank other judge <div style="border: 1px solid black; padding: 5px; display: inline-block;"> RCFUP 3 5 2 1 4 3 1 4 5 2 1 2 3 4 5 5 4 3 2 1 </div> All 5 d^2 attempted & added. Dep ranks att'd Dep 2 nd M1 <div style="border: 1px solid black; padding: 5px; display: inline-block;"> $\frac{43 - 15^2/5}{\sqrt{((55 - 15^2/5)(55 - 15^2/5))}}$ Corr sub in ≥ 2 S's M1 All correct: M1 </div>
Total		5	
3i	${}^{15}C_7$ or ${}^{15!}/7!8!$ 6435	M1 A1 2	
ii	${}^6C_3 \times {}^9C_4$ or ${}^{6!}/3!3! \times {}^{9!}/4!5!$ 2520	M1 A1 2	Alone except allow $\div {}^{15}C_7$ Or ${}^6P_3 \times {}^9P_4$ or ${}^{6!}/3! \times {}^{9!}/5!$ Allow $\div {}^{15}P_7$ NB not ${}^{6!}/3! \times {}^{9!}/4!$ 362880
Total		4	
4ia	$\frac{1}{3}$ oe	B1 1	<div style="border: 1px solid black; padding: 5px; display: inline-block;"> B\leftrightarrowW MR: max (a)B0(b)M1M1(c)B1M1 </div>
b	P(BB) + P(WB) attempted $= \frac{4}{10} \times \frac{3}{9} + \frac{6}{10} \times \frac{4}{9}$ or $\frac{2}{15} + \frac{4}{15}$ $= \frac{2}{5}$ oe	M1 M1 A1 3	Or $\frac{4}{10} \times \frac{3}{9}$ OR $\frac{6}{10} \times \frac{4}{9}$ correct NB $\frac{4}{10} \times \frac{4}{10} + \frac{6}{10} \times \frac{4}{10} = \frac{2}{5}$: M1M0A0
c	Denoms 9 & 8 seen or implied $\frac{3}{9} \times \frac{2}{8} + \frac{6}{9} \times \frac{3}{8}$ $= \frac{1}{3}$ oe	B1 M1 A1 3	Or $\frac{2}{15}$ as numerator Or $\frac{2/15}{4/10}$ <div style="border: 1px solid black; padding: 5px; display: inline-block;"> Or $\frac{\frac{4}{10} \times \frac{6}{9} \times \frac{3}{8} + \frac{4}{10} \times \frac{3}{9} \times \frac{2}{8}}{\frac{4}{10} \times \frac{6}{9} \times \frac{3}{8} + \frac{4}{10} \times \frac{3}{9} \times \frac{2}{8}}$ </div> May not see wking
ii	P(Blue) not constant or discs not indep, so no	B1 1	Prob changes as discs removed Limit to no. of discs. Fixed no. of discs Discs will run out Context essential: "disc" or "blue" NOT fixed no. of trials NOT because without repl Ignore extra
Total		8	

5i	1991 100 000 to 110 000	B1 ind B1 ind 2	Or fewer in 2001 Allow digits 100 to 110
5ia	Median = 29 to 29.9 Quartiles 33 to 34, 24.5 to 26 = 7.5 to 9.5 140 to 155 23 to 26.3%	B1 M1 A1 M1 A1 5	Or one correct quartile and subtr NOT from incorrect wking ×1000, but allow without Rounded to 1 dp or integer 73.7 to 77% : SC1
b	Older Median (or ave) greater } % older mothers greater oe} % younger mothers less oe}	B1 B1 B1 3	Or 1991 younger Any two Or 1991 steeper so more younger: B2 NOT mean gter Ignore extra
Total		10	

6ia	Correct subst in \geq two S formulae $\frac{767 - \frac{60 \times 72}{8}}{\sqrt{(1148 - \frac{60^2}{8})(810 - \frac{72^2}{8})}}$ or $\frac{227}{\sqrt{698}\sqrt{162}}$ $= 0.675$ (3 sfs)	M1 M1 A1 3	Any version All correct. Or $\frac{767-8x7.5x9}{\sqrt{((1148-8x7.5^2)(810-8x9^2))}}$ or correct substn in any correct formula for r
b	1 y always increases with x or ranks same oe	B1 B1 2	+ve grad thro'out. Increase in steps. Same order. Both ascending order Perfect RANK corr'n Ignore extra NOT Increasing proportionately
iiia	Closer to 1, or increases because nearer to st line	B1 B1 2	Corr'n stronger. Fewer outliers. "They" are outliers Ignore extra
b	None, or remains at 1 Because y still increasing with x oe	B1 B1 2	Σd^2 still 0. Still same order. Ignore extra NOT differences still the same. NOT ft (i)(b)
iii	13.8 to 14.0	B1 1	
iv	(iii) or graph or diag or my est Takes account of curve	B1 B1 2	Must be clear which est. Can be implied. "This est" probably \Rightarrow using equn of line Straight line is not good fit. Not linear. Corr'n not strong.
Total		12	
7i	P(contains voucher) constant oe Packets indep oe	B1 B1 2	Context essential NOT vouchers indep
ii	0.9857 or 0.986 (3 sfs)	B2 2	B1 for 0.9456 or 0.946 or 0.997(2) or for 7 terms correct, allow one omit or extra NOT $1 - 0.9857 = 0.0143$ (see (iii))
iii	$(1 - 0.9857)$ $= 0.014(3)$ (2 sfs)	B1ft 1	Allow 1- their (ii) correctly calc'd
iv	B(11, 0.25) or 6 in 11 wks stated or impl ${}^{11}C_6 \times 0.75^5 \times 0.25^6$ (= 0.0267663) $P(6 \text{ from } 11) \times 0.25$ $= 0.00669$ or 6.69×10^{-3} (3 sfs)	B1 M1 M1 A1 4	or $0.75^a \times 0.25^b$ ($a + b = 11$) or ${}^{11}C_6$ dep B1
Total		9	

8i	$\sqrt{0.04} (= 0.2)$ $(1 - \text{their } \sqrt{0.04})^2$ $= 0.64$	M1 M1 A1 3	
ii	1 - p seen $2p(1 - p) = 0.42$ or $p(1 - p) = 0.21$ oe $2p^2 - 2p + 0.42 (= 0)$ or $p^2 - p + 0.21 (= 0)$ $\frac{2 \pm \sqrt{((-2)^2 - 4 \times 0.42)}}{2 \times 2}$ or $\frac{1 \pm \sqrt{((-1)^2 - 4 \times 0.21)}}{2 \times 1}$ or $(p - 0.7)(p - 0.3) = 0$ or $(10p - 7)(10p - 3) = 0$ $p = 0.7$ or 0.3	B1 M1 M1 M1 A1 5	$2pq = 0.42$ or $pq = 0.21$ Allow $pq = 0.42$ or opp signs, correct terms any order ($= 0$) oe Correct Dep B1M1M1 Any corr subst'n or fact'n Omit 2 in 2 nd line: max B1M1M0M0A0 One corr ans with no or inadeq wking: SC1 eg $0.6 \times 0.7 = 0.42 \Rightarrow p = 0.7$ or 0.6 $p^2 + 2pq + q^2 = 1$ B1 $p^2 + q^2 = 0.58$ } $p = 0.21/q$ } $p^4 - 0.58p^2 + 0.0441 = 0$ M1 corr subst'n or fact'n M1 1 - p seen B1 $2p(1 - p) = 0.42$ or $p(1 - p) = 0.21$ M1 $p^2 - p = -0.21$ $p^2 - p + 0.25 = -0.21 + 0.25$ oe } M1 OR $(p - 0.5)^2 - 0.25 = -0.21$ oe } $(p - 0.5)^2 = 0.04$ M1 $(p - 0.5) = \pm 0.2$ $p = 0.3$ or 0.7 A1
Total		8	
9ia	$1 / \frac{1}{5}$ $= 5$	M1 A1 2	
b	$(\frac{4}{5})^3 \times \frac{1}{5}$ $= \frac{64}{625}$ or 0.102 (3 sfs)	M1 A1 2	
c	$(\frac{4}{5})^4$ $= \frac{256}{625}$ or a.r.t 0.410 (3 sfs) or 0.41	M1 A1 2	or $1 - (\frac{1}{5} + \frac{4}{5} \times \frac{1}{5} + (\frac{4}{5})^2 \times \frac{1}{5} + (\frac{4}{5})^3 \times \frac{1}{5})$ NOT $1 - (\frac{4}{5})^4$
iiia	$P(Y=1) = p, P(Y=3) = q^2 p, P(Y=5) = q^4 p$		$P(Y=1) + P(Y=3) + P(Y=5) = p + q^2 p + q^4 p$ $p, p(1 - p)^2, p(1 - p)^4$ q^{-1}, q^{-3}, q^{-5} or any of these with $1 - p$ instead of q "Always q to even power $\times p$ " Either associate each term with relevant prob Or give indication of how terms derived \geq two terms
b	Recog that c.r. $= q^2$ or $(1 - p)^2$ $S_\infty = \frac{p}{1 - q^2}$ or $\frac{p}{1 - (1 - p)^2}$ $P(\text{odd}) = \frac{1 - q}{1 - q^2}$ $= \frac{1 - q}{(1 - q)(1 + q)}$ Must see this step for A1 $(= \frac{1}{1 + q})$ AG	M1 M1 M1 A1 4	or eg $r = \frac{q^2 p}{p}$ $(= \frac{p}{2p - p^2}) = \frac{p}{p(2 - p)}$ $(= \frac{1}{2 - p}) = \frac{1}{2 - (1 - q)}$




Total		11	
-------	--	----	--

Mark Scheme 4733

June 2007

1	(i)	$\hat{\mu} = 4830.0/100 = 48.3$ $249509.16/100 - (\text{their } \bar{x}^2)$ $\times 100/99$ $= 163.84$	B1 M1 M1 A1	4	48.3 seen Biased estimate: 162.2016: can get B1M1M0 Multiply by $n/(n-1)$ Answer, 164 or 163.8 or 163.84
	(ii)	No, Central Limit theorem applies, so can assume distribution is normal	B2	2	“No” with statement showing CLT is understood (though CLT does not need to be mentioned) [SR: No with reason that is not wrong: B1]
2		$B(130, 1/40)$ $\approx \text{Po}(3.25)$ $e^{-\lambda} \frac{\lambda^x}{x!}$ $= 0.180$	B1 M1 A1√ M1 A1	5	$B(130, 1/40)$ stated or implied Poisson, <i>or</i> correct N on their $B(n, p)$ Parameter their np , <i>or</i> correct parameter(s)√ Correct formula, or interpolation Answer, 0.18 or a.r.t. 0.180 [SR: N(3.25, 3.17) or N(3.25, 3.25): B1M1A1]
3	(i)	Binomial	B1	1	Binomial stated or implied
	(ii)	Each element equally likely Choices independent	B1 B1	2	All elements, or selections, equally likely stated Choices independent [not just “independent”] [can get B2 even if (i) is wrong]
4	(i)	Two of: Distribution symmetric No substantial truncation Unimodal/Increasingly unlikely further from μ , etc	B1 B1	2	One property Another definitely different property Don’t give both marks for just these two “Bell-shaped”: B1 only unless “no truncation”
	(ii)	Variance $8^2/20$ $z = \frac{47.0 - 50.0}{\sqrt{8^2/20}} = -1.677$ $\Phi(1.677) = 0.9532$	M1 A1 A1 A1	4	Standardise, allow cc, don’t need n Denominator (8 or 8^2 or $\sqrt{8}$) \div (20 or $\sqrt{20}$ or 20^2) z -value, a.r.t. -1.68 or $+1.68$ Answer, a.r.t. 0.953
5	(i)	$H_1: \lambda > 2.5$ or 15	B1	1	$\lambda > 2.5$ or 15, allow μ , don’t need “ H_1 ”
	(ii)	Use parameter 15 $P(> 23)$ $1 - 0.9805 = 0.0195$ or 1.95%	M1 M1 A1	3	$\lambda = 15$ used [N(15, 15) gets this mark only] Find $P(> 23$ or $\geq 23)$, final answer < 0.5 eg 0.0327 or 0.0122 Answer, 1.95% or 2% or 0.0195 or 0.02 [SR: 2-tailed, 3.9% gets 3/3 here]
	(iii)	$P(\leq 23 \lambda = 17) = 0.9367$ $P(\leq 23 \lambda = 18) = 0.8989$ Parameter = 17 $\lambda = 17/6$ or 2.83	M1 A1 M1	3	One of these, or their complement: .9367, .8989, 0.9047, 0.8551, .9317, .8933, .9907, .9805 Parameter 17 [17.1076], needs $P(\leq 23)$, cwo [SR: if insufficient evidence can give B1 for 17] Their parameter $\div 6$ [2.85] [SR: Solve $(23.5 - \lambda)/\sqrt{\lambda} = 1.282$ M1; 18.05 A0]
6	(i)	$H_0: p = 0.19, H_1: p < 0.19$ where p is population proportion $0.81^{20} + 20 \times 0.81^{19} \times 0.19$ $= 0.0841$ Compare 0.1	B2 M1 A1 A1 B1		Correct, B2. One error, B1, but x or \bar{x} or r : B0 Binomial probabilities, allow 1 term only Correct expression [0.0148 + 0.0693] Probability, a.r.t. 0.084 Explicit comparison of “like with like”
	or	Add binomial probs until ans > 0.1 Critical region ≤ 1	A1 B1		$[P(\leq 2) = 0.239]$
		Reject H_0 Significant evidence that proportion of e 's in language is less than 0.19	M1 A1√	8	Correct deduction and method [needs $P(\leq 1)$] Correct conclusion in context [SR: N(3.8, 3.078): B2M1A0B1M0]
	(ii)	Letters not independent	B1	1	Correct modelling assumption, stated in context Allow “random”, “depends on message”, etc

<p>7 (i)</p> 	<p>B1 B1 B1</p> <p>3</p>	<p>Horizontal straight line Positive parabola, symmetric about 0 Completely correct, including correct relationship between two Don't need vertical lines or horizontal lines outside range, but don't give last B1 if horizontal line continues past "±1"</p>
<p>(ii) S is equally likely to take any value in range, T is more likely at extremities</p>	<p>B2</p> <p>2</p>	<p>Correct statement about distributions (not graphs) [Partial statement, or correct description for one only: B1]</p>
<p>(iii)</p> $\int_t^1 \frac{3}{2}x^2 dx = \left[\frac{x^3}{2} \right]_t^1$ <p>$\frac{1}{2}(1 - t^3) = 0.2$ or $\frac{1}{2}(t^3 + 1) = 0.8$ $t^3 = 0.6$ $t = 0.8434$</p>	<p>M1 B1 M1 M1 A1</p> <p>5</p>	<p>Integrate $f(x)$ with limits $(-1, t)$ or $(t, 1)$ [recoverable if t used later] Correct indefinite integral Equate to 0.2, or 0.8 if $[-1, t]$ used Solve cubic equation to find t Answer, in range $[0.843, 0.844]$</p>
<p>8 (i)</p> $\frac{64.2 - 63}{\sqrt{12.25/23}} = 1.644$ <p>$P(z > 1.644)$ $= 0.05$</p>	<p>M1 dep A1 dep M1 A1</p> <p>4</p>	<p>Standardise 64.2 with $\sqrt{12.25}$ $z = 1.644$ or 1.645, must be + Find $\Phi(z)$, answer < 0.5 Answer, a.r.t. 0.05 or 5.0%</p>
<p>(ii) (a)</p> $63 + 1.645 \times \frac{3.5}{\sqrt{50}}$ <p>≥ 63.81</p>	<p>M1 B1 A1</p> <p>3</p>	<p>$63 + 3.5 \times k / \sqrt{50}$, k from Φ^{-1}, not $-$ $k = 1.645$ (allow 1.64, 1.65) Answer, a.r.t. 63.8, allow $>$, \geq, $=$, c.w.o.</p>
<p>(b)</p> $P(< 63.8 \mid \mu = 65)$ $\frac{63.8 - 65}{3.5/\sqrt{50}} = -2.3956$ <p>0.0083</p>	<p>M1 M1 A1 A1</p> <p>4</p>	<p>Use of correct meaning of Type II Standardise their c with $\sqrt{50}$ $z = (\pm) 2.40$ [or -2.424 or -2.404 etc] Answer, a.r.t. 0.008 [eg, 0.00767]</p>
<p>(iii) B better: Type II error smaller (and same Type I error)</p>	<p>B2✓</p> <p>2</p>	<p>This answer: B2. "B because sample bigger": B1. [SR: Partial answer: B1]</p>
<p>9 (a)</p> <p>$np > 5$ and $nq > 5$ $0.75n > 5$ is relevant $n > 20$</p>	<p>M2 A1</p> <p>3</p>	<p>Use either $nq > 5$ or $npq > 5$ [SR: If M0, use $np > 5$, or "n = 20" seen: M1] Final answer $n > 20$ or $n \geq 20$ only</p>
<p>(b) (i)</p> $70.5 - \mu = 1.75\sigma$ $\mu - 46.5 = 2.25\sigma$ <p>Solve simultaneously $\mu = 60$ $\sigma = 6$</p>	<p>M1 A1 B1 M1 A1✓ A1✓</p> <p>6</p>	<p>Standardise once, and equate to Φ^{-1}, \pm cc Standardise twice, signs correct, cc correct Both 1.75 and 2.25 Correct solution method to get one variable μ, a.r.t. 60.0 or ± 154.5 σ, a.r.t. 6.00 [Wrong cc (below): A1 both] [SR: σ^2: M1A0B1M1A1A0]</p>
<p>(ii)</p> $np = 60, npq = 36$ $q = 36/60 = 0.6$ $p = 0.4$ $n = 150$	<p>M1 dep depM1 A1✓ A1✓</p> <p>4</p>	<p>$np = 60$ and $npq = 6^2$ or 6 Solve to get q or p or n $p = 0.4$ ✓ on wrong cc or z $n = 150$ ✓ on wrong cc or z</p>

σ	μ	q	$p (\pm 0.01)$	n
6	60	0.6	0.4	150
6.25	60.062	5	0.6504	171.8
6.25	60.562	5	0.6450	170.6
6.25	59.562	5	0.6558	173.0
6.5	60.125	0.7027	0.2973	202.2
6	59.5	0.6050	0.3950	150.6

Mark Scheme 4734

June 2007

1	$\int_0^1 ax \, dx + \int_1^{\infty} \frac{a}{x^2} \, dx = 1$	M1		For sum of integrals =1
	$[ax^2]_0^1 + \left[-\frac{a}{x} \right]_1^{\infty} = 1$	A1		For second integral.
	$a + a = 1$	A1		For second a
	$a = 1/2$	A1	4	Or from $F(x)$ M1A1 then $F(\infty)=1$ M1, $a=1/2$ A1
2	(i) $\bar{X}_I \square N(5, \frac{0.7^2}{20})$	B1		If no parameters allow in (ii)
	$\bar{X}_E \square N(4.5, \frac{0.5^2}{25})$	B1	2	If 0.7/20, 0.5/25 then B1 for both, with means in (ii)
	(ii) Use $\bar{X}_I - \bar{X}_E \square N(0.5, \sigma^2)$	M1A1		OR $\bar{X}_I - \bar{X}_E - 1 \square N(-0.5, \sigma^2)$
	$\sigma^2 = 0.49/20 + 0.25/25$	B1		cao
	$1 - \Phi([1-0.5]/\sigma)$	M1		RH probability implied. If 0.7, 0.5
	$= 0.0036$ or 0.0035	A1	5	in σ^2 , M1A1B0M1A1 for 0.165
3	Assumes differences form a random sample from a normal distribution.			
	$H_0: \mu = 0, H_1: \mu > 0$	B1	B1	Other letters if defined; or in words
	$\bar{x} = 17.2/12; s^2 = 10.155$ AEF	B1B1		Or $(12/11)(136.36/12 - (17.2/12)^2)$ aef
	EITHER: $t = \frac{\bar{x}}{\sqrt{s^2/12}}$ (+ or -)	M1		With 12 or 9.309/11
	=1.558	A1		Must be positive. Accept 1.56
	1.363 seen	B1		
	1.558 > 1.363, so reject H_0 and accept that there that the readings from the aneroid device overestimate blood pressure on average	B1√		Allow CV of 1.372 or 1.356 evidence Explicit comparison of CV(not - with +) and conclusion in context.
	OR: For critical region or critical value of \bar{x}			
	$1.363\sqrt{(s^2/12)}$	M1B1		B1 for correct t
	Giving 1.25(3)	A1		
	Compare 1.43(3) with 1.25(3)			
	Conclusion in context	B1√	8	



4	(i)	Proper				
		P	F			
	Trial	P 31	11 42	B1		Two correct
	F 5	13 18	B1		Others correct	
		36	24 60		2	

	(ii) (H_0 : Trial results and Proper results are independent.)					
	E-values:	25.2	16.8	M1		One correct. Ft marginals in (i)
		10.8	7.2	A1		All correct
	$\chi^2 = 5.3^2(25.2^{-1} + 10.8^{-1} + 16.8^{-1} + 7.2^{-1})$			M1		Allow two errors
	= 9.289			A1		With Yates' correction
				A1		art 9.29
	Compare correctly with 7.8794			M1		Or 7.88
	There is evidence that results are not independent.			A1 ✓	7	Ft χ^2_{calc}

5	(i) $e^{-\mu} = 0.45$			M1		
	$\mu_G = 0.799 \approx 0.80$ AG			A1	2	0.799 or 0.798 or better seen

	(ii) $\mu_U \approx 1.8$			B1		
	Total, $T \sim \text{Po}(2.6)$			M1		May be implied by answer 0.264
	$P(>3) = 0.264$			A1	3	From table or otherwise

	(iii) $e^{-2.6} 2.6^6 / 6!$			B1		Or 0.318 from table
	$e^{-5.2} 5.2^4 / 4!$			B1		
	Multiply two probabilities			M1		
	Answers rounding to 0.0053 or 0.0054			A1	4	

4734

Mark Scheme

June 2015

6	<p>(i) $\hat{p} = 62/200=0.31$</p> <p>Use $\hat{p}_\alpha \pm z \sqrt{\frac{\hat{p}_\alpha(1-\hat{p}_\alpha)}{200}}$</p> <p>$z=1.96$</p> <p>Correct variance estimate (0.2459,0.3741)</p>	<p>B1</p> <p>M1</p> <p>B1</p> <p>A1√</p> <p>A1</p>	<p>5</p> <p>ae f</p> <p>With 200 or 199</p> <p>Seen ft \hat{p} art (0.246,0.374)</p>
<hr/>			
	<p>(ii)EITHER: Sample proportion has an approximate normal distribution</p> <p>OR: Variance is an estimate</p>	<p>B1</p>	<p>1</p> <p>Not \hat{p} is an estimate, unless variance mentioned</p>
<hr/>			
	<p>(iii) $H_0: p_\alpha = p_\beta, H_1: p_\alpha \neq p_\beta$</p> <p>$\hat{p}=(62+35)/(200+150)$</p>	<p>B1</p>	<p>ae f</p>
<hr/>			
	<p>EITHER: $z=(\pm) \frac{62/200-35/150}{\sqrt{\hat{p}\hat{q}(200^{-1}+150^{-1})}}$</p> <p>=1.586</p> <p>(-1.96 <) 1.586 < 1.96</p> <p>Do not reject H_0- there is insufficient evidence of a difference in proportions.</p>	<p>M1</p> <p>B1√</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>s^2 with, $\hat{p}, 200, 150$ (or 199,149)</p> <p>Evidence of correct variance estimate. Ft \hat{p}</p> <p>Rounding to 1.58 or 1.59</p> <p>Correct comparison with ± 1.96</p> <p>SR: If variance $p_1q_1/n_1+p_2q_2/n_2$ used then: B0M1B0A1(for $z=1.61$ or 1.62)M1A1 Max 4/6.</p>
	<p>OR: $p_{sa} - p_{s\beta} = z s$</p> <p>$s = \sqrt{(0.277 \times 0.723(200^{-1} + 150^{-1}))}$</p> <p>CV of $p_{sa} - p_{s\beta} = 0.0948$ or 0.095</p> <p>Compare $p_{sa} - p_{s\beta} = 0.0767$ with their 0.0948</p> <p>Do not reject H_0 and accept that there is insufficient evidence of a difference in proportions</p>	<p>M1</p> <p>B1√</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>Ft \hat{p}</p> <p>Conditional on $z=1.96$</p>
			6

7	(i) $G(y) = P(Y \leq y)$ $= P(X^2 \geq 1/y)$ [or $P(X > 1/\sqrt{y})$] $= 1 - F(1/\sqrt{y})$ $= \begin{cases} 0 & y \leq 0, \\ y^2 & 0 \leq y \leq 1, \\ 1 & y > 1. \end{cases}$	M1 A1 A1		May be implied by following line Accept strict inequalities
		A1	4	Or $F(x) = P(X \leq x) = P(Y \geq 1/x^2)$ M1 $= 1 - P(Y < 1/x^2)$ A1 $= 1 - G(y)$;etc A1 A1
	(ii) Differentiate their $G(y)$ to obtain $g(y) = 2y$ for $0 < y \leq 1$ AG obtained	M1	A1	2 Only from G correctly
	(iii) $\int_0^1 2y(\sqrt[3]{y}) dy$ $= [6y^{7/3}/7]$ $= 6/7$	M1 B1 A1		Unsimplified, but with limits OR: Find $f(x)$, $\int_1^\infty x^{-2/3} f(x) dx$ M1 $= [4x^{14/3}/(14/3)]; 6/7$ B1A1 OR: Find $H(z)$, $Z = Y^{1/3}$
8	(i) $P(20 \leq y < 25) = \Phi(0) - \Phi(-5/\sqrt{20})$ Multiply by 50 to give 18.41 AG 18.41 for $25 \leq y < 30$ and 6.59 for $y < 20$, $y \geq 30$	M1 A1 A1 A1	4	
	(ii) $H_0: N(25,20)$ fits data $\chi^2 = 3.59^2/6.59 + 8.59^2/18.41 + 6.41^2/18.41$ $+ 1.41^2/6.59$ $= 8.497$ $8.497 > 7.815$ Accept that $N(25,20)$ is not a good fit	B1 M1√ A1 M1 A1		OR $Y \sim N(25,20)$ ft values from (i) art 8.5 5
	(iii) Use $24.91 \pm z\sqrt{(20/50)}$ $z = 2.326$ $(23.44, 26.38)$	B1 A1	M1 3	With $\sqrt{(20/50)}$ art (23.4, 26.4) Must be interval
	(iv) No- Sample size large enough to apply CLT Sample mean will be (approximately) normally distributed whatever the distribution of Y	B1 B1		Refer to large sample size 2 Refer to normality of sample mean

Mark Scheme 4735

June 2007

Statistics 4

1 (i) Use $P(A' \cap B') = 1 - P(A \cup B)$ Use $P(A \cap B) = P(A) + P(B) - P(A \cup B)$ $= c - 0.1$	M1 M1 A1 3	Or $c = 1 - P(A \cup B)$
(ii) $P(B A) = (c - 0.1) / 0.3$ Use $0 \leq p \leq 1$ to obtain $0.1 \leq c \leq 0.4$ AG	B1√ M1 A1 3	Shown clearly
2 $H_0: m_n = m_s, H_1: m_n \neq m_s$ Use Wilcoxon rank sum test 59 64 68 77 80 85 88 90 98 N N N S N S N S S $R_m = 4 + 6 + 8 + 9 = 27$ $40 - 27 = 13$ $W = 13$ Compare correctly with correct CV, !2 Do not reject H_0 . There is no evidence of a difference in the median pulse rates of the two populations.	B1 M1 A1 B1 B1 M1 A1 7	Medians; both hypotheses 'Population medians' if words Rank and identify M0 if normal approx. used Quote critical region or state that $13 > 12$. M0 if $W=27$ Conclusion in context.
3 (i) Use marginal distributions to obtain $E(X) = -0.4, E(Y) = 1.5$ $E(XY) = -0.24 + 0.04 - 0.52 + 0.12$ $Cov(X, Y) = -0.6 + 0.6 = 0$ AG	M1 A1A1 M1 A1 5	
(ii) $P(X = -1 Y = 2) = 0.26 / 0.5 = 0.52$ $P(X = 0 Y = 2) = 0.18 / 0.5 = 0.36$ $P(X = 1 Y = 2) = 0.12$	M1 A1 2	Correct method for any one All correct SR: B1 if no method indicated

4 (i) $H_0: m = 2.70, H_1: m > 2.7$ Subtract 2.70 from each value and count the number of positive signs Obtain 13 Use $B(20, \frac{1}{2})$ to obtain $P(X \geq 13) = 0.1316$ (0.132) Compare correctly with 0.05 Do not reject H_0 . Conclude that there is insufficient evidence to claim that median level of impurity is greater than 2.70	B1 M1 A1 M1 A1 M1 A1 7	In terms of medians Allow just 'medians' here For finding tail probability Or CR: $X \geq 15$ M1A1 Or: $N(10, 5), p=0.132$
(ii) Wilcoxon signed rank test Advantage: More powerful (uses more formation) Disadvantage: This test requires a symmetric population distribution, not required for sign test	B1 B1 B1 3	Smaller P(Type II) Not 'more time taken'
5 (i) $\int_0^{\infty} \frac{1}{(\alpha - 1)!} x^{\alpha-1} e^{-x} dx = 1$, result follows	B1 1	
(ii) $M_X(t) = \int_0^{\infty} \frac{1}{(\alpha - 1)!} x^{\alpha-1} e^{-x} e^{xt} dx$ $= \int_0^{\infty} \frac{1}{(\alpha - 1)!} x^{\alpha-1} e^{-x(1-t)} dx$ $x = u/(1-t), dx = du/(1-t)$ and limits unchanged $= \int_0^{\infty} \frac{1}{(\alpha - 1)!} \frac{u^{\alpha-1}}{(1-t)^{\alpha-1}} \frac{e^{-u}}{1-t} du$ $= \frac{1}{(\alpha - 1)! (1-t)^{\alpha}} \int_0^{\infty} u^{\alpha-1} e^{-u} du$ $= (1-t)^{-\alpha}$ AG	M1 M1 A1 A1 A1 5	Attempt to differentiate With evidence
(iii) EITHER: $M'(t) = \alpha(1-t)^{\alpha-1}$ $M''(t) = \alpha(\alpha+1)(1-t)^{\alpha-2}$ Substitute $t=0$ $E(X) = \alpha$ $\text{Var}(X) = \alpha(\alpha+1) - \alpha^2$ $= \alpha$ OR: $(1-t)^{-\alpha} = 1 + at + \frac{1}{2} \alpha(\alpha+1)t^2 + \dots$ $E(X) = \alpha$ $\text{Var}(X) = E(X^2) - [E(X)]^2$ $= \alpha(\alpha+1) - \alpha^2 ; \alpha$	B1 B1 M1 A1 M1 A1 M1A1 B1 M1 A1A1 6	AEF M0 if t involved

6 (i) $q + pt$	B1 1	Accept $qt^0 + pt^1$
(ii) $(q + pt)^n (= G_S(t))$ Binomial	B1 B1 2	
(iii) $E(S) = G'(1) = np(q+p)$ $= np$ $\text{Var}(S) = G''(1) + G'(1) - [G'(1)]^2$ $= n(n-1)p^2(p+q) + np - n^2p^2$ $= npq$	M1A1 A1 M1 A1 A1 6	AEF, properly obtained
(iv) $(\frac{1}{2} + \frac{1}{2}t)^{10} e^{-(1-t)}$ Find coefficient of t^2 $(\frac{1}{2})^{10} (1 + 10t + \frac{1}{2} \times 10 \times 9t^2)$ $e^{-1} (1 + t + \frac{1}{2}t^2)$ Required coefficient $= e^{-1} 2^{-10} (1/2 + 10 + 45)$ $= 0.0199$	M1 M1 A1 A1 M1 A1 6	Seen May be implied OR: $P(Y=0)P(Z=2) + \dots$ M1, Z is Po(1) M1 Ans: A1A1A1; A1 Not from $e^{-(1-t)} = 1 - (1-t) + (1-t)^2/2$ No more than one term missing--
7 (i) $E(T_1) = 2E(\bar{X}) = 2 \times \frac{1}{2} \theta = \theta$ (So T_1 is an unbiased estimator of θ)	M1A1 2	SR: B1 if $\bar{X} = \int_0^\theta \frac{x}{\theta} d\theta$
(ii) $E(U) = \int_0^\theta \frac{nu^n}{\theta^n} du ; \left[\frac{nu^{n+1}}{\theta^n(n+1)} \right] ; \frac{n\theta}{n+1}$ $E(U^2) = \int_0^\theta \frac{nu^{n+1}}{\theta^n} du ; \frac{n}{n+2} \theta^2$ $\text{Var}(U) = E(U^2) - [E(U)]^2$ $= \frac{n\theta^2}{(n+1)^2(n+2)}$ AG	M1A1A1 M1A1 A1 6	
(iii) $\text{Var}(T_2) = \theta^2/[n(n+2)]$ $\text{Var}(T_1) = 4\text{Var}(X)/n ; \theta^2/3n$ $\text{Var}(T_2)/\text{Var}(T_1)$ $3/(n+2)$ < 1 for $n > 1$ So T_2 is more efficient than T_1	B1 M1A1 M1 M1A1 A1 7	For comparison of var. T_1, T_2 Idea used.

Mark Scheme 4736
June 2007

SOLUTIONS

4736

D1

June 2007

FINAL

1	(i)	Example: $N-P-Q-T-S-R-N$ or: $P-Q-S-P$	B1 1	Any valid cycle (closed and does not repeat vertices, need not be a Hamiltonian cycle)																				
	(ii)	It passes through Y twice	B1 1	Or, it includes a cycle (accept 'loop')																				
	(iii)	5	B1 1																					
	(iv)	A : neither B : semi-Eulerian	B1 B1 2	If graphs are not specified, assume A is first																				
	(v)	A : 2 B : 1	B1 B1 2	If graphs are not specified, assume A is first																				
	(vi)	There are 4 odd nodes (N, P, S and Z) To connect these we must add 2 arcs	M1 A1 2 9	$A: 1, B: 0 \Rightarrow B1, nntv$ Seen or implied For 2																				
2	(i)	$d + f + g = 120$	B1 1	For this equality. Condone an inequality																				
	(ii)	"(Area of) grass is not more than 4 times (area of) decking"	B1 1	Identifying the constraint in words (not just 'grass is less than or equal to 4 times decking' though)																				
	(iii)	$d \leq f$	B1 1	Do <u>not</u> accept $d < f$																				
	(iv)	$g \geq 40$ $\min d = 10$ $\min f = 20$	B1 B1 B1 3	Do <u>not</u> accept $g > 40$ $d \geq 10$ $f \geq 20$																				
	(v)	$5g + 10d + 20f$ or $g + 2d + 4f$	B1 1	Or any positive multiple of this																				
	(vi)	Minimise $g + 2d + 4f$ Subject to $d + f + g = 120$ $g - 4d + s = 0$ $d - f + t = 0$ $g \geq 40,$ and $d \geq 10, f \geq 20, s \geq 0, t \geq 0$	M1 B1 A1 3 10	For a reasonable attempt at setting up the minimisation problem using their expressions For dealing with this slack variable correctly (variables on LHS and constant on RHS) For a completely correct formulation (accept d and $f \geq 0$, or their min values for d, f)																				
3	(i)	<table style="display: inline-table; border: none;"> <tr> <td></td> <td style="text-align: center;">8 6 9 7 5</td> <td style="text-align: center;">Comps</td> <td style="text-align: center;">Swaps</td> </tr> <tr> <td>After 1st pass:</td> <td style="text-align: center;">6 8 9 7 5</td> <td style="text-align: center;">1</td> <td style="text-align: center;">1</td> </tr> <tr> <td>After 2nd pass:</td> <td style="text-align: center;">6 8 9 7 5</td> <td style="text-align: center;">1</td> <td style="text-align: center;">0</td> </tr> <tr> <td>After 3rd pass:</td> <td style="text-align: center;">6 7 8 9 5</td> <td style="text-align: center;">3</td> <td style="text-align: center;">2</td> </tr> <tr> <td>After 4th pass:</td> <td style="text-align: center;">5 6 7 8 9</td> <td style="text-align: center;">4</td> <td style="text-align: center;">4</td> </tr> </table>		8 6 9 7 5	Comps	Swaps	After 1st pass:	6 8 9 7 5	1	1	After 2nd pass:	6 8 9 7 5	1	0	After 3rd pass:	6 7 8 9 5	3	2	After 4th pass:	5 6 7 8 9	4	4	M1 M1 M1 A1 B1 B1 6	Bubble sort or decreasing order loses first 4 marks 1st pass correct 2nd pass correct, follow through from 1st pass 3rd pass correct, follow through from 2nd pass 4th pass correct Counting comparisons for at least three passes Counting swaps for at least three passes
		8 6 9 7 5	Comps	Swaps																				
After 1st pass:	6 8 9 7 5	1	1																					
After 2nd pass:	6 8 9 7 5	1	0																					
After 3rd pass:	6 7 8 9 5	3	2																					
After 4th pass:	5 6 7 8 9	4	4																					
(ii)	Step 1 $A = 8 6 9 7 5$ Step 2 $A = 6 9 7 5$ $X = 8$ Step 3 $A = 9 7 5$ $B = 6$ Step 4 $A = 7 5$ $C = 9$ Step 4 $A = 5$ $B = 6 7$ Step 4 A is empty $B = 6 7 5$ Step 6 $N = 3$ Step 7 $A = 6 7 5 8 9$ Step 8 Display 6 7 5 8 9	M1 M1 M1 M1 A1 5 11	For identifying that $6 \rightarrow B$ or the sublist $\{6\}$ For identifying that $9 \rightarrow C$ or the sublist $\{9\}$ For identifying that $7 \rightarrow B$ For identifying that $5 \rightarrow B$ For the final A list or the display correct																					

4	(i)	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <th><i>P</i></th> <th><i>x</i></th> <th><i>y</i></th> <th><i>s</i></th> <th><i>t</i></th> <th><i>u</i></th> <th></th> </tr> <tr> <td>1</td> <td>-3</td> <td>5</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> </tr> <tr> <td>0</td> <td>1</td> <td>5</td> <td>1</td> <td>0</td> <td>0</td> <td>12</td> </tr> <tr> <td>0</td> <td>1</td> <td>-5</td> <td>0</td> <td>1</td> <td>0</td> <td>10</td> </tr> <tr> <td>0</td> <td>3</td> <td>10</td> <td>0</td> <td>0</td> <td>1</td> <td>45</td> </tr> </table>	<i>P</i>	<i>x</i>	<i>y</i>	<i>s</i>	<i>t</i>	<i>u</i>		1	-3	5	0	0	0	0	0	1	5	1	0	0	12	0	1	-5	0	1	0	10	0	3	10	0	0	1	45	B1	For correct use of three slack variable columns
	<i>P</i>	<i>x</i>	<i>y</i>	<i>s</i>	<i>t</i>	<i>u</i>																																	
	1	-3	5	0	0	0	0																																
	0	1	5	1	0	0	12																																
0	1	-5	0	1	0	10																																	
0	3	10	0	0	1	45																																	
			B1	For $\pm (-3 \ 5)$ in objective row																																			
			B1	For 1 5 12, 1 -5 10 and 3 10 45 in constraint rows																																			
			3																																				
	(ii)	Pivot on second 1 in <i>x</i> column <i>x</i> column has a negative entry in objective row $12 \div 1 = 12$, $10 \div 1 = 10$, $45 \div 3 = 15$ Least non-negative ratio is 10 so pivot on the second 1	B1	For correct pivot choice (cao)																																			
			B1	For 'negative in top row for <i>x</i> ', or equivalent, and a correct explanation of choice of row 'least ratio $10 \div 1$ ' (ft their pivot column)																																			
			2																																				
	(iii)	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <th><i>P</i></th> <th><i>x</i></th> <th><i>y</i></th> <th><i>z</i></th> <th><i>s</i></th> <th><i>t</i></th> <th></th> </tr> <tr> <td>1</td> <td>0</td> <td>-10</td> <td>0</td> <td>3</td> <td>0</td> <td>30</td> </tr> <tr> <td>0</td> <td>0</td> <td>10</td> <td>1</td> <td>-1</td> <td>0</td> <td>2</td> </tr> <tr> <td>0</td> <td>1</td> <td>-5</td> <td>0</td> <td>1</td> <td>0</td> <td>10</td> </tr> <tr> <td>0</td> <td>0</td> <td>25</td> <td>0</td> <td>-3</td> <td>1</td> <td>15</td> </tr> </table> <p>$x = 10, y = 0$ $P = 30$</p>	<i>P</i>	<i>x</i>	<i>y</i>	<i>z</i>	<i>s</i>	<i>t</i>		1	0	-10	0	3	0	30	0	0	10	1	-1	0	2	0	1	-5	0	1	0	10	0	0	25	0	-3	1	15	M1	For correct method evident for objective row
<i>P</i>	<i>x</i>	<i>y</i>	<i>z</i>	<i>s</i>	<i>t</i>																																		
1	0	-10	0	3	0	30																																	
0	0	10	1	-1	0	2																																	
0	1	-5	0	1	0	10																																	
0	0	25	0	-3	1	15																																	
			M1	For a correct method evident for pivot row																																			
			M1	For a correct method evident for other rows																																			
			A1	For correct tableau CAO																																			
			B1	For correct values from their tableau																																			
			B1	For correct value from their tableau																																			
			6																																				
	(iv)	$11 + 5(0.2) = 12$ or $s = 0$ $11 - 5(0.2) = 10$ or $t = 0$ $3(11) + 10(0.2) = 35$ or $u = 10$ so all the constraints are satisfied	B1	For showing (not just stating) that constraints are satisfied																																			
		$P = 3(11) - 5(0.2) = 32$ which is bigger than 30 from (iii)	B1	For calculating 32, or equivalent (eg $3x$ has increased by 3 but $-5y$ has only decreased by 1)																																			
			2																																				
			13																																				

<p>5</p>	<p>(i)</p>	<p>Shortest path from J to B: J G H E B Length of path: 125 metres</p>	<p>M1 M1 A1 B1 B1 B1 B1</p>	<p>ANSWERED ON INSERT</p> <p>For correct initial temporary labels at F, G, I</p> <p>For correctly updating F and label at H</p> <p>For all temporary labels correct (including A) (allow extra 100 at C, 105 at D, 75 at H only)</p> <p>For order of becoming permanent correct</p> <p>For all permanent labels correct (A need not have a permanent label)</p> <p>For correct route (condone omission of J or B)</p> <p>For 125</p>
	<p>(ii)</p>	<p>Odd nodes: B C E J</p> <p>$BC = 60$ $BE = 35$ $BJ = 125$ $EJ = 90$ $CJ = 95$ $CE = 70$ 150 130 195</p> <p>Repeat BE and CJ (or BE, JI, IC)</p> <p>130 + 765 Shortest route: 895 metres</p>	<p>B1 M1 A1 M1 A1</p>	<p>For identifying or using B C E J or implied</p> <p>For any three of these weights correct, or implied or ft from their (i)</p> <p>For identifying the pairing BE, CJ to repeat or 130 (not ft)</p> <p>For 765 + their 130 (a valid pairs total)</p> <p>For 895 (cao)</p>
	<p>(iii)</p>	<p>Travelling salesperson problem</p>	<p>B1 M1 A1 B1</p>	<p>For graph structure correct</p> <p>For a reasonable attempt at arc weights (at least 9 correct, including the three given)</p> <p>For all arc weights correct</p> <p>For identifying TSP by name</p>

6	(i)	<table border="1"> <tr> <td></td> <td>1</td> <td>5</td> <td>2</td> <td>4</td> <td>3</td> <td>6</td> </tr> <tr> <td></td> <td>A</td> <td>B</td> <td>C</td> <td>D</td> <td>E</td> <td>F</td> </tr> <tr> <td>A</td> <td>-</td> <td>6</td> <td>3</td> <td>-</td> <td>-</td> <td>-</td> </tr> <tr> <td>B</td> <td>6</td> <td>-</td> <td>5</td> <td>6</td> <td>-</td> <td>14</td> </tr> <tr> <td>C</td> <td>3</td> <td>5</td> <td>-</td> <td>8</td> <td>4</td> <td>10</td> </tr> <tr> <td>D</td> <td>-</td> <td>6</td> <td>8</td> <td>-</td> <td>3</td> <td>8</td> </tr> <tr> <td>E</td> <td>-</td> <td>-</td> <td>4</td> <td>3</td> <td>-</td> <td>-</td> </tr> <tr> <td>F</td> <td>-</td> <td>14</td> <td>10</td> <td>8</td> <td>-</td> <td>-</td> </tr> </table>		1	5	2	4	3	6		A	B	C	D	E	F	A	-	6	3	-	-	-	B	6	-	5	6	-	14	C	3	5	-	8	4	10	D	-	6	8	-	3	8	E	-	-	4	3	-	-	F	-	14	10	8	-	-		ANSWERED ON INSERT
			1	5	2	4	3	6																																																				
			A	B	C	D	E	F																																																				
		A	-	6	3	-	-	-																																																				
		B	6	-	5	6	-	14																																																				
		C	3	5	-	8	4	10																																																				
		D	-	6	8	-	3	8																																																				
		E	-	-	4	3	-	-																																																				
		F	-	14	10	8	-	-																																																				
		Order: <i>A C E D B F</i>	M1	For choosing row C in column A																																																								
Minimum spanning tree:	M1 dep	For choosing more than one entry from column C																																																										
	A1	For correct entries chosen																																																										
Total weight: 23 miles	B1	For correct order, listed or marked on arrows or table, or arcs listed <i>AC CE ED CB DF</i>																																																										
	B1	For tree (correct or follow through from table, provided solution forms a spanning tree)																																																										
	B1	For 23 (or follow through from table or diagram, provided solution forms a spanning tree)																																																										
	6																																																											
(ii)	M1	MST for reduced network = 18																																																										
	M1	Two shortest arcs from B = 5 + 6 = 11																																																										
	A1	Lower bound = 29 miles																																																										
(iii)	M1	<i>F-D-E-C-A-B-F</i>																																																										
	A1	For correct tour																																																										
	M1	8 + 3 + 4 + 3 + 6 + 14 = 38 miles																																																										
	A1	For a substantially correct attempt at sum																																																										
	A1	For 38 (cao)																																																										

Mark Scheme 4737
June 2007

SOLUTIONS

4737




D2

June 2007

FINAL

1	(i)	house 1	house 2	house 3	house 4	B1	For copying the table, with row and column headings (accept consistent scalings)	
		A	500	400	700			600
		B	300	200	400			350
		C	500	300	750			680
	D	0	0	0	0	B1	2	For dummy row (Daniel) with all equal values
	(ii)	Reduce rows	100	0	300	200	M1	For a substantially correct attempt at reducing rows and columns
			100	0	200	150		
			200	0	450	380		
			0	0	0	0		
		Columns are already reduced						A1
Cross out using two lines						2		
		100		300	200	M1	For covering zeros using minimum number of lines, clearly seen or implied from augmenting	
		100		200	150			
		200		450	380			
Augment by 100						M1 dep	For a single augmentation by 100 (ft their matrix) (accept either way of augmenting by 100)	
		0	0	200	100			
		0	0	100	50	A1	ft For a correct augmented matrix (ft their matrix)	
		100	0	350	280			
		0	100	0	0	3		
(ii)	Cross out using three lines			200	100	M1	For covering zeros using minimum number of lines a second time, clearly seen or implied from augmenting	
				100	50			
				350	280			
	Augment by 50						M1 dep	For a single augmentation by 50 (ft their matrix) (accept either way of augmenting by 50)
			0	0	150	50		
			0	0	50	0	A1	ft For a correct augmented matrix (ft their matrix)
			100	0	300	230		
			50	150	0	0		
	Complete matching						B1	For a complete matching achieved, must follow from an attempt at reducing or augmenting a matrix, not just implied from a list of the matching
		0	0	150	50			
		0	0	50	0			
		100	0	300	230			
		50	150	0	0	4		
(iii)	Allclean	should clean	house 1				B1	For A = 1, B = 4, C = 2 (may also list D = 3) cao
	Brightenupp	should clean	house 4					
	Clean4U	should clean	house 2					
Cost = £1150						B1	2	For 1150 cao
						B1	13	

2	(i)	$4p - (1-p)$ $= 5p - 1$ $-2p + 5(1-p) = 5 - 7p$ $4(1-p) = 4 - 4p$	M1 A1	For $4p - 1(1-p)$ or equivalent, seen or implied For $5p - 1$ or $-1 + 5p$	cao
	(ii)		M1 A1 ft A1 ft A1 ft	For correct structure to graph with a horizontal axis that extends from 0 to 1, but not more than this, and with consistent scales. For line $E = 5p - 1$ plotted from (0,-1) to (1, 4) For line $E = 5 - 7p$ plotted from (0, 5) to (1,-2) For line $E = 4 - 4p$ plotted from (0, 4) to (1, 0)	cao cao cao
	(iii)	$p = 0.5$ $5(0.5) - 1$ $= 1.5$ points per game Bea may not play her best strategy	B1 M1 A1 B1	For this or ft their graph For substituting their p into any of their equations (must be seen, cannot be implied from value) For 1.5 For this or equivalent	cao
	(iv)	1.5 If Amy plays using her optimal strategy, Bea should never play strategy Z Assuming that Bea knows that Amy will make a random choice between P and Q so that each has probability 0.5, it does not matter how she chooses between strategies X and Y .	B1 ft M1 A1	Accept -1.5, ft from (iii) For identifying that she should not play Z For a full description of how she should play (If the candidate assumes that Bea does not know then Bea should play P with probability $\frac{7}{12}$ and Q with probability $\frac{5}{12}$).	3 15

<p>3 (i)</p>  <p>A dummy is needed after C because D follows both B and C. A dummy is needed after D because F and G both follow D.</p>	<p>M1 A1 B1 B1</p> <p style="text-align: right;">4</p>	<p>M1 A substantially correct network Condone arrows missing or wrong way round, no end and/or extra dummies Do NOT allow activity on node formulation</p> <p>A1 A correct network, with arrows on at least the dummy activities, with no extra dummies and a single end point.</p> <p>B1 A valid explanation</p> <p>B1 A valid explanation</p>
<p>(ii)</p>  <p>Minimum completion time = 14 days Critical activities are A, C, D, F</p>	<p>M1 A1 M1 A1 B1 B1</p> <p style="text-align: right;">6</p>	<p>M1 A substantially correct forward pass A1 Early event times correct (ft their network if possible)</p> <p>M1 A substantially correct backwards pass A1 Late event times correct (ft their network if possible)</p> <p>B1 For 14 cao</p> <p>B1 For these four activities and no others cao</p> <p>In both cases these need to be stated, not implied from the diagram</p>
<p>(iii)</p> 	<p>M1 M1 dep</p> <p style="text-align: right;">3</p>	<p>M1 For a reasonable attempt at using the number of workers for the different activities Scales and labels required and some days with 4 workers.</p> <p>M1 dep For a reasonable attempt with no overhanging blocks</p>
<p>(iv)</p>	<p>B1 B1</p> <p style="text-align: right;">2</p> <p style="text-align: right;">15</p>	<p>B1 Earliest finish for E > latest start for F</p> <p>B1 For delaying the start of F (by 1 day)</p>

4	(i)	stage	state	action	working	minimax		ANSWERED ON INSERT
		1	0	0	4	4		Values only credited when seen in table
			1	0	3	3		
			2	0	2	2		
		2	0	0	$\max(6,4) = 6$	3		
				1	$\max(2,3) = 3$			
				2	$\max(3,2) = 3$			
			1	0	$\max(2,4) = 4$	4	M1	For calculating the maxima as 4, 4, 5
				1	$\max(4,3) = 4$		A1 2	For calculating the minimax as 4
				2	$\max(5,2) = 5$			
		2	0	$\max(2,4) = 4$	3	B1	For completing 4, 3, 2 in the brackets	
			1	$\max(3,3) = 3$		M1	For calculating the maxima as 4, 3, 4 (method)	
			2	$\max(4,2) = 4$		A1 3	For calculating the minimax as 3 cao	
		3	0	0	$\max(5,3) = 5$	3	B1	For using their minimax values from stage 2
				1	$\max(5,4) = 5$		M1	For calculating the maxima for their values
				2	$\max(2,3) = 3$		A1 4	For calculating the maxima as 5, 5, 3 cao For calculating the minimax as 3 cao
	(ii)	3					M1	For the value from their tabulation
							A1	For 3 (irrespective of their tabulation) cao
							M1 dep	For reading route from their tabulation
							A1 4	For this route (irrespective of their tabulation) cao
	(iii)						B1	For the graph structure correct
							M1	For a substantially correct attempt at the weights (no more than two definite errors or omissions)
							A1	For weights unambiguously correct
							3 16	

5	(i)	$S - E - I - T$	B1 1	ANSWERED ON INSERT For this route (not in reverse) cao
	(ii)	6 litres per second From A to G	B1 2	For 6 For direction AG
	(iii)	$6 + 2 + 4 + 0 + 8$ = 20 litres per second	M1 M1 A1 3	For a substantially correct attempt with $DF = 0$ For dealing with $EI (= 8 \text{ or } = 2 + 6)$ For 20 cao Method marks may be implied from answer
	(iv)	eg flow 5 along $S - A - G - T$ and 2 along $S - C - F - H - G - T$	M1 A1 2	For describing a valid flow augmenting route For correctly flowing 7 from S to T
		Diagram correctly augmented	M1 M1 A1 3	For a reasonable attempt at augmenting a flow For correctly augmenting a flow For a correct augmentation by a total of 7
		Cut $\{S, A, B, C, D, E, F, G, H, I\}, \{T\}$	B1	For identifying cut or arcs GT and IT
		This cut has a value of 13 and the flow already found is $6 + 7 = 13$ litres per second. Or This is the maximum flow since the arcs GT and IT are both saturated, so no more can flow into T .	B1 2 13	For explaining how this shows that the flow is a maximum, but NOT just stating max flow = min cut

Advanced GCE Mathematics (3892 – 2, 7890 - 2)
June 2007 Assessment Series

Unit Threshold Marks

<i>Unit</i>		Maximum Mark	a	b	c	d	e	u
4721	Raw	72	60	52	44	36	29	0
	UMS	100	80	70	60	50	40	0
4722	Raw	72	56	48	40	33	26	0
	UMS	100	80	70	60	50	40	0
4723	Raw	72	57	50	43	36	29	0
	UMS	100	80	70	60	50	40	0
4724	Raw	72	61	54	47	40	33	0
	UMS	100	80	70	60	50	40	0
4725	Raw	72	54	46	39	32	25	0
	UMS	100	80	70	60	50	40	0
4726	Raw	72	60	53	46	39	33	0
	UMS	100	80	70	60	50	40	0
4727	Raw	72	57	50	43	36	29	0
	UMS	100	80	70	60	50	40	0
4728	Raw	72	57	49	42	35	28	0
	UMS	100	80	70	60	50	40	0
4729	Raw	72	59	51	44	37	30	0
	UMS	100	80	70	60	50	40	0
4730	Raw	72	62	54	46	38	31	0
	UMS	100	80	70	60	50	40	0
4731	Raw	72	51	43	36	29	22	0
	UMS	100	80	70	60	50	40	0
4732	Raw	72	55	48	42	36	30	0
	UMS	100	80	70	60	50	40	0
4733	Raw	72	56	48	41	34	27	0
	UMS	100	80	70	60	50	40	0

4734	Raw	72	56	49	42	36	30	0
	UMS	100	80	70	60	50	40	0
4735	Raw	72	60	51	43	35	27	0
	UMS	100	80	70	60	50	40	0
4736	Raw	72	62	55	48	42	36	0
	UMS	100	80	70	60	50	40	0
4737	Raw	72	61	53	46	39	32	0
	UMS	100	80	70	60	50	40	0

Specification Aggregation Results

Overall threshold marks in UMS (i.e. after conversion of raw marks to uniform marks)

	Maximum Mark	A	B	C	D	E	U
3890/3891/3892	300	240	210	180	150	120	0
7890/7891/7892	600	480	420	360	300	240	0

The cumulative percentage of candidates awarded each grade was as follows:

	A	B	C	D	E	U	Total Number of Candidates
3890	31.2	47.9	62.0	74.4	84.9	100	13873
3891	20.0	20.0	20.0	20.0	20.0	100	10
3892	58.5	75.6	87.9	94.7	97.5	100	1384
7890	45.3	66.9	82.2	92.4	97.7	100	9663
7891	0	0	0	100	100	100	1
7892	58.2	78.1	89.1	96.0	98.8	100	1487

For a description of how UMS marks are calculated see;
http://www.ocr.org.uk/exam_system/understand_ums.html

Statistics are correct at the time of publication

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

(General Qualifications)

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity



OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553