

**OXFORD CAMBRIDGE AND RSA EXAMINATIONS**  
**Advanced Subsidiary General Certificate of Education**  
**Advanced General Certificate of Education**

**MATHEMATICS**

**4723**

Core Mathematics 3

Wednesday **18 JANUARY 2006** Afternoon 1 hour 30 minutes

Additional materials:  
8 page answer booklet  
Graph paper  
List of Formulae (MF1)

**TIME** 1 hour 30 minutes

**INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- The total number of marks for this paper is 72.
- Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.
- **You are reminded of the need for clear presentation in your answers.**

---

**This question paper consists of 4 printed pages.**


1 Show that  $\int_2^8 \frac{3}{x} dx = \ln 64$ . [4]

2 Solve, for  $0^\circ < \theta < 360^\circ$ , the equation  $\sec^2 \theta = 4 \tan \theta - 2$ . [5]

3 (a) Differentiate  $x^2(x+1)^6$  with respect to  $x$ . [3]

(b) Find the gradient of the curve  $y = \frac{x^2 + 3}{x^2 - 3}$  at the point where  $x = 1$ . [3]

4




The function  $f$  is defined by  $f(x) = 2 - \sqrt{x}$  for  $x \geq 0$ . The graph of  $y = f(x)$  is shown above.

(i) State the range of  $f$ . [1]

(ii) Find the value of  $ff(4)$ . [2]

(iii) Given that the equation  $|f(x)| = k$  has two distinct roots, determine the possible values of the constant  $k$ . [2]

5



The diagram shows the curves  $y = (1 - 2x)^5$  and  $y = e^{2x-1} - 1$ . The curves meet at the point  $(\frac{1}{2}, 0)$ . Find the exact area of the region (shaded in the diagram) bounded by the  $y$ -axis and by part of each curve. [8]

6 (a)

$t$	0	10	20
$X$	275	440	

The quantity  $X$  is increasing exponentially with respect to time  $t$ . The table above shows values of  $X$  for different values of  $t$ . Find the value of  $X$  when  $t = 20$ . [3]


(b) The quantity  $Y$  is decreasing exponentially with respect to time  $t$  where

$$Y = 80e^{-0.02t}.$$

(i) Find the value of  $t$  for which  $Y = 20$ , giving your answer correct to 2 significant figures. [3]

(ii) Find by differentiation the rate at which  $Y$  is decreasing when  $t = 30$ , giving your answer correct to 2 significant figures. [3]

7



The diagram shows the curve with equation  $y = \cos^{-1} x$ .

(i) Sketch the curve with equation  $y = 3 \cos^{-1}(x - 1)$ , showing the coordinates of the points where the curve meets the axes. [3]

(ii) By drawing an appropriate straight line on your sketch in part (i), show that the equation  $3 \cos^{-1}(x - 1) = x$  has exactly one root. [1]


(iii) Show by calculation that the root of the equation  $3 \cos^{-1}(x - 1) = x$  lies between 1.8 and 1.9. [2]

(iv) The sequence defined by

$$x_1 = 2, \quad x_{n+1} = 1 + \cos\left(\frac{1}{3}x_n\right)$$

converges to a number  $\alpha$ . Find the value of  $\alpha$  correct to 2 decimal places and explain why  $\alpha$  is the root of the equation  $3 \cos^{-1}(x - 1) = x$ . [5]

[Questions 8 and 9 are printed overleaf.]



The diagram shows part of the curve  $y = \ln(5 - x^2)$  which meets the  $x$ -axis at the point  $P$  with coordinates  $(2, 0)$ . The tangent to the curve at  $P$  meets the  $y$ -axis at the point  $Q$ . The region  $A$  is bounded by the curve and the lines  $x = 0$  and  $y = 0$ . The region  $B$  is bounded by the curve and the lines  $PQ$  and  $x = 0$ .

(i) Find the equation of the tangent to the curve at  $P$ . [5]

(ii) Use Simpson's Rule with four strips to find an approximation to the area of the region  $A$ , giving your answer correct to 3 significant figures. [4]

(iii) Deduce an approximation to the area of the region  $B$ . [2]

9 (i) By first writing  $\sin 3\theta$  as  $\sin(2\theta + \theta)$ , show that

$$\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta. \quad [4]$$

(ii) Determine the greatest possible value of

$$9 \sin\left(\frac{10}{3}\alpha\right) - 12 \sin^3\left(\frac{10}{3}\alpha\right),$$

and find the smallest positive value of  $\alpha$  (in degrees) for which that greatest value occurs. [3]

(iii) Solve, for  $0^\circ < \beta < 90^\circ$ , the equation  $3 \sin 6\beta \operatorname{cosec} 2\beta = 4$ . [6]