

Mathematics

Advanced GCE

Unit 4723: Core Mathematics 3

Mark Scheme for June 2012

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2012

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

Annotations and abbreviations

Annotation in scoris	Meaning
✓ and *	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining
U1	Mark for correct units
G1	Mark for a correct feature on a graph
M1 dep* or dep*M	Method mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
A2	Accuracy mark awarded 2

Subject-specific Marking Instructions for GCE Mathematics Pure strand

- a. Annotations should be used whenever appropriate during your marking.

The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded.

- b. An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme assists in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must be marked on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the work must be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such methods must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, award marks to the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) contact your Team Leader.

- c. The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate to state an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of a mark is specified.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, eg wrong working following a correct answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply if a candidate passes through the correct answer as part of a wrong argument.

- d. When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme indicates otherwise; and similarly where there are several B marks allocated. (The notation 'dep **' is used to indicate that a part is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate is wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, if two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e. The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect work. Otherwise, A and B marks are given for correct work only — differences in notation are of course permitted. A (accuracy) mark is given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will be given for 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is in a different image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f. Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise. Candidates are expected to give numerical answers to an appropriate degree of accuracy, with 3 significant figures often being the expected standard. Variations in the degree of accuracy to which an answer is given (e.g. 2 or 4 significant figures where 3 is expected) should not be penalised, while answers which are grossly over- or under-specified should normally result in the loss of a mark. The mark scheme will specify any particular cases where the accuracy of the answer may be a marking issue should be detailed in the mark scheme rationale. In any doubt, contact your Team Leader.
- g. **Rules for replaced work**
 If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should mark the candidate requests.
 If there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the (complete) attempt and ignore the others.
 NB Follow these maths-specific instructions rather than those in the assessor handbook.

- h. For a *genuine* misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain the same according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is withheld. This may differ for some units. This is achieved by withholding one A mark in the question.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

Question	Answer	Marks	Guidance
1	Attempt process for finding critical values Obtain $\frac{4}{3}$ Obtain 6 Attempt process for inequality involving two critical values Obtain $x < \frac{4}{3}$, $x > 6$	M1 A1 A1 M1 A1 [5]	squaring both sides, 2 linear eqns, ineqs, ... sketch, table, ...; implied by plausible soln A0 for use of \leq and/or \geq
2	(i)	<u>EITHER</u> Attempt use of at least one logarithm property correctly applied to $\ln\left(\frac{ep^2}{q}\right)$ Obtain 261 legitimately with necessary detail seen <u>OR</u> Express $\frac{ep^2}{q}$ in form e^n Obtain e^{261} and hence 261	M1 not including $\ln e = 1$; such as... = $\ln ep^2 - \ln q$ for example A2 AG; award A1 if nothing wrong but not quite enough detail or if there is one slip on way to 261 [3] M1 with correct treatment of powers A2 AG; award A1 if nothing wrong but not quite enough detail to be fully convincing
2	(ii)	Introduce logarithms and bring power down Obtain $n \ln 5 > 580$ State single integer 361	M1 relating $n \ln 5$ to a constant; if using base 5 or base 10, no powers must remain on right-hand side A1 or equiv (such as $n > 580 \log_5 e$ or $n \log 5 > 580 \log e$); allow eqn at this stage A1 not $n > 360$ nor $n \geq 361$ [3]

Question		Answer	Marks	Guidance
3	(i)	Use $\sec \theta = \frac{1}{\cos \theta}$ Attempt to express in terms of $\tan \theta$ only Obtain $\tan^2 \theta = 36$ and hence $\tan \theta = 6$	B1 M1 A1 [3]	AG; necessary detail needed (but no need to justify exclusion of $\tan \theta = -6$)
3	(ii) (a)	Substitute 6 in attempt at formula Obtain $\frac{5}{7}$	M1 A1 [2]	of form $\frac{\tan \theta \pm \tan 45^\circ}{1 \mp \tan \theta \tan 45^\circ}$ with different signs in numerator and denominator or exact equiv
3	(ii) (b)	Substitute 6 in attempt at formula Obtain $-\frac{12}{35}$	M1 A1 [2]	of form $\frac{\tan \theta + \tan \theta}{1 \pm \tan \theta \tan \theta}$ or exact equiv; allow $\frac{12}{-35}$
4	(a)	Obtain integral of form $k(6x+1)^{\frac{1}{2}}$ Obtain $6(6x+1)^{\frac{1}{2}}$ Substitute both limits and subtract Obtain $30 - 6$ and hence 24	*M1 A1 M1 A1 [4]	any constant k or (unsimplified) equiv dep *M AG; necessary detail needed
4	(b)	Attempt expansion of integrand Integrate e^{kx} to obtain $\frac{1}{k}e^{kx}$ Obtain $\frac{1}{2}e^{2x} + 4e^x + 4x$ Obtain $\frac{1}{2}e^2 + 4e - \frac{1}{2}$	M1 M1 A1 A1 [4]	to obtain (at least) 3 terms for any constant k other than 1 allow $+c$ at this stage or equiv in terms of e simplified to three terms; no $+c$ now

Question		Answer	Marks	Guidance	
5	(i)	Sketch (more or less) correct $y = 14 - x^2$	B1	assessed separately from other graph; must exist in all four quadrants; ignore any intercepts given	
		Sketch (more or less) correct $y = k \ln x$	B1	assessed separately from other graph; must exist in first and fourth quadrants; if clearly meets y-axis award B0; if clear maximum point in first quadrant award B0	
		Indicate one root ('blob' on sketch or written reference to one intersection or ...)	B1	dependent on both curves being correct in first quadrant and there being no possibility, from their graphs, of further points of intersection elsewhere	
			[3]		
5	(ii)	(a)	Calculate values for at least 2 integers	M1	
			Obtain correct values for $x = 3$ and $x = 4$	A1	$14 - x^2 - 3 \ln x : 1.7 \quad -6.2$
			State 3 and 4	A1	$14 - x^2, 3 \ln x: 5, 3.3 \quad -2, 4.2$
			[3]	following correct calculations	
5	(ii)	(b)	Obtain correct first iterate	B1	having started with any positive value; B1 available if 'iteration' never goes beyond a first iterate;
			Attempt iteration process	M1	implied by plausible sequence of values
			Obtain at least 3 correct iterates in all	A1	showing at least 2 d.p.
			A1	answer required to exactly 2 d.p; not given for 3.24 as the final iterate in a sequence, i.e. needs an indication (perhaps just underlining) that value of α found	
			[4]	$[3 \rightarrow 3.27172 \rightarrow 3.23173 \rightarrow 3.23743 \rightarrow 3.23661$ $3.5 \rightarrow 3.20027 \rightarrow 3.24196 \rightarrow 3.23596 \rightarrow 3.23682$ $4 \rightarrow 3.13706 \rightarrow 3.25118 \rightarrow 3.23465 \rightarrow 3.23701]$	

Question	Answer	Marks	Guidance
6 (i)	Attempt use of chain rule Obtain $9h(3h^2 + 4)^{\frac{1}{2}}$ Substitute 0.6 in attempt at first derivative Obtain 12.17	*M1 A1 M1 A1 [4]	to obtain derivative of form $kh(3h^2 + 4)^n$, any non-zero constants k and n condone retention of -8 or (unsimplified) equiv; no -8 here dep *M; condone retention of -8 here; implied by their value following wrong derivative if no working seen or greater accuracy
6 (ii)	State or imply that $\frac{dh}{dt} = -0.015$ or 0.015 Carry out multiplication of $(\pm)0.015$ and answer from part (i) Obtain 0.18 or -0.18 (whatever this value is claimed to be)	B1 M1 A1 [3]	implied by use in calculation with part (i) answer or greater accuracy; condone absence or misuse of negative signs throughout; ignore units; allow for answer rounding to 0.18 following slight inaccuracy due to use of 12.18 or 12.2 or ...
7	Show composition of functions Obtain $2\sqrt[3]{12 - a} + 5 = 9$ Obtain $a = 4$ <u>EITHER</u> Attempt to find $g(x)$ Obtain $(2x + 5)^3 + 4 = 68$ Attempt solution of equation Obtain $-\frac{1}{2}$ <u>OR</u> State or imply $f(x) = g^{-1}(68)$ Attempt solution of equation of form $2x + 5 = \sqrt[3]{68 - 4}$ Obtain $-\frac{1}{2}$	M1 A1 A1 *M1 A1ft M1 A1 [7] B2 M1 A1	the right way round; or equiv or equiv obtaining $px^3 + q$ or $py^3 + q$ form following their value of a dep *M; earned at stage $2x + 5 = \dots$; if expanding to produce cub attempt at linear and quadratic factors and no others; dependent on correct work throughout

Question		Answer	Marks	Guidance
8	(i)	State $R = 5$ Attempt to find value of α Obtain 53.1	B1 M1 A1 [3]	implied by correct value or its complement allow $\tan^{-1}\frac{4}{3}$
8	(ii) (a)	Attempt to find at least one value of $\theta + \alpha$ Obtain 1 correct value of θ (-64.7 or 138) Attempt correct process to find the second value Obtain second value of θ (138 or -64.7)	M1 A1 M1 A1 [4]	(should be -168.5 or -11.5 or 191.5 or ...) allow ± 0.1 in answer and greater accuracy involving a positive value of $\sin^{-1}(-\frac{1}{3})$ and subtraction of their α allow ± 0.1 in answer and greater accuracy; and no others between -180 and 180
8	(ii) (b)	Use -1 as minimum or 1 as maximum value of $\sin(\theta + \alpha)$ Relate $-5k + c$ to -37 and $5k + c$ to 43 Attempt solution of pair of linear eqns Obtain $k = 8$ and $c = 3$	*M1 A1 M1 A1 [4]	as equations or inequalities dep *M; must be equations now SC: both $k = 8$ and $c = 3$ obtained with no working or from unconvincing working, award B2 (i.e. max 2/4)

Question	Answer	Marks	Guidance
9 (i)	Attempt use of product rule to produce the form $\ln 2y + y \times \frac{a}{by}$ Obtain correct $\ln 2y + y \times \frac{2}{2y} \dots$ Obtain complete $\ln 2y + 1 - 1$ and confirm	M1 A1 A1 [3]	or equiv AG; necessary detail needed
9 (ii)	Attempt to rearrange eqn to $x = \dots$ or $x^2 = \dots$ Obtain $x = \sqrt{\ln 2y}$ or $x^2 = \ln 2y$ State or imply volume is $\int \pi \ln 2y \, dy$ Integrate using result of part (i) Attempt to use limits $\frac{1}{2}$ and $\frac{1}{2}e^4$ correctly with expression involving y Obtain $\frac{1}{2}\pi(3e^4 + 1)$	M1 A1 A1ft M1 M1 A1 [6]	obtaining form $p \ln qy$ following their $x = \dots$ or $x^2 = \dots$; condone absence of dy ; condone presence of dx ; no need for limits here; π may be implied by its first appearance later in solution or equiv involving two terms; dependent on correct work throughout part (ii)
9 (iii)	Subtract answer to part (ii) from $2\pi e^4 \dots$ Obtain $\frac{1}{2}\pi(e^4 - 1)$	M1 A1 [2]	... or its decimal equivalent or exact equiv involving two terms

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2012

