

June 2007.

Q1e4.

1) $f(x) = \frac{3x+1}{(x+2)(x-3)} = \frac{A}{x+2} + \frac{B}{x-3}$ $A(x-3) + B(x+2) \equiv 3x+1$ $5B = 10$ $B = 2$
 $-5A = -5$ $A = 1$

(i) $f(x) = \frac{1}{x+2} + \frac{2}{x-3}$ $f'(x) = -\frac{1}{(x+2)^2} - \frac{2}{(x-3)^2} = -1 \left(\frac{1}{(x+2)^2} + \frac{2}{(x-3)^2} \right)$ hence $f'(x)$ is always negative for all x .



2) $\int_0^1 x^2 e^x dx$ $u = x^2$ $\frac{du}{dx} = 2x$ $\frac{dv}{dx} = e^x$ $v = e^x$ $\int_0^1 x^2 e^x dx = x^2 e^x - \int e^x 2x dx$

$\int_0^1 e^x 2x dx$ $u = 2x$ $\frac{du}{dx} = 2$ $\frac{dv}{dx} = e^x$ $v = e^x$ $\int_0^1 e^x 2x dx = 2x e^x - \int e^x 2 dx$

$\int_0^1 x^2 e^x dx = [x^2 e^x - (2x e^x - 2e^x)]_0^1 = [x^2 e^x - 2x e^x + 2e^x]_0^1 = (e^1 - 2e^1 + 2e^1) - (0 - 0 + 2)$

$\int_0^1 x^2 e^x dx = e^1 - 2$.

3) $y = \sin x$

 $V = \pi \int y^2 dx = \pi \int \sin^2 x dx$

$\cos 2A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$

$\sin^2 A = \frac{1}{2}(1 - \cos 2A)$


$V = \pi \int \frac{1}{2} (1 - \cos 2x) dx = \frac{\pi}{2} \left[x - \frac{\sin 2x}{2} \right]_0^\pi = \frac{\pi}{2} \left[\left(\pi - \frac{1}{2} \sin 2\pi \right) - \left(0 - \frac{1}{2} \sin 0 \right) \right] = \frac{\pi^2}{2}$

4) $(2+x)^{-2} = \left(2 \left(1 + \frac{x}{2} \right) \right)^{-2} = \frac{1}{4} \left(1 + \frac{x}{2} \right)^{-2} = \frac{1}{4} \left\{ 1 + (-2) \left(\frac{x}{2} \right) + \frac{(-2)(-3)}{2!} \left(\frac{x}{2} \right)^2 + \frac{(-2)(-3)(-4)}{3!} \left(\frac{x}{2} \right)^3 + \dots \right\}$
 $= \frac{1}{4} \left\{ 1 - x + \frac{3x^2}{4} - \frac{1}{2} \frac{x^3}{2} + \dots \right\} = \frac{1}{4} - \frac{x}{4} + \frac{3x^2}{16} - \frac{x^3}{8} + \dots$
 $\left| \frac{x}{2} \right| < 1 \quad -2 < x < 2$

(ii) $(1+x^2) \left\{ \frac{1}{4} - \frac{x}{4} + \frac{3x^2}{16} - \frac{x^3}{8} + \dots \right\}$ coefficient $x^3 \left(-\frac{1}{4} - \frac{1}{8} \right) = -\frac{3}{8}$

5. $x = \cos t$ $y = 3 + 2 \cos 2t$ $\frac{dy}{dx} = \frac{4 \sin 2t}{-2 \sin t} = \frac{4 \times 2 \sin t \cos t}{-2 \sin t} = -4 \cos t$
 $\frac{dx}{dt} = -\sin t$ $\frac{dy}{dt} = -2 \sin 2t$ max value of $-4 \cos t = 4$ when $\cos t = -1$ $t = \pi$

$3 + 2 \cos 2t = 3 + 2(2 \cos^2 t - 1) = 3 + 4 \cos^2 t - 2 = 1 + 4 \cos^2 t$ $y = 1 + 4x^2$



6. $x^2 + 3xy + 4y^2 = 58$ $2x + 3x \frac{dy}{dx} + 3y + 4 \times 2y \frac{dy}{dx} = 0$
 $(3x + 8y) \frac{dy}{dx} + 2x + 3y = 0$ $\frac{dy}{dx} = \frac{-(2x + 3y)}{3x + 8y}$ grad $\log t(2, 3) = -\frac{13}{30}$

grad of normal $+\frac{30}{13}$ at $(2,3)$

$$(y-3) = +\frac{30}{13}(x-2)$$

$$13y - 39 = +30x - 60$$

$$-30x - 13y = 21 = 0$$

7) $x^2+4 \overline{) 2x^3 + 3x^2 + 9x + 12}$

quotient $2x+3$ remainder $\frac{x}{x^2+4}$

$$\begin{array}{r} 2x^3 + 0 + 8x \\ \underline{3x^2 + 0 + 12} \\ 3x^2 + 0 + 12 \\ \underline{0 + x + 0} \end{array}$$

$$\frac{2x^3+3x^2+9x+12}{x^2+4} = 2x+3 + \frac{x}{x^2+4} \quad A=2 \quad B=3 \quad C=1 \quad D=0$$


$$\int_1^3 (2x+3 + \frac{x}{x^2+4}) dx = \frac{2x^2}{2} + 3x + \frac{1}{2} \int_1^3 \frac{2x}{x^2+4} dx = \left[x^2 + 3x + \frac{1}{2} \ln|x^2+4| \right]_1^3$$

$$= (9 + 9 + \frac{1}{2} \ln 13) - (1 + 3 + \frac{1}{2} \ln 5) = 14 + \frac{1}{2} (\ln 13 - \ln 5) = 14 + \frac{1}{2} \ln \frac{13}{5}$$

8) $\frac{dh}{dt} = \frac{6-h}{20} \quad \int \frac{1}{6-h} dh = \int \frac{1}{20} dt \quad \ln|6-h| = \frac{t}{20} + k$


$\ln|6-h| = -\frac{t}{20} - k \quad t=0 \quad h=1 \quad -\ln 5 = -k \quad -\ln|6-h| + \ln 5 = +\frac{t}{20}$

$20 \ln \left| \frac{5}{6-h} \right| = t \quad \text{height at planting is 1m} \quad h=2m \quad t = 20 \ln \frac{5}{4} \quad t = 4.46 \text{ years (3sf)} \quad 4.4629$



$$e^{\frac{t}{20}} = \frac{5}{6-h} \quad 6-h = \frac{5}{e^{\frac{t}{20}}} = 3.03265 \quad h = 2.9673$$

$$h = 2.97 \text{ (3sf)}$$



9(i) $l_1 \cdot l_2 = \begin{pmatrix} -6 \\ 8 \\ -2 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix} = -6 + 24 - 4 = 14 \quad |l_1| |l_2| \cos \theta = 14$

$$\cos \theta = \frac{14}{\sqrt{36+64+4} \times \sqrt{1+9+4}} = \frac{14}{\sqrt{104} \sqrt{14}} = \frac{14}{\sqrt{16 \times 91}} = \frac{14}{4\sqrt{91}}$$

(ii) $\begin{pmatrix} -6 \\ 8 \\ -2 \end{pmatrix} = -2 \begin{pmatrix} 3 \\ c \\ 1 \end{pmatrix} \quad \therefore c = -4$
parallel.

(iii) $L_2 \quad L_3$ intersect $\begin{pmatrix} 3 \\ -8 \\ 0 \end{pmatrix} + t \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} + u \begin{pmatrix} 3 \\ c \\ 1 \end{pmatrix}$

(a) $3+t = 2+3u \quad t = 3u-1 \quad 2t = 6u-2 = 3+u \quad 5u = 5 \quad u=1 \quad t=2$

(c) $2t = 3+u$

(y) $-8+6 = 1+c \quad c = -3$