

Mathematics

Advanced GCE

Unit **4734**: Probability and Statistics 3

Mark Scheme for June 2012

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2012

Annotations

Annotation	Meaning
✓ and ✗	
BOD	Benefit of doubt
FT or √	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining
U1	Mark for correct units
G1	Mark for a correct feature on a graph
M1 dep*	Method mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working

Subject-specific Marking Instructions

- a Annotations should be used whenever appropriate during your marking.

The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded.

- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme assist in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must be the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the work must be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such methods must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, but is to the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) contact your Team Leader.

- c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate to show the intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of a mark is specified.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a candidate who passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme indicates otherwise; and similarly where there are several B marks allocated. (The notation 'dep *' is used to indicate that a part is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate is wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, if two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously indicated work. Otherwise A and B marks are given for correct work only — differences in notation are of course permitted. A (accuracy) mark is given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will be given for 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is in the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.

Candidates are expected to give numerical answers to an appropriate degree of accuracy. 3 significant figures may be expected, but this always needs to be considered in the context of the problem in hand. For example, in quoting probabilities from tables, we generally expect *some* evidence of interpolation and so quotation to 4 decimal places will often be appropriate. However, this does not always apply – quotations of the standard critical points for significance tests such as 1.96, 1.645, 2.576 (not 2.57) will commonly suffice, especially if the calculated value of a test statistic is nowhere near any of these values. Discretion *must* be exercised in such cases.

Discretion must also be exercised in the case of small variations in the degree of accuracy to which an answer is given. If 2 significant figures are expected (either because of an explicit instruction or because the general context of a problem requires it), loss of an accuracy ("A") mark is likely to be appropriate; but if 4 significant figures are given, this should not be penalised.

Likewise, answers which are slightly deviant from what is expected in a very minor manner (for example, an attempt at interpolation, as 0.6418 whereas 0.6417 was expected) should not be penalised. However, over- or under-specified should normally result in the loss of a mark. This includes cases such as, for example, if the test statistic is (say) 2.128888446667 merely because that is the value that happened to come off the candidate's calculator. This applies to answers that are given as final stages of calculations; intermediate working should usually be carried out to a greater degree of accuracy to avoid the danger of premature approximation.

The situation regarding any particular cases where the accuracy of the answer may be a marking issue should be dealt with in the scheme rationale. If in doubt, contact your Team Leader.

g Rules for replaced work

If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examine the candidate requests.

If there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the (complete) attempt and ignore the others.

NB Follow these maths-specific instructions rather than those in the assessor handbook.

h Genuine misreading (of numbers or symbols, occasionally even of text) occurs. If this results in the object and/or difficulty being considerably changed, it is likely that all the marks for that question, or section of the question, will be lost. However, often such that the object and/or difficulty remain substantially unaltered; these cases are considered below.

The simple rule is that *all* method ("M") marks [and of course all independent ("B") marks] remain accessible but at least some ("A") marks do not. It is difficult to legislate in an overall sense beyond this global statement because misreads, even if the object and/or difficulty remains unchanged, can vary greatly in their effects. For example, a misread of 1.02 as 10.2 (perhaps of a sample mean) may well be catastrophic; whereas a misread of 1.6748 as 1.6746 may have so slight an effect as to be unnoticeable in the candidate's work.

A misread should normally attract *some* penalty, though this would often be only 1 mark and should rarely if ever be more. Commonly in sections of questions where there is a numerical answer either at the end of the section or to be obtained from (eg the value of a test statistic), this answer will have an "A" mark that may actually be designated as "cao" [correct answer only] – should be interpreted *strictly* – if the misread has led to failure to obtain this value, then this "A" mark must be withheld. If other marks have been earned. It will also often be the case that such a mark is implicitly "cao" even if not explicitly designated as such.

On the other hand, we commonly allow "fresh starts" within a question or part of question. For example, if a candidate's value of a test statistic is generally allowed (and often explicitly stated as such within the mark scheme) a candidate may exhibit knowledge of how to compare it with a critical value and draw conclusions. Such "fresh starts" are allowed for any earlier misreads.

A misread may be of a symbol rather than a number – for example, an algebraic symbol in a mathematical expression. Such misreads are more likely to bring about a considerable change in the object and/or difficulty of the question; but, if they do not, they are treated as far as possible in the same way as numerical misreads, *mutatis mutandis*. This also applied to misreads of units, which are rare but can cause major problems in fair marking.

The situation regarding any particular cases that arise while you are marking for which you feel you need detailed guidance should be discussed with your Team Leader.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

Question	Answer	Marks	
1	(i) $0.14 \pm zs$ $z = 1.96$ $s^2 = 0.14 \times 0.86/100$ $(0.072(0), 0.2080)$	M1 B1 A1 A1 [4]	Or /99 $(0.0716, 0.208(4))$ from /99 A0
	(ii) Equate $\phi((500 - \mu)/\sqrt{(50)})$ to 0.14 $\Rightarrow (500 - \mu)/\sqrt{(50)} = -1.0803$ $\Rightarrow \mu = 508$ (3SF)	M1 A1 A1 [3]	
2	Assumes population of time differences is normal $H_0: \mu_F = \mu_S, H_1: \mu_F \neq \mu_S$ Differences 6 -3 7 3 Sample mean = 3.25 Sample variance = 4.5^2 = 20.25 TS = $3.25/(\sqrt{4.5}/2)$ = 1.44(4) Not ≥ 2.353 do not reject H_0 There is insufficient evidence of a difference in the mean times of the two populations	B1 B1 M1 A1 A1 M1 A1 M1 A1 ft [9]	Need population. Allow $\mu_d=0$ etc. One tail B1B0M1A1A1M1A1M1(CV=1.638)A0 Comparison with correct t and correct first conclusion ft. Ft TS Not over-assertive.

Question	Answer	Marks	
3	$TS = (39/60 - 12/40 - 0.15)/s$ $s^2 = 0.65 \times 0.35/60 + 0.3 \times 0.7/40 (=0.0090417 \text{ (5sf)})$ $TS = 2.10(3)$ Compare TS with 1.96 and reject H_0 Reports that there is evidence (at 5% SL) that. popn resp to coloured more than 15% greater than popn resp to plain	M1 M1A1 A1 M1 A1ft [6]	No marks for hyps in this Q. M1 for “variance” involving 40 and 60 Or “1.78%” < 2.5% Ft TS Not over-assertive
4	(i) $\bar{t} = 348/24 (=14.5)$ $s^2 = (5195.5 - 348^2/24)/23(=6.5)$ $348/24 \pm t.s / \sqrt{(24)}$ $t = 2.069$ $(13.423, 15.557)$	B1 B1 M1 B1 A1 [5]	Allow $z=1.96$ Rounding to (13.4, 15.6)
	(ii) $H_0: \mu = 15 \text{ (Or } \geq), H_1: \mu < 15$ $\alpha (14.5 - 15) / \sqrt{(6.5/24)}$ $= -0.961$ $> -1.319, \text{ do not reject } H_0$ $\beta \bar{x} \leq 15 - 1.139 \sqrt{6.5/24}$ $= 14.3(1)$ < 14.5 Insufficient evidence at the 5% SL that the machine is not working acceptably	B1 B1 M1 A1 M1 M1 A1 M1 A1ft [6]	Must have /24 t -value t -value t -value ft-TS, not over-assertive

Question	Answer	Marks		
5	(i) $(E(X)=16, E(Y)=28) E(Y-X)=12$ $(\text{Var}(X)=8, \text{Var}(Y)=28) 28+32 \times \frac{1}{2} \times \frac{1}{2}$ $\text{Var}(Y-X) = 36$	B1 M1 A1 [3]		
	(ii) $(X \approx N(16,8))$ since $32 \times \frac{1}{2} > 5$ and $32 \times \frac{1}{2} \times \frac{1}{2} > 5$ $(Y \approx N(28,28))$ since $E(Y) = 28$ is large oe (eg>15) $Y-X \approx N(12, 36)$	M1 M1 B1ft [3]	from $np > 5$ and $nq > 5$ or $npq > 5$ N(parameters from (i))	
	(iii) $P(Y-X \leq -2.5) + P(Y-X \geq 2.5)$ $= \Phi(-2.417) + 1 - \Phi(-1.583)$ $0.0079 + 0.9433 = 0.951(2)$	M1 A1 M1 A1 [4]	(-2.417, -1.583) Correct use of z and Φ for their value(s) Allow M1A0 if only 1 interval, except M1A1 if 0.0079 or 0.9433 seen (SC)	No M ± 2 or A1

Question	Answer	Marks	
6	(i) Total area from $t = 0$ to 4 is 1 and $\frac{1}{2}a + 3a = 1$ Solve to give $a = 2/7$	M1 A1 [2]	Any method
6	(ii) $0 < t < 1$ $t^2/7$ $0 \leq t \leq 1$ $\int_0^1 2t/7 dt = 2t^2/14 - 1/7$ $1 < t < 4$	B1 M1 A1 B1 [4]	Ft a for B1 and A1 For $t < 0$ and $t > 4$
	(iii) $G(Y) = P(Y \leq y) = P(T \leq y^3)$ $= F(y^2)$ $= \begin{cases} 0 & y < 0, \\ y^4/7 & 0 \leq y \leq 1, \\ (2y^2 - 1)/7 & 1 < y \leq 2, \\ 1 & y \geq 2 \end{cases}$ $g(y) = G'(y) = \begin{cases} 4y^3/7 & 0 \leq y \leq 1 \\ 4y/7 & 1 < y \leq 2 \\ 0 & \text{otherwise} \end{cases}$	M1 A1 M1 A1 ft M1 A1 B1 [7]	Allow < Seen or implied Possible to score M0A0, then M1A1M1A1B1 , For using F correctly For correct expressions ft a and F(t) For differentiating Correctly, allow from eg $2y^2/7$ Correct ranges for y seen

Question	Answer	Marks																					
7	(i) $H_0: p_1 = p_2 = p_3 (=1/3)$, H_1 : Not all equal oe eg in words E-values all 30 $\chi^2 = (1^2 + 5^2 + 4^2)/30$ $= 1.4$ (Critical value = 5.991) Compare χ^2 with CV and do not reject H_0 . Insufficient evidence that groups not randomly chosen.oe	B1 B1 M1 A1 M1 A1ft [6]	$p = \frac{1}{3}$ only insufficient. Accept 1.39, 1.399 etc With valid comparison or Accept that groups randomly chosen.																				
	(ii) (H_0 : Outcome indep of treatment) <table border="1" style="margin-left: 20px;"> <tr> <td>Treatment</td> <td>A</td> <td>B</td> <td>C</td> </tr> <tr> <td>Improved</td> <td>14</td> <td>18</td> <td>25</td> </tr> <tr> <td>Not Improved</td> <td>17</td> <td>7</td> <td>9</td> </tr> </table> <table border="1" style="margin-left: 20px;"> <tr> <td>E-values</td> <td>19.6</td> <td>15.8</td> <td>21.5</td> </tr> <tr> <td></td> <td>11.4</td> <td>9.2</td> <td>12.5</td> </tr> </table>	Treatment	A	B	C	Improved	14	18	25	Not Improved	17	7	9	E-values	19.6	15.8	21.5		11.4	9.2	12.5	B1 M1 A1	For correct contingency table At least 2 correct FT table All correct CAO
Treatment	A	B	C																				
Improved	14	18	25																				
Not Improved	17	7	9																				
E-values	19.6	15.8	21.5																				
	11.4	9.2	12.5																				
	(iii) $\chi^2 = 5.6^2(19.6^{-1} + 11.4^{-1}) + 2.2^2(15.8^{-1} + 9.2^{-1}) + 3.5^2(21.5^{-1} + 12.5^{-1})$ $= 6.7$ > 5.991 and reject H_0 Sufficient evidence at the 5% SL that the outcome depends on treatment Treatment A shows fewer improved than expected, Treatments B and C show more So abandon Treatment A	M1 A1 A1 M1 A1ft [8] M1 A1 [2]	At least 2 correct ft All correct Ft TS if attempt at correct table made. Or consider proportion(s) improved, 0.45, 0.72, 0.74 (M1)																				

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2012

