



GCE AS MARKING SCHEME

SUMMER 2018

**AS (NEW)
MATHEMATICS – UNIT 2 APPLIED MATHEMATICS A
2300U20-1**

INTRODUCTION

This marking scheme was used by WJEC for the 2018 examination. It was finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conference was held shortly after the paper was taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conference was to ensure that the marking scheme was interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conference, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about this marking scheme.

AS Mathematics Unit 2: Applied Mathematics A

Solutions and Mark Scheme Summer 2018

SECTION A – Statistics

Qu. No.	Solution	Mark	Notes
1.	$P(X = 7) = \binom{16}{7} \times 0.3^7 \times 0.7^9$ $P(X = 7) = 0.1009(6 \dots)$	M1 A1 [2]	M0 if no calculation shown. Accept 0.101.
2(a)	(The set of) students who study Mathematics and not Drama.	E1	Do not accept reference to 'number of students' or 'probability'
(b) (i)	$\frac{6}{40}$ OR $\frac{3}{20}$ OR 0.15	B1	
(ii)	$P(M \cup F) = \frac{13 + 2 + 10 + 4}{40} \text{ oe}$ $= \frac{29}{40} \text{ OR } 0.725$	M1 A1	
(c)	$P(M) = \frac{16}{40} \quad P(D) = \frac{10}{40} \quad P(M \cap D) = \frac{4}{40}$ $P(M) \times P(D) = \frac{16}{40} \times \frac{10}{40} = \frac{1}{10} (= P(M \cap D))$ <p>Since $P(M) \times P(D) = P(M \cap D)$ they are statistically independent.</p>	B1 B1 E1 [7]	All 3 correct (0.4, 0.25, 0.1) Correctly evaluating 'their P(M)' x 'their P(D)' provided at least one correct. Accept alternative method FT candidate's probabilities provided B1 awarded. Convincing.


Qu. No.	Solution	Mark	Notes
5(a)	Strong linear relationship	E1	
	The higher the hydration the lower the pH.	E1	
(b)(i)	Each additional ml of water per 100g of flour decreases the pH by 0.02 <u>on average</u> .	E1	Or equivalent
	The intercept would imply that at zero hydration the pH would be 5.4.	E1	
(ii)	$y = 5.4 - 0.02 \times 20$	B1	From use of regression line.
	$y = 5$	E1	
	Any correct comment. e.g. Outside the data set Extrapolation, etc.		
		[6]	
6(a)(i)	$1.5(21 - 16)$ $= 7.5$	B1	B1: 7.5 May be implied by sight of 8.5 or 28.5. M1: Correct method for either using "their 7.5" A1: 8.5 and 28.5 both correct.
	$16 - 7.5 = 8.5$ (therefore no outliers below 8.5)	M1	
	$21 + 7.5 = 28.5$ (therefore outliers above 28.5)	A1	
	40 is an outlier but there may be others.	E1	E1: stating that 40 is an outlier but must also state or imply that the summary stats don't show if there are any others.
(b)	Positive skew.	B1	Accept skewed to the right.
	Appropriate comment. e.g. A few tutors are very expensive. The bulk of the tutors are relatively cheap.	E1	
(c) (i)	Mean will decrease.	E1	
(ii)	Median may stay the same or it may decrease.	E1	Must imply that we don't know unless we know the individual values.
(d)	Two appropriate comments. e.g. Dafydd's lessons are cheaper on average than Basel's. Dafydd's lessons are more variable in cost than Basel's Both are positively skewed.	E2	E1 for one appropriate comment. E0 if omission of "on average" ISW unless contradicts a previous correct statement. Do not allow "data is more variable" without reference to cost somewhere in the answer.
		[10]	

Section B – Mechanics

Q	Solution	Mark	Notes
7	$x = \int 6t^2 - 8t - 5 \, dt$ $x = \frac{6}{3}t^3 - \frac{8}{2}t^2 - 5t + (C)$ $x = 2t^3 - 4t^2 - 5t + (C)$ <p>when $t = 1, x = -4$</p> $C = -4 - 2 + 4 + 5 = 3$ $x = 2t^3 - 4t^2 - 5t + 3$	M1 A1 A1	at least 1 term with increased power cao

Q	Solution	Mark	Notes
8(a)	Apply N2L to both particles.	M1	dim correct for at least 1
			Allow $T \pm 3g = 3a$, T and $5g$ opposing.
	$T = 3a$	B1	first correct equation
	$5g - T = 5a$	A1	second correct equation
	$5g = 8a$	m1	
	$a = 6.125 \text{ (ms}^{-2}\text{)}$	A1	cao
	$T = 18.375 \text{ (N)}$	A1	cao
8(b)	If the pulley is rough, the tension in the string on either side of the pulley would not be the same.	E1	

Q	Solution	Mark	Notes
9.	$R = (2 + 3 + 4)\mathbf{i} + (5 - 22 - 23)\mathbf{j}$	M1	
	$R = 9\mathbf{i} - 40\mathbf{j}$	A1	cao si
	$ R = \sqrt{9^2 + 40^2}$	M1	ft R
	$ R = 41 \text{ (N)}$	A1	ft R only if 2 non-zero components
	$\theta = \tan^{-1}\left(-\frac{40}{9}\right)$	M1	
	$\theta = -77.32^\circ \text{ or } 282.68^\circ$	A1	cao direction clearly indicated eg angle in fourth quadrant, diagram with resultant marked.

Q	Solution	Mark	Notes
10(a)	Apply N2L to lift and man $8000 - (770 + 68)g = (770 + 68)a$ $a = -0.25 \text{ (ms}^{-2}\text{) (correct to 2 d.p.)}$ SC	M1 A1 A1	Dim correct equation. Tension and wt opposing. correct equation. cao
	(a) Apply N2L to lift only $8000 - 770g = 770a$ $a = 0.59 \text{ (ms}^{-2}\text{) (correct to 2 d.p.)}$	M1 A1	Dim correct equation. Tension and wt opposing. cao
10(b)	As the acceleration is negative, the lift is slowing down. B0 if SC in (a)	B1	depends on M1 in (a)
10(c)	 Apply N2L to man $R - 68g = 68a$ $R - 68g = 68 \times (-0.25)$ $R = 649.16 \text{ (N)}$	M1 A1 A1	Dim correct equation. Reaction and weight opposing. cao Accept answers rounding to 649.

Q Solution Mark Notes

11(a) Distance moved during constant speed
 $= 15 \times 120 = 1800$ B1

Distance moved during deceleration
 $= 0.5(u+v) \times t, \quad u=15, v=0, t=12$ M1 oe
 $= 0.5(15+0) \times 12 = 90$

$AB = 1890$ (m) A1 cao


11(b) During acceleration
 Use $v=u+at$ with $u=0, a=(\pm)2, t=8$ M1

$v = (\pm)16$ A1

During deceleration
 Use $v=u+at$ with $u=16, v=0, a=(\pm)1.6$ M1

$0 = 16 - 1.6t, \quad t = 10$

Time from B to C = 18 s A1



B1 $v-t$ graph +ve portion with 15, 120, 132
 Labelled
 B1 negative portion
 B1 all correct, units, labels

11(d) Distance $AB = 1890$
 Distance = area under graph

Distance $BC = 0.5 \times 18 \times 16$ M1 used. oe

Distance $CB = 144$

Distance $AC = 1890 - 144$

Distance $AC = 1746$ (m) A1 ft answer from (a)