

MECHANICS 2 (A) TEST PAPER 7 : ANSWERS AND MARK SCHE

1. $a = 2 - 16t^3 = 0$ when $t^3 = 8$ $t = 2$ M1 A1 M1 A1 A1 5

2. $8\pi(3) + 8(10) + 3(13.5) = (11 + 8\pi)\bar{x}$ M1 M1 A1 A1
 $\bar{x} = (24\pi + 120.5) \div (8\pi + 11) = 5.42$ cm M1 A1 A1 7

3. (a) Net resisting force = $2000 - 1600g \sin 7^\circ = 89.1$ N M1 A1 A1
 $1500 = 89.1v$ $v = 16.8$ ms⁻¹ M1 A1
- (b) Now accelerating force = 100 N = $1600a$ $a = 0.0625$ ms⁻² M1 A1 7

4. (a) Displacement = $12\mathbf{i} + 15\mathbf{j}$ Distance = $3\sqrt{41}$ M1 A1 A1
 $|\mathbf{F}| = \sqrt{41}$, so work done = $3\sqrt{41} \times \sqrt{41} = 123$ J M1 A1
- (b) $\mathbf{a} = 4\mathbf{i} + 5\mathbf{j}$ $12\mathbf{i} + 15\mathbf{j} = \frac{1}{2}(4\mathbf{i} + 5\mathbf{j})t^2$ $t = \sqrt{6}$ B1 M1 A1
 $\mathbf{v} = \sqrt{6}(4\mathbf{i} + 5\mathbf{j})$ $|\mathbf{v}| = \sqrt{246} = 15.7$ ms⁻¹ M1 A1 10

5. (a) (i) Momentum : $30mu + 15mu = 6mv + 5mkv$ M1 A1
 $45u = (6 + 5k)v$ $v = \frac{45u}{5k+6}$ M1 A1
- (ii) Elasticity : $(kv - v) / (3u - 5u) = -e$ M1 A1
 $(k - 1)v = (-2u)(-e)$ $v = \frac{2eu}{k-1}$ M1 A1
- (b) $\frac{45u}{5k+6} = \frac{2eu}{k-1}$ $e = \frac{45(k-1)}{2(5k+6)}$ M1 A1
 $0 \leq e \leq 1$, so $0 \leq 45k - 45 \leq 10k + 12$ $1 \leq k \leq \frac{57}{35}$ M1 A1 A1 13

6. (a) $s = \frac{1}{2}gt^2 = \frac{1}{2} \times 9.8 \times 3.3^2 = 53.4$ m M1 A1
- (b) Ball, being lighter, may be affected by air resistance : include this B1 B1
- (c) $x = (7 \cos 30^\circ)t = \frac{7\sqrt{3}}{2}t$ $y = (7 \sin 30^\circ)t - \frac{1}{2}gt^2 = \frac{7}{2}t - 4.9t^2$ M1 A1 M1 A1
- (d) $t = \frac{2x}{7\sqrt{3}}$ $y = \frac{x}{\sqrt{3}} - 4.9\left(\frac{2x}{7\sqrt{3}}\right)^2 = \frac{\sqrt{3}}{3}x - \frac{2}{15}x^2$ M1 A1 A1
- (e) When $x = 10$, $t = 1.65$ $v_x = 3.5\sqrt{3}$, $v_y = 3.5 - 1.65g = -12.67$ M1 A1 A1
 $v = \sqrt{(6.062^2 + 12.67^2)} = 14.0$ ms⁻¹ M1 A1 16

7. (a) $R = mg$, $F = S$ M(B) : $mga \cos \alpha = 2a S \sin \alpha$ B1 B1 M1 A1
 $S = mg / 2 \tan \alpha = \frac{2mg}{3} = F$ A1
Resultant force at B = $\sqrt{[(mg)^2 + \left(\frac{2mg}{3}\right)^2]} = \frac{\sqrt{13}}{3}mg$ M1 A1
- (b) Angle = $\tan^{-1}(3/2) = 56^\circ$ to horizontal M1 A1 A1
- (c) M(B) : $mga \cos \alpha + 6mgx \cos \alpha = 2a S \sin \alpha$ M1 A1 A1
 $S = \frac{2mg(a+6x)}{3a}$ When $S = 2mg$, $a + 6x = 3a$ A1 M1 A1
 $6x = 2a$ $x = \frac{a}{3}$ A1 17