


Take  $g = 9.8 \text{ ms}^{-2}$  and give all answers correct to 3 significant figures where necessary.

1. A particle of mass  $m$  kg moves in a horizontal straight line. Its initial speed is  $u \text{ ms}^{-1}$  and the only force acting on it is a variable resistance of magnitude  $mkv$  N, where  $v \text{ ms}^{-1}$  is the speed of the particle after  $t$  seconds and  $k$  is a constant.

Show that  $v = ue^{-kt}$ . (7 marks)

2. A particle  $P$  of mass  $m$  kg moves in a horizontal circle at one end of a light inextensible string of length 40 cm, as shown. The other end of the string is attached to a fixed point  $O$ . The angular velocity of  $P$  is  $\omega \text{ rad s}^{-1}$ .



If the angle  $\theta$  which the string makes with the vertical must not exceed  $60^\circ$ , calculate the greatest possible value of  $\omega$ . (7 marks)

3. A particle  $P$  of mass  $m$  kg is attached to one end of a light elastic string of natural length 0.5 m and modulus of elasticity  $\frac{mg}{2}$  N. The other end of the string is attached to a fixed point  $O$  and  $P$  hangs vertically below  $O$ .

(a) Find the stretched length of the string when  $P$  rests in equilibrium. (3 marks)


(b) Find the elastic potential energy stored in the string in the equilibrium position. (2 marks)

$P$ , which is still attached to the string, is now held at rest at  $O$  and then lowered gently into its equilibrium position.

(c) Find the work done by the weight of the particle as it moves from  $O$  to the equilibrium position. (2 marks)

(d) Explain the discrepancy between your answers to parts (b) and (c). (1 mark)


4. A particle  $P$ , of mass  $m$  kg, is attached to two light elastic strings, each of natural length  $l$  m and modulus of elasticity  $3mg$  N. The other ends of the strings are attached to the fixed points  $A$  and  $B$ , where  $AB$  is horizontal and  $AB = 2l$  m.



If  $P$  rests in equilibrium vertically below the mid-point of  $AB$ , with each string making an angle  $\theta$  with the vertical, show that

$$\cot \theta - \cos \theta = \frac{1}{6}. \quad \text{(8 marks)}$$

5. A small bead  $P$ , of mass  $m$  kg, can slide on a smooth circular ring, with centre  $O$  and radius  $r$  m, which is fixed in a vertical plane.  $P$  is projected from the lowest point  $L$  of the ring with speed  $\sqrt{(3gr)} \text{ ms}^{-1}$ . When  $P$  has reached a position such that  $OP$  makes an angle  $\theta$  with the downward vertical, as shown, its speed is  $v \text{ ms}^{-1}$ .




- (a) Show that  $v^2 = gr(1 + 2 \cos \theta)$ . (5 marks)
- (b) Show that the magnitude of the reaction  $R$  N of the ring on the bead is given by
- $$R = mg(1 + 3 \cos \theta).$$
- (4 marks)
- (c) Find the values of  $\cos \theta$  when
- (i)  $P$  is instantaneously at rest, (ii) the reaction  $R$  is instantaneously zero. (2 marks)
- (d) Hence show that the ratio of the heights of  $P$  above  $L$  in cases (i) and (ii) is 9 : 8. (3 marks)

6. A light elastic string, of natural length 0.8 m, has one end fastened to a fixed point  $O$ . The other end of the string is attached to a particle  $P$  of mass 0.5 kg. When  $P$  hangs in equilibrium, the length of the string is 1.5 m.

- (a) Calculate the modulus of elasticity of the string. (3 marks)
- $P$  is displaced to a point 0.5 m vertically below its equilibrium position and released from rest.
- (b) Show that the subsequent motion of  $P$  is simple harmonic, with period 1.68 s. (5 marks)
- (c) Calculate the maximum speed of  $P$  during its motion. (3 marks)
- (d) Show that the time taken for  $P$  to first reach a distance 0.25 m from the point of release is 0.28 s, to 2 significant figures. (4 marks)

7. (a) Show that the centre of mass of a uniform solid hemisphere of radius  $r$  is at a distance  $\frac{3r}{8}$  from the centre  $O$  of the plane face. (7 marks)

The figure shows the vertical cross-section of a rough solid hemisphere at rest on a rough inclined plane inclined at an angle  $\theta$  to the horizontal, where  $\sin \theta = \frac{3}{10}$ .



- (b) Indicate on a copy of the figure the three forces acting on the hemisphere, clearly stating what they are, and paying special attention to their lines of action. (3 marks)
- (c) Given that the plane face containing the diameter  $AB$  makes an angle  $\alpha$  with the vertical, show that  $\cos \alpha = \frac{4}{5}$ . (6 marks)