

MECHANICS 3 (A) TEST PAPER 7 : ANSWERS AND MARK SCHEMES

1. $0.27 = mr\omega^2 = 0.6r(1.5^2)$ $r = 0.2 \text{ m}$ M1 A1 A1

2. Vert. : $R + T \sin 60^\circ = 0.7g$ Horiz. : $T \cos 60^\circ = F$ M1 A1 M1 A1
 $F = 0.25R$, so $0.5T = 0.25(6.86 - 0.866T)$ $T = 2.394$ M1 A1
 $T = 6.86x + 0.5$, so $x = 1.196 + 6.86 = 0.174 \text{ m} \approx 17 \text{ cm}$ M1 A1 8

3. (a) Energy conserved, so $mg(1 + \cos 30^\circ) = \frac{1}{2}mv^2$ M1 A1
Hence $v^2 = 2g(1 + \cos 30^\circ) = g(2 + \sqrt{3})$ A1
 $a_y = \frac{v^2}{r} = g(2 + \sqrt{3})$ towards O; $a_x = 0$ (no horizontal force) A1 A1
(b) At bottom, $R = \frac{mv^2}{r} + mg = mg(2 + \sqrt{3}) + mg = 0.1g(3 + \sqrt{3})$ M1 A1 A1 8

4. (a) $v^2 = n^2(a^2 - x^2)$ $36 = n^2(a^2 - 16)$, $16 = n^2(a^2 - 36)$ M1 A1 A1
 $36(a^2 - 36) = 16(a^2 - 16)$ $20a^2 = 1040$ $a = 7.21 \text{ m}$ M1 A1 A1
(b) $n^2 = 1$ $n = 1$ Period = $\frac{2\pi}{n} = 2\pi \text{ s}$ M1 A1 A1 9

5. (a) $x^2 + y^2 = r^2$ $\bar{x} \int_0^r \pi y^2 dx = \int_0^r \pi xy^2 dx$ B1 M1 A1
 $\bar{x} \int_0^r r^2 - x^2 dx = \pi \int_0^r r^2 x - x^3 dx$ $\frac{2x^2}{3} \bar{x} = \frac{r^4}{4}$ $\bar{x} = \frac{3r}{8}$ M1 A1 A1 A1
(b) $M(O) : \frac{2}{3}\pi r^3 \cdot \frac{3r}{8} = \pi \left(\frac{3r}{4}\right)^2 \cdot kr \cdot \frac{kr}{2}$ $k^2 = \frac{8}{9}$ $k = \frac{2}{3}\sqrt{2}$ M1 A1 A1 A1
(c) $\tan \theta = \frac{3r}{4} + \frac{2r\sqrt{2}}{3} = \frac{9}{8\sqrt{2}}$ $\theta = 38.5^\circ$ M1 A1 A1 14

6. (a) $g = \frac{kM(1)}{R^2}$, so $k = \frac{gR^2}{M}$ M1 A1
(b) $\frac{mv^2}{r} = \frac{kMm}{r^2}$ $v^2 = \frac{gR^2}{M} \cdot \frac{M}{r} = \frac{gR^2}{r}$ $T = \frac{2\pi r}{v} = 2\pi \sqrt{\frac{r^3}{gR^2}}$ M1 A1 M1 A1
(c) Diagram (d) Along XE, as X in circular orbit so central force B1; B2
(e) $\frac{mv^2}{r} = \frac{gR^2}{M} \left(\frac{m^2}{3r^2} \cos 30^\circ + \frac{mM}{r^2} + \frac{m^2}{3r^2} \cos 30^\circ \right)$ M1 A1 A1
 $\frac{mv^2}{r} = \frac{gR^2 m}{Mr^2} \left(\frac{m\sqrt{3}}{3} \times 2 + M \right)$ $v^2 = \frac{gR^2}{Mr} \left(M + \frac{m\sqrt{3}}{3} \right)$ M1 A1
 $T_1 = \frac{2\pi r}{v} = 2\pi r \sqrt{\frac{3Mr}{gR^2(3M+m\sqrt{3})}} = 2\pi \sqrt{\frac{r^3}{gR^2} \frac{3M}{3M+m\sqrt{3}}} = T \sqrt{\frac{3M}{3M+m\sqrt{3}}}$ M1 A1 16

7. (a) In eq. position, $\frac{\lambda}{3} l = mg$ $\lambda = 3mg$ M1 A1
At depth x below eq. position, $mg - T = mx$ M1
 $mg - \frac{3mg}{3l}(l+x) = mx$ $x = -\frac{g}{l}x$ SHM, with $\omega^2 = \frac{g}{l}$ A1 A1
(b) $x = 2l \cos \omega t$ When $x = -l$, $\omega t = \frac{2\pi}{3}$ $t = \frac{2\pi}{3\omega} = \frac{2\pi}{3} \sqrt{\frac{l}{g}}$ B1 M1 A1 A1
(c) When released, E.P.E. = $\frac{3mg(9l^2)}{2(3l)} = \frac{9mg^2 l}{2}$ M1 A1
At max. height H , P.E. = $mgH = \frac{9mg^2 l}{2}$ $H = \frac{9l}{2}$ M1 A1
(d) When slack, $v^2 = 3gl$ $0 = \sqrt{(3gl) - gt_H^2}$ $t_H = \sqrt{\frac{3l}{g}}$ $T_h = \frac{2\pi}{3} \sqrt{\frac{l}{g}} + \sqrt{3} \sqrt{\frac{l}{g}}$ M1 A1 M1 A1 17