


MECHANICS (A) UNIT 3

TEST PAPER

Take $g = 9.8 \text{ ms}^{-2}$ and give all answers correct to 3 significant figures where necessary.


1. A cyclist travels on a banked track inclined at 8° to the horizontal. He moves in a horizontal circle of radius 10 m at a constant speed of $v \text{ ms}^{-1}$. If there is no sideways frictional force on the cycle, calculate the value of v . (6 marks)

2. The figure shows a particle P , of mass 0.8 kg, attached to the ends of two light elastic strings. AP has natural length 20 cm and modulus of elasticity $\lambda \text{ N}$. BP has natural length 20 cm and modulus of elasticity $\mu \text{ N}$. A and B are fixed to points on the same horizontal level so that $AB = 50 \text{ cm}$. When P is suspended in equilibrium, $AP = 30 \text{ cm}$ and $BP = 40 \text{ cm}$. Calculate the values of λ and μ . (9 marks)




3. Suraiya, whose mass is $m \text{ kg}$, takes a running jump into a swimming pool so that she begins to swim in a straight line with speed 0.2 ms^{-1} . She continues to move in the same straight line, the only force acting on her being a resistance of magnitude $mv^2 \sin\left(\frac{t}{100}\right) \text{ N}$, where $v \text{ ms}^{-1}$ is her speed at time t seconds after entering the pool and $0 \leq t \leq 50\pi$.
 - (a) Find an expression for v in terms of t . (7 marks)
 - (b) Calculate her greatest and least speeds during her motion. (3 marks)

4. A uniform lamina is in the shape of the region enclosed by the coordinate axes and the curve with equation $y = 1 + \cos x$, as shown.



- (a) Show by integration that the centre of mass of the lamina is at a distance $\frac{\pi^2 - 4}{2\pi}$ from the y -axis. (9 marks)
- Given that the centre of mass is at a distance 0.75 units from the x -axis, and that P is the point $(0, 2)$ and O is the origin $(0, 0)$,
- (b) find, to the nearest degree, the angle between the line OP and the vertical when the lamina is freely suspended from P . (3 marks)

5. A particle P , of mass 0.5 kg, rests on the surface of a rough horizontal table. The coefficient of friction between P and the table is 0.5. P is connected to a particle Q , of mass 0.2 kg, by a light inextensible string passing through



[Turn over ...]

MECHANICS 3 (A) TEST PAPER 10 Page 2

5. continued ...

a small smooth hole at a point O on the table, such that the distance OQ is 0.4 m. Q moves in a horizontal circle while P remains in limiting equilibrium.


(a) Calculate the angle θ which OQ makes with the vertical. (4 marks)

(b) Show that the speed of Q is 1.33 ms^{-1} . (3 marks)

The motion is altered so that Q hangs at rest below O and P moves in a horizontal circle on the table with speed 0.84 ms^{-1} , at a constant distance r m from O but tending to slip *away* from O .

(c) Find the value of r . (5 marks)

6. The figure shows a swing consisting of two identical vertical light springs attached symmetrically to a light horizontal cross-bar and supported from a strong fixed horizontal beam. When a mass of 24 kg is placed at the mid-point of the cross-bar, both springs extend by 30 cm to the position A , as shown.



Each spring has natural length l m and modulus of elasticity λ N.

(a) Show that $\lambda = 392l$. (2 marks)


The 24 kg mass is left on the bar and the bar is then displaced downwards by a further 20 cm.

(b) Prove that the system comprising the bar and the mass now performs simple harmonic motion with the centre of oscillation at the level A . (5 marks)

(c) Calculate the number of oscillations made per second in this motion. (3 marks)

(d) Find the maximum acceleration which the mass experiences during the motion. (2 marks)

7. A particle P of mass m kg is attached to points C and D on the same horizontal level by means of two light inextensible strings CP and DP , both of length 40 cm. P is projected with speed $u \text{ ms}^{-1}$ so as to move in a vertical circle in a plane perpendicular to CD , so that angle $PCD = \text{angle } PDC = \theta$ throughout the motion.



If u is just large enough for the strings to remain taut as P describes this circular path,

(a) show that $u^2 = 2g \sin \theta$. (8 marks)

The string DP breaks when P is at its lowest point. P then immediately starts to move in a horizontal circle on the end of the string CP .

(b) Prove that $\tan \theta = \frac{1}{5}\sqrt{5}$. (6 marks)