

1. SIGHT OF  $x^{\frac{1}{2}}$  &  $x^{-\frac{1}{2}}$  B1  
 $4x^{\frac{3}{2}} - 12x^{\frac{1}{2}}$  M1  
 CORRECT METHOD E.g.  $(108-36)-(4-12)$  OR  $72+8$  M1  
 80 c.a.o A1

2. a)  $2(-2)^3 - 7(-2)^2 - 5(-2) + 4$  OR  $-16 - 28 + 10 + 4$  M1  
 OR SENSIBLE ATTEMPT ON DIVISION M1  
 -30 c.a.o A1

b)  $2 \times 4^3 - 7 \times 4^2 - 5 \times 4 + 4$  OR  $128 - 112 - 20 + 4$  M1  
 (DO NOT ACCEPT DIVISION HERE)  
 OBTAINS ZERO + CONCLUSION A1

c)  $(x-4)(2x^2+bx+c)$  M1  
 $(x-4)(2x^2+x-1)$  M1  
 $(x-4)(2x-1)(x+1)$  A1

3 a)  $\frac{1}{6}$  o.e B1

b)  $\frac{90}{1 - \frac{1}{6}}$  M1  
 108 c.a.o A1

4.  $3x^2 - 10x + 3$  B1  
 $3x^2 - 10x + 3 < 0$  OR  $f(x) < 0$  B1

$(3x-1)(x-3)$  M1  
 SIGHT OF 3 &  $\frac{1}{3}$  AS "CRITICAL VALUES" A1

~~$\frac{1}{3} < x < 3$~~  OR EQUIVALENT METHOD M1  
 $\frac{1}{3} < x < 3$  OR  $\frac{1}{3} \leq x \leq 3$  A1

5. USE OF  $1 - \cos^2 3x$  (MUST BE IN  $3x$ ) **B1**  
 SIMPLIFIED TO 3 TERM QUADRATIC e.g.  $3\cos^2 3x + 7\cos 3x + 2$  **M1**  
 $(3\cos 3x + 1)(\cos 3x + 2)$  OR SIMILAR INPUT FACTORIZATION **M1**  
 SIGHT OF  $-\frac{1}{3}$  AND  $-2$  **A1**  
 SIGHT OF  $109.47\dots$  **A1**  
 SIGHT OF  $250.53\dots$  **A1**  
 $36.5^\circ, 156.5^\circ, 83.5^\circ$  **A2** -1 e e o o

6. a) SIGHT OF "tan" OR  $\tan \theta = \frac{2.2}{0.9}$  **M1**  
 $1.18247\dots$  **A1**

$\pi - 2 \times "1.18247\dots"$  **M1**  
 SIGNS  $0.7766$  **A1**  $\uparrow$  **M1**  
 dtp

b) USE OF PYTHAGORAS **M1**  
 SIGHT OF  $2.37697$  **A1**  
 $"2.37697" \times 0.7766$  **M1**  
 A.W.R.T  $8.05$  c.a.o **A1**

c)  $\frac{1}{2} \times ("2.376\dots")^2 \times 0.7766$  OR  $2.194\dots$  **M1**  
 $\frac{1}{2} \times 0.4 \times 2.2$  OR  $0.99$  **M1**  
 A.W.R.T  $4.17 - 4.18$  **A1**

7.  $\frac{1}{2} \times 3 \times 18$  or 27 M1

SLANT  $\int_{-1}^2 18 - x - x^4 dx$  B1

$18x - \frac{1}{2}x^2 - \frac{1}{5}x^5$  (ALLOW ONE ERROR) M1

$(36 - 2 - \frac{30}{5}) - (-18 - \frac{1}{2} + \frac{1}{5})$  or  $\frac{138}{5} - (-\frac{183}{10})$  M1

$\frac{459}{10}$  or 45.9 A1

$\frac{189}{10}$  or 18.9 A1

8.  $1 + nkx + \frac{1}{2}n(n-1)k^2x^2$  o.e. M2

ATTEMPTS SOLUTION OF  $\frac{1}{2}n(n-1) = 120$  M1

$n = 16$  SEEN A1

$n/k = 40$  B1

$k = \frac{5}{2}$  o.e. A1 ~~It~~ from "that n"

9.  $\frac{\log_4(x-4)}{\log_4 16}$  B1

IMPLIES  $\log_4 16 = 2$  B1

$2(\log_4 x - \log_4(x-4)) = 2$  M1

$\log_4 \left( \frac{x^2}{x-4} \right)$  A1

$\frac{x^2}{x-4} = 16$  M1

$x^2 - 16x + 64$  A1

$(x-8)^2$  M1

$x = 8$  c.a.o. A1

10. a)  $(4, 2)$  B1

ATTEMPT TO FIND DISTANCE FROM "THESE C" TO ANY CORNER M1  
 $\sqrt{10}$  A1

b) "ATTEMPTS"  $|SR|$  OR  $|PQ|$  (OR THEIR MIDPOINTS) M1

SHOWS  $\sqrt{20}$  OR  $2\sqrt{5}$  A1

$(x-4)^2 + (y-2)^2 = 5$  A1 A1

11.

$\frac{dy}{dx} = 3x^2 + 2ax + b$  B1

$-5 = (-1)^3 + a(-1)^2 + b(-1) - 10 = -5$  o.e. B1

$3(-1)^2 + 2a(-1) + b = 0$  o.e. B1

$a - b = 6$   
OR  $2a - b = 3$  ) A1

SOLVES AND OBTAINS  $a = -3$  both A1  
 $b = -9$

ATTEMPT TO FACTORIZE e.g.  $(x+1)(x-3)$  M1

$Q(3, -37)$  A1

$\frac{d^2y}{dx^2} = 6a - 6$  M1

$\left. \frac{d^2y}{dx^2} \right|_{x=3} = 12 > 0$  & STATES (LOCAL) MIN A1