

1Y6B - FPI PAPER Q - QUESTION 1

a) using $|zw| = |z||w|$

$$\Rightarrow |z_1 z_3| = 16$$

$$\Rightarrow |z_1| |z_3| = 16$$

$$\Rightarrow |2-2i| |z_3| = 16$$

$$\Rightarrow \sqrt{4+4} |z_3| = 16$$

$$\Rightarrow \sqrt{8} |z_3| = 16$$

$$\Rightarrow \sqrt{2} \sqrt{8} |z_3| = 16\sqrt{2}$$

$$\Rightarrow 4 |z_3| = 16\sqrt{2}$$

$$\Rightarrow |z_3| = 4\sqrt{2}$$

b) using $\arg\left(\frac{z}{w}\right) = \arg z - \arg w$

$$\Rightarrow \arg\left(\frac{z_3}{z_2}\right) = \frac{7\pi}{12}$$

$$\Rightarrow \arg z_3 - \arg z_2 = \frac{7\pi}{12}$$

$$\Rightarrow \arg z_3 - \arg(\sqrt{3}+i) = \frac{7\pi}{12}$$

$$\Rightarrow \arg z_3 - \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{7\pi}{12}$$

$$\Rightarrow \arg z_3 - \frac{\pi}{6} = \frac{7\pi}{12}$$

$$\Rightarrow \arg z_3 = \frac{3\pi}{4}$$

1YGB - FPI PART 1 - QUESTIONS

c) finally if $z_3 = a + bi$, $|z_3| = 4\sqrt{2}$, $\arg z_3 = \frac{3\pi}{4}$

$$\begin{aligned} |a + bi| &= 4\sqrt{2} \\ \sqrt{a^2 + b^2} &= 4\sqrt{2} \\ a^2 + b^2 &= 32 \end{aligned}$$

$$\begin{aligned} \arg z_3 &= \frac{3\pi}{4} \\ \arctan \frac{b}{a} + \pi &= \frac{3\pi}{4} \\ \arctan \frac{b}{a} &= -\frac{\pi}{4} \end{aligned}$$

(OR SIMPLY INSPECTION)


$$\begin{aligned} \frac{b}{a} &= \tan\left(-\frac{\pi}{4}\right) \\ \frac{b}{a} &= -1 \\ b &= -a \end{aligned}$$

$$\begin{aligned} a^2 + a^2 &= 32 \\ 2a^2 &= 32 \\ a^2 &= 16 \\ a &= -4 \end{aligned}$$

(AS z_3 IS IN THE 2ND QUADRANT)
a b = +4

finally $\frac{z_3}{z_1} = \frac{-4 + 4i}{2 - 2i} = \frac{-2(2 - 2i)}{2 - 2i} = -2$ ✓

ALTERNATIVE FOR PART C



$$\begin{aligned} z_3 &= r(\cos \theta + i \sin \theta) \\ z_3 &= 4\sqrt{2} \left(\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4} \right) \\ z_3 &= 4\sqrt{2} \left(-\frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}} \right) \\ z_3 &= -4 + 4i \quad \text{etc} \end{aligned}$$

1YGB - P1 PAPER Q - QUESTION 2

OBTAIN RELATIONSHIPS FOR THE ROOTS OF THE GIVEN QUADRATIC

$$x^2 + 2x + 3 = 0 \implies \begin{cases} \alpha + \beta = \frac{-2}{1} = -2 & \leftarrow -\frac{b}{a} \\ \alpha\beta = \frac{3}{1} = 3 & \leftarrow \frac{c}{a} \end{cases}$$

PROCEED AS FOLLOWS

$$\begin{aligned} A &= \alpha - \frac{1}{\beta^2} \\ B &= \beta - \frac{1}{\alpha^2} \end{aligned}$$

$$\begin{aligned} \bullet \underline{A+B} &= \left(\alpha - \frac{1}{\beta^2}\right) + \left(\beta - \frac{1}{\alpha^2}\right) = (\alpha + \beta) - \left(\frac{1}{\alpha^2} + \frac{1}{\beta^2}\right) \\ &= (\alpha + \beta) - \frac{\beta^2 + \alpha^2}{\alpha^2\beta^2} = (\alpha + \beta) - \frac{\alpha^2 + \beta^2}{(\alpha\beta)^2} \\ &= (\alpha + \beta) - \frac{(\alpha + \beta)^2 - 2\alpha\beta}{(\alpha\beta)^2} = -2 - \frac{(-2)^2 - 2 \times 3}{3^2} = \underline{\underline{-\frac{16}{9}}} \end{aligned}$$

$$\begin{aligned} \bullet \underline{AB} &= \left(\alpha - \frac{1}{\beta^2}\right)\left(\beta - \frac{1}{\alpha^2}\right) = \alpha\beta - \frac{1}{\alpha} - \frac{1}{\beta} + \frac{1}{\alpha^2\beta^2} \\ &= \alpha\beta - \left(\frac{1}{\alpha} + \frac{1}{\beta}\right) + \frac{1}{(\alpha\beta)^2} = \alpha\beta - \left(\frac{\beta + \alpha}{\alpha\beta}\right) + \frac{1}{(\alpha\beta)^2} \\ &= 3 - \frac{-2}{3} - \frac{1}{3^2} = 3 + \frac{2}{3} + \frac{1}{9} = \underline{\underline{\frac{34}{9}}} \end{aligned}$$

HENCE THE REQUIRED EQUATION WILL BE

$$\implies x^2 - (A+B)x + AB = 0$$

$$\implies x^2 - \left(-\frac{16}{9}\right)x + \frac{34}{9} = 0$$

$$\implies \underline{\underline{9x^2 + 16x + 34 = 0}}$$

NYGB - FPI PAPER Q - QUESTION 3

$$f(n) = 4^n + 6n - 1, \quad n \in \mathbb{N}$$

BASE CASE

$$f(1) = 4^1 + 6 \times 1 - 1 = 4 + 6 - 1 = 9, \quad \text{IF RESULT HOLDS FOR } n=1$$

INDUCTIVE HYPOTHESIS

SUPPOSE THAT THE RESULT HOLDS FOR $n=k, k \in \mathbb{N}$ IF $f(k) = 9m$ WHERE $m \in \mathbb{N}$

$$\Rightarrow f(k+1) - f(k) = [4^{k+1} + 6(k+1) - 1] - [4^k + 6k - 1]$$

$$\Rightarrow f(k+1) - 9m = 4^{k+1} + 6k + 6 - 1 - 4^k - 6k + 1$$

$$\Rightarrow f(k+1) - 9m = 4^{k+1} - 4^k + 6$$

$$\Rightarrow f(k+1) - 9m = 4 \times 4^k - 4^k + 6$$

$$\Rightarrow f(k+1) - 9m = 3 \times 4^k + 6$$

$$\Rightarrow f(k+1) = 9m + 6 + 3[f(k) - 6k + 1]$$

$$\Rightarrow f(k+1) = 9m + 6 + 3f(k) - 18k + 3$$

$$\Rightarrow f(k+1) = 9m - 18k + 9 + 3(9m)$$

$$\Rightarrow f(k+1) = 36m - 18k + 9$$

$$\Rightarrow f(k+1) = 9[4m - 2k + 1]$$

$$\begin{aligned} f(k) &= 4^k + 6k - 1 \\ 4^k &= f(k) - 6k + 1 \end{aligned}$$

CONCLUSION

IF THE RESULT HOLDS FOR $n=k, k \in \mathbb{N}$, THEN IT ALSO HOLDS FOR $n=k+1$
SINCE THE RESULT HOLDS FOR $n=1$, THEN IT MUST HOLD FOR ALL n

1YGB - FPI PAPER Q - QUESTION 4

AS THE POLYNOMIAL EQUATION HAS REAL COEFFICIENTS ANY SOLUTIONS MUST APPEAR IN CONJUGATE PAIRS, SO $z = 3 \pm i$ ARE SOLUTIONS

$$\begin{aligned} [z - (3+i)][z - (3-i)] &= [(z-3) - i][(z-3) + i] \\ &= (z-3)^2 - i^2 \\ &= z^2 - 6z + 9 + 1 \\ &= z^2 - 6z + 10 \end{aligned}$$

BY LONG DIVISION

$$\begin{array}{r} z^2 - 6z + 10 \quad | \quad \begin{array}{r} 2z^2 - 2z + 1 \\ \hline 2z^4 - 14z^3 + 33z^2 - 26z + 10 \\ -2z^4 + 12z^3 - 20z^2 \\ \hline -2z^3 + 13z^2 - 26z + 10 \\ +2z^3 - 12z^2 + 20z \\ \hline z^2 - 6z + 10 \\ -z^2 + 6z - 10 \\ \hline \end{array} \end{array}$$

Hence $2z^2 - 2z + 1 = 0$

$$4z^2 - 4z + 2 = 0$$

$$4z^2 - 4z + 1 = -1$$

$$(2z-1)^2 = -1$$

$$2z-1 = \pm i$$

$$2z = 1 \pm i$$

$$z = \frac{1}{2} \pm \frac{1}{2}i$$

∴ THE FULL SOLUTION SET IS $3+i, 3-i, \frac{1}{2} + \frac{1}{2}i, \frac{1}{2} - \frac{1}{2}i$

- 1 -

LYGB - FPI PAPER Q - QUESTION 5


a)

$$\underline{A} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}$$

$$\begin{aligned} \underline{i} &\mapsto \underline{i} \\ \underline{j} &\mapsto \underline{k} \\ \underline{k} &\mapsto -\underline{j} \end{aligned}$$

$\det \underline{A} = 1$ (NO REFLECTION)

ROTATION ABOUT THE z AXIS,
BY 90°, ANTICLOCKWISE IN A
RIGHT HAND SENSE




$$\underline{B} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{aligned} \underline{i} &\mapsto -\underline{i} \\ \underline{j} &\mapsto -\underline{j} \\ \underline{k} &\mapsto \underline{k} \end{aligned}$$

$\det \underline{B} = -1$ (REFLECTION INVOLVED)

REFLECTION ABOUT THE xz PLANE



b) COMPOSE IN THE CORRECT ORDER

$$\underline{C} = \underline{B}\underline{A} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

$\det \underline{C} = -1$ (REFLECTION INVOLVED)


LYGB - FPI PAPER Q - QUESTION 5

$$i \mapsto \bar{i}$$

$$j \mapsto \bar{k}$$

$$k \mapsto \bar{j}$$

LOOKING AT THE yz PLANE,
FROM THE "POSITIVE" x



\therefore REFLECTION ABOUT THE PLANE $y=z$

- 1 -

YGB - FPI PAPER Q - QUESTION 6

USING THE STANDARD SUMMATION FORMULAS

$$\sum_{r=1}^k r = \frac{1}{2}k(k+1)$$

$$\sum_{r=1}^k r^3 = \frac{1}{4}k^2(k+1)^2$$

Hence we now have

$$\begin{aligned} \sum_{r=n}^{2n} (r^3 - 2r) &= \sum_{r=n}^{2n} r^3 - 2 \sum_{r=n}^{2n} r \\ &= \left[\frac{1}{4}(2n)^2(2n+1)^2 - \frac{1}{4}(n-1)^2n^2 \right] - 2 \left[\frac{1}{2}(2n)(2n+1) - \frac{1}{2}(n-1)n \right] \\ &= n^2(2n+1)^2 - \frac{1}{4}n^2(n-1)^2 - 2n(2n+1) + n(n-1) \\ &= \frac{1}{4}n \left[4n(2n+1)^2 - n(n-1)^2 - 8(2n+1) + 4(n-1) \right] \end{aligned}$$

As it will be a mess to expand to full w.r.t. it's best to factorize the terms inside the bracket in primes.

$$\begin{aligned} &= \frac{1}{4}n \left[n \left[4(2n+1)^2 - (n-1)^2 \right] - 4 \left[4n+2+n-1 \right] \right] \\ &= \frac{1}{4}n \left[n \left[\underbrace{4(2n+1)^2 - (n-1)^2}_{\text{diff. of squares}} \right] - 4(3n+3) \right] \\ &= \frac{1}{4}n \left[n \left[2(2n+1) - (n-1) \right] \left[2(2n+1) + (n-1) \right] - 12(n+1) \right] \\ &= \frac{1}{4}n \left[n \left[4n+2-n+1 \right] \left[4n+2+n-1 \right] - 12(n+1) \right] \\ &= \frac{1}{4}n \left[n(3n+3)(5n+1) - 12(n+1) \right] \\ &= \frac{1}{4}n \left[3n(n+1)(5n+1) - 12(n+1) \right] \end{aligned}$$

1YGB - FPI PAPER Q - QUESTION 6

$$\begin{aligned}
&= \frac{3}{4}n [n(n+1)(5n+1) - 4(n+1)] \\
&= \frac{3}{4}n(n+1)[n(5n+1) - 4] \\
&= \frac{3}{4}n(n+1)(5n^2+n-4) \\
&= \frac{3}{4}n(n+1)(5n-4)(n+1) \\
&= \underline{\underline{\frac{3}{4}n(n+1)^2(5n-4)}}
\end{aligned}$$

ALTERNATIVE BY EXPANDING A CUBIC AFTER THE INITIAL FACTORING

$$\begin{aligned}
\dots &= \frac{1}{4}n [4n(2n+1)^2 - n(n-1)^2 - 8(2n+1) + 4(n-1)] \\
&= \frac{1}{4}n [16n^3 + 16n^2 + 4n - n^3 + 2n^2 - n - 16n - 8 + 4n - 4] \\
&= \frac{1}{4}n [15n^3 + 18n^2 - 9n - 12] \\
&= \frac{3}{4}n [5n^3 + 6n^2 - 3n - 4]
\end{aligned}$$

LOOKING FOR FACTORS

$$n=1 \quad 5+6-3-4 \neq 0$$

$$n=-1 \quad -5+6+3-4=0 \quad \text{Hence } (n+1) \text{ is a factor}$$

LONG DIVIDE

| | | |
|-------|------------|------------------------|
| | | $5n^3 + 6n^2 - 3n - 4$ |
| $n+1$ | $5n^2+n-4$ | $-5n^3 - 5n^2$ |
| | | $n^2 - 3n - 4$ |
| | | $-n^2 - n$ |
| | | $-4n - 4$ |
| | | $4n + 4$ |

$$\dots = \frac{3}{4}n(n+1)(5n^2+n-4)$$

$$= \frac{3}{4}n(n+1)(5n-4)(n+1)$$

$$= \underline{\underline{\frac{3}{4}n(n+1)^2(5n-4)}}$$

✓ BRUCE

- 1 -
IYGB - FPI PAPER Q - QUESTION 7

a) VOLUME OF REVOLUTION IN CARTESIAN, ABOUT THE x AXIS

$$V = \pi \int_{x_1}^{x_2} (y(x))^2 dx = \pi \int_2^6 \left(\frac{x+1}{\sqrt{x-1}} \right)^2 dx = \pi \int_2^6 \frac{(x+1)^2}{x-1} dx$$

BY SUBSTITUTION

OR

MANIPULATION

• $u = x-1$ OR $x = u+1$

$$\frac{du}{dx} = 1$$

$$du = dx$$

• $x=2 \mapsto u=1$

$x=6 \mapsto u=5$

$$\Rightarrow V = \pi \int_1^5 \frac{(x+1)^2}{u} du$$

$$\Rightarrow V = \pi \int_1^5 \frac{(u+1)^2}{u} du$$

$$\Rightarrow V = \pi \int_1^5 \frac{(u+2)^2}{u} du$$

$$\Rightarrow V = \pi \int_1^5 \frac{u^2 + 4u + 4}{u} du$$

$$\Rightarrow V = \pi \int_1^5 \left(u + 4 + \frac{4}{u} \right) du$$

$$\Rightarrow V = \pi \left[\frac{1}{2}u^2 + 4u + 4 \ln|u| \right]_1^5$$

$$\Rightarrow V = \pi \left[\left(\frac{25}{2} + 20 + 4 \ln 5 \right) - \left(\frac{1}{2} + 4 + 4 \ln 1 \right) \right]$$

$$\Rightarrow V = \pi \left[28 + 4 \ln 5 \right]$$

$$\Rightarrow V = \pi \int_2^6 \frac{[(x-1)+2]^2}{x-1} dx$$

$$\Rightarrow V = \pi \int_2^6 \frac{(x-1)^2 + 4(x-1) + 4}{x-1} dx$$

$$\Rightarrow V = \pi \int_2^6 \left(\frac{(x-1)^2}{x-1} + \frac{4(x-1)}{x-1} + \frac{4}{x-1} \right) dx$$

$$\Rightarrow V = \pi \int_2^6 \left(x-1 + 4 + \frac{4}{x-1} \right) dx$$

$$\Rightarrow V = \pi \int_2^6 \left(x+3 + \frac{4}{x-1} \right) dx$$

$$\Rightarrow V = \pi \left[\frac{1}{2}x^2 + 3x + 4 \ln|x-1| \right]_2^6$$

$$\Rightarrow V = \pi \left[(18 + 18 + 4 \ln 5) - (2 + 6 + 4 \ln 1) \right]$$


$$\Rightarrow V = \pi \left[28 + 4 \ln 5 \right]$$

AS OPPOSITE

IYGB - FPI PAPER Q - QUESTION 7

b)

LOOKING AT THE SIMILAR SHAPES



$$V = \pi(28 + 4\ln 5)$$

$$V = ?$$

$$V = V \times (\text{SCALE FACTOR})^3$$

$$V = \pi(28 + 4\ln 5) \times (1.5)^3$$

$$V \approx \underline{365}$$

1YGB - FPI PAPER Q - QUESTION 8

a) ELIMINATE TO CARTESIAN FIRST

$$\left. \begin{aligned} x &= 4 + \lambda + 5\mu \\ y &= 8 + 2\lambda - 4\mu \\ z &= -5 + \lambda + 7\mu \end{aligned} \right\} \Rightarrow \lambda = z + 5 - 7\mu$$

SUBSTITUTE INTO THE FIRST TWO EQUATIONS

$$\left. \begin{aligned} x &= 4 + (z + 5 - 7\mu) + 5\mu \\ y &= 8 + 2(z + 5 - 7\mu) - 4\mu \end{aligned} \right\} \Rightarrow$$

$$\begin{aligned} x &= 9 + z - 2\mu && \times 9 \\ y &= 18 + 2z - 18\mu && \times (-1) \end{aligned}$$

$$\left. \begin{aligned} 9x &= 81 + 9z - 18\mu \\ -y &= -18 - 2z + 18\mu \end{aligned} \right\} \text{ADDING}$$

$$9x - y = 63 + 7z$$

$$9x - y - 7z = 63$$

FINALLY THE EQUATION OF THE PLANE CAN BE WRITTEN AS

$$(9, -1, -7) \cdot (x, y, z) = 63$$

$$\underline{\underline{\underline{\underline{r \cdot \begin{pmatrix} 9 \\ -1 \\ -7 \end{pmatrix} = 63}}}}}$$

b) DETERMINE THE EQUATION OF A LINE THROUGH P (12, -1, 44)
& IN THE DIRECTION OF THE NORMAL

$$r = (12, -1, 44) + t(9, -1, -7)$$

$$(x, y, z) = (9t + 12, -t - 1, -7t + 44)$$

1YGB - FPI PAPER Q - QUESTION 8

SOLVE SIMULTANEOUSLY WITH THE EQUATION OF THE PLANE

$$x = 9t + 12$$

$$y = -t - 1$$

$$z = 44 - 7t$$

$$9x - y - 7z = 63$$

$$\Rightarrow 9(9t + 12) - (-t - 1) - 7(44 - 7t) = 63$$


$$\Rightarrow 81t + t + 49t + 108 + 1 - 308 = 63$$

$$\Rightarrow 131t = 262$$

$$\Rightarrow t = 2$$

USING $t = 4$ WE OBTAIN THE REFLECTION

$$P'(48, -5, 16)$$



ALTERNATIVE/VARIATION

- USE $t = 2$ TO FIND $Q(30, -3, 30)$

↑
THIS USES ON
 $9x - y - 7z = 63$

- THEN USE "MIDPOINT PATTERNS"

