

- 1 -

IYGB - SYNOPTIC PAPER F - QUESTION 1

$$\underline{f(x) = f(x+2)}$$

$$\Rightarrow x^2 - 7x + 6 = (x+2)^2 - 7(x+2) + 6$$

$$\Rightarrow x^2 - 7x = x^2 + 4x + 4 - 7x - 14$$

$$\Rightarrow 10 = 4x$$

$$\Rightarrow x = \frac{5}{2}$$

1YGB - SYNOPTIC PAPER F - QUESTION 2

a) START BY UNRAILING THE EQUATION

$$\Rightarrow H = kt^n$$

$$\Rightarrow \log H = \log(kt^n)$$

$$\Rightarrow \log H = \log k + \log t^n$$

$$\Rightarrow \log H = \log k + n \log t$$

$$\Rightarrow \log H = n(\log t) + \log k$$

$$Y \quad m \quad X \quad C$$

b) MODIFYING THE TABLE OF VALUES

t	5	10	20	40	50
H	4.1	8.5	18.0	42.0	50.0

X = log t	0.70	1	1.30	1.60	1.70
Y = log H	0.61	0.93	1.26	1.62	1.70

PUTTING THESE VALUES ACCURATELY

1YGB - SYNOPTIC PAPER F - QUESTION 2

$Y = \log H$

c) ● "y intercept is $\log k$

$$\log_{10} k \approx -0.18$$

$$k \approx 10^{-0.18}$$


$$k \approx 0.66$$

● GRADIENT IS n

$(0, -0.18)$ & $(1, 0.93)$

$$n \approx \frac{0.93 - (-0.18)}{1}$$

$$n \approx 1.1$$



1YGB - SYNOPTIC PAPER F - QUESTION 3

COMPLETING THE SQUARE

$$y = x^2 + 8x + 12$$

$$y = (x+4)^2 - 4^2 + 12$$

$$y = (x+4)^2 - 16 + 12$$

$$y = (x+4)^2 - 4$$

NOW WE HAVE

$$x^2 \xrightarrow{\text{REPLACE } x \text{ BY } x+4} (x+4)^2 \xrightarrow{\text{SUBTRACT 4 FROM THE FUNCTION TO GIVE}}$$

TRANSLATION, BY 4 UNITS TO THE "LEFT"

$$(x+4)^2 - 4$$

TRANSLATION BY 4 UNITS "DOWNWARDS"

∴ TRANSLATION BY THE VECTOR $\begin{pmatrix} -4 \\ -4 \end{pmatrix}$



YGB - SYNOPTIC PAPER F - QUESTION 4

$$x_n = \frac{k - 5x_{n-1}}{x_{n-1}}$$

- $x_1 = 1$
- $x_2 = \frac{k - 5x_1}{x_1} = \frac{k - 5 \times 1}{1} = k - 5$
- $x_3 = \frac{k - 5x_2}{x_2} = \frac{k - 5(k - 5)}{k - 5} = \frac{k - 5k + 25}{k - 5} = \frac{25 - 4k}{k - 5}$

Now we are given THAT $x_3 > 6$ & $k > 5$ so $k - 5 > 0$

$$\Rightarrow x_3 > 6$$

$$\Rightarrow \frac{25 - 4k}{k - 5} > 6$$

$$\Rightarrow 25 - 4k > 6(k - 5)$$

$$\Rightarrow 25 - 4k > 6k - 30$$

$$\Rightarrow -10k > -55$$

$$\Rightarrow k < \frac{55}{10} = \frac{11}{2}$$

$$\therefore \underline{5 < k < \frac{11}{2}}$$

- 1 -

1 YGB - SYNOPTIC PAPER F - QUESTIONS

a) REWRITE THE EQUATION OF THE CIRCLE IN THE "STANDARD FORM"

$$x^2 + y^2 - 10x - 6y + 14 = 0$$

$$x^2 - 10x + y^2 - 6y + 14 = 0$$


$$(x-5)^2 - 25 + (y-3)^2 - 9 + 14 = 0$$

$$(x-5)^2 + (y-3)^2 = 20$$


CENTRE AT (5,3), RADIUS = $\sqrt{20} = 2\sqrt{5}$

LOOKING AT THE DIAGRAM OPPOSITE

$$k = 3 \pm 2\sqrt{5}$$



b) LOOKING AT THE DIAGRAM BELOW & USING PYTHAGORAS' THEOREM



$$\Rightarrow |MB|^2 + |MC|^2 = |CB|^2$$

$$\Rightarrow |MB|^2 + 2^2 = (2\sqrt{5})^2$$

$$\Rightarrow |MB|^2 + 4 = 20$$

$$\Rightarrow |MB|^2 = 16$$

$$\Rightarrow |MB| = 4$$

$$\therefore |AB| = 8$$

c) SOLVING THE TWO EQUATIONS SIMULTANEOUSLY

$$\left. \begin{aligned} x - 2y - 9 &= 0 \\ (x-5)^2 + (y-3)^2 &= 20 \end{aligned} \right\}$$

$$\Rightarrow x = 2y + 9$$

$$\Rightarrow [(2y+9)-5]^2 + (y-3)^2 = 20$$

YGB - SYNOPSIS PAPER F - QUESTIONS

$$\Rightarrow (2y+4)^2 + (y-3)^2 = 20$$

$$\Rightarrow 4y^2 + 16y + 16 + y^2 - 6y + 9 = 20$$

$$\Rightarrow 5y^2 + 10y + 5 = 0$$

$$\Rightarrow y^2 + 2y + 1 = 0$$

$$\Rightarrow (y+1)^2 = 0 \quad (\text{EXPECTED A REPEATED ROOT})$$


$$\Rightarrow y = -1$$

4 USING $x = 2y + 9$
 $x = 7$

\therefore $D(7, -1)$ //

- 1 -

IXGB - SYNOPTIC PAPER F - QUESTION 6



CONSTRAINT ON THE LENGTH OF THE WIRE

$$\Rightarrow P = 60$$

$$\Rightarrow 3x + 2x + x\theta = 60$$

$$\Rightarrow 2\theta + 5x = 60$$

$$\Rightarrow \underline{x\theta = 60 - 5x}$$

// AS REQUIRED

b) TOTAL AREA IS GIVEN BY

$$\frac{1}{2}x^2 \sin \frac{\pi}{3} + \frac{1}{2}x^2\theta \quad \leftarrow \quad \frac{1}{2}r^2\theta$$

$$\Rightarrow A = \frac{1}{2}x^2 \frac{\sqrt{3}}{2} + \frac{1}{2}x(2\theta)$$

$$\Rightarrow A = \frac{1}{4}\sqrt{3}x^2 + \frac{1}{2}x(60 - 5x)$$

$$\Rightarrow A = \frac{1}{4}\sqrt{3}x^2 + 30x - \frac{5}{2}x^2$$

$$\Rightarrow A = \frac{1}{4}\sqrt{3}x^2 - \frac{5}{2}x^2 + 30x$$

$$\Rightarrow \underline{A = \frac{1}{4}(\sqrt{3}-10)x^2 + 30x}$$

// AS REQUIRED

c) DIFFERENTIATE & SOLVE FOR ZERO

$$\Rightarrow \frac{dA}{dx} = \frac{1}{2}(\sqrt{3}-10)x + 30$$

$$\Rightarrow 0 = \frac{1}{2}(\sqrt{3}-10)x + 30$$

$$\Rightarrow 0 = (\sqrt{3}-10)x + 60$$

$$\Rightarrow -60 = (\sqrt{3}-10)x$$

$$\Rightarrow x = \frac{-60}{\sqrt{3}-10}$$

$$\Rightarrow \underline{x = 7.26}$$

//

LYGB - SYNOPTIC PAPER F - QUESTION 6

$$d) \Rightarrow A = \frac{1}{4}(\sqrt{3}-10)x^2 + 30x$$

$$\Rightarrow A_{\text{MAX}} = \frac{1}{4}(\sqrt{3}-10)(7.26\dots)^2 + 30(7.26\dots)$$

$$\Rightarrow \underline{A_{\text{MAX}} = 109 \text{ cm}^2}$$

(3sf)

AND FINALLY JUSTIFYING

$$\Rightarrow \frac{dA}{dx} = \frac{1}{2}(\sqrt{3}-10)x + 30$$

$$\Rightarrow \frac{d^2A}{dx^2} = \frac{1}{2}(\sqrt{3}-10)$$

$$\Rightarrow \left. \frac{d^2A}{dx^2} \right|_{x=7.26} = -4.13\dots < 0$$

INDEED A MAXIMUM

LYGB - SYNOPTIC PAPER F - QUESTION 7

$$x^2 + (k-1)x + (k+2) = 0, k \in \mathbb{R}$$

REPEATED ROOTS $\Rightarrow b^2 - 4ac = 0$
 $\Rightarrow (k-1)^2 - 4 \times 1 \times (k+2) = 0$
 $\Rightarrow (k-1)^2 - 4(k+2) = 0$
 $\Rightarrow k^2 - 2k + 1 - 4k - 8 = 0$
 $\Rightarrow k^2 - 6k - 7 = 0$
 $\Rightarrow (k+1)(k-7) = 0$
 $\Rightarrow k = \begin{matrix} -1 \\ 7 \end{matrix}$

THERE ARE NOW TWO CASES TO CONSIDER

● IF $k = -1$

$$x^2 - 2x + 1 = 0$$

$$(x-1)^2 = 0$$

$$x = 1$$



● IF $k = 7$

$$x^2 + 6x + 9 = 0$$

$$(x+3)^2 = 0$$

$$x = -3$$



NGB - SYNOPTIC PAPER F - QUESTIONS


a) EVN INFLUES SYMMETRY ABOUT THE y AXIS - SKETCH FOR $x \geq 0$

$$f_1(0) = 0^2 - 2 \times 0 = 0$$

$$f_1(3) = 3^2 - 2 \times 3 = 3$$

$$f_2(3) = 6 - 3 = 3$$

HENCE WE HAVE NOTING THAT $x^2 - 2x = x(x-2)$



b) SOLVING $f(x) = \frac{5}{4}$ $0 < x < 3$ or $x > 3$

$$\bullet x^2 - 2x = \frac{5}{4}$$

$$4x^2 - 8x = 5$$

$$4x^2 - 8x - 5 = 0$$

$$(2x + 1)(2x - 5) = 0$$


$$x = \begin{cases} \frac{5}{2} \\ -\frac{1}{2} \end{cases}$$

$$\bullet 6 - x = \frac{5}{4}$$

$$24 - 4x = 5$$

$$19 = 4x$$

$$x = \frac{19}{4}$$



BUT $f(x)$ IS EVN

$$\therefore x = \pm \frac{5}{2}, \pm \frac{19}{4}$$

- 1 -

LYGB - SYNOPTIC PAPER F - QUESTION 9

METHOD A

$$\begin{aligned}
 \ln(2\sqrt{e}) - \frac{1}{3}\ln\left(\frac{8}{e^2}\right) - \ln\left(\frac{1}{3}e\right) &= \frac{1}{3}\left[3\ln(2e^{\frac{1}{2}}) - \ln\left(\frac{8}{e^2}\right) - 3\ln\left(\frac{e}{3}\right)\right] \\
 &= \frac{1}{3}\left[\ln(8e^{\frac{3}{2}}) - \ln\left(\frac{8}{e^2}\right) - \ln\left(\frac{e^3}{27}\right)\right] \\
 &= \frac{1}{3}\left[\ln(8e^{\frac{3}{2}}) + \ln\left(\frac{e^2}{8}\right) + \ln\left(\frac{27}{e^3}\right)\right] \\
 &= \frac{1}{3}\ln\left[\cancel{8}e^{\frac{3}{2}} \times \frac{e^2}{\cancel{8}} \times \frac{27}{e^3}\right] \\
 &= \frac{1}{3}\ln\left[27e^{\frac{1}{2}}\right] \\
 &= \frac{1}{3}\left[\ln 27 + \ln e^{\frac{1}{2}}\right] \\
 &= \frac{1}{3}\left[\ln 3^3 + \frac{1}{2}\ln e\right] \\
 &= \frac{1}{3}\left[3\ln 3 + \frac{1}{2} \times 1\right] \\
 &= \underline{\underline{\ln 3 + \frac{1}{6}}} \quad // \quad a=6 \quad b=3
 \end{aligned}$$

METHOD B

$$\begin{aligned}
 \ln(2\sqrt{e}) - \frac{1}{3}\ln\left(\frac{8}{e^2}\right) - \ln\left(\frac{1}{3}e\right) &= \ln 2 + \ln \sqrt{e} - \frac{1}{3}[\ln 8 - \ln e^2] - \left[\ln \frac{1}{3} + \ln e\right] \\
 &= \ln 2 + \ln e^{\frac{1}{2}} - \frac{1}{3}[\ln 2^3 - 2\ln e] - \left[\ln \frac{1}{3} + 1\right] \\
 &= \ln 2 + \frac{1}{2}\ln e - \frac{1}{3}[3\ln 2 - 2 \times 1] - \left[\ln \frac{1}{3} + 1\right] \\
 &= \ln 2 + \frac{1}{2} \times 1 - \frac{1}{3}[3\ln 2 - 2] - \left[\ln \frac{1}{3} + 1\right] \\
 &= \cancel{\ln 2} + \frac{1}{2} - \cancel{\ln 2} + \frac{2}{3} - \ln \frac{1}{3} - 1 \\
 &= \frac{1}{6} - \ln \frac{1}{3} \\
 &= \underline{\underline{\frac{1}{6} + \ln 3}} \quad // \quad \text{As before}
 \end{aligned}$$

YGB - SYNOPTIC PAPER F - QUESTION 10

START WITH THE STATIONARY POINTS

$$f(x) = 3x^4 - 8x^3 - 6x^2 + 24x - 8$$

$$f'(x) = 12x^3 - 24x^2 - 12x + 24$$

SOLVING FOR ZERO

$$\Rightarrow 12x^3 - 24x^2 - 12x + 24 = 0$$

$$\Rightarrow x^3 - 2x^2 - x + 2 = 0$$

$$\Rightarrow x^2(x-2) - (x-2) = 0$$

$$\Rightarrow (x-2)(x^2-1) = 0$$

$$\Rightarrow (x-2)(x-1)(x+1) = 0$$

$$x = \begin{cases} 2 \\ 1 \\ -1 \end{cases} \quad y = \begin{cases} 0 \\ 5 \\ -27 \end{cases} \quad \therefore \begin{matrix} (2, 0) \\ (1, 5) \\ (-1, -27) \end{matrix}$$

NOW AS THE FUNCTION IS STATIONARY ON THE x AXIS IT MUST HAVE A REPEATED ROOT AT x=2 (NO INFLEXION AS THERE ARE TWO MORE STATIONARY VALUES).

HENCE DIVIDE BY (x-2)²

$$x^2 - 4x + 4 \begin{array}{r} 3x^2 + 4x - 2 \\ \hline 3x^4 - 8x^3 - 6x^2 + 24x - 8 \\ -3x^4 + 12x^3 - 12x^2 \\ \hline 4x^3 - 18x^2 + 24x - 8 \\ -4x^3 + 16x^2 - 16x \\ \hline -2x^2 + 8x - 8 \\ 2x^2 - 8x + 8 \\ \hline 0 \end{array}$$

$$\therefore f(x) = (x-2)^2 (3x^2 + 4x - 2)$$

$$b^2 - 4ac = 16 - 4(3)(-2) = 40$$


-2-

IYGB - SYNOPTIC PAPER F - QUESTION 10

GETTING THE "ZEROS" OF THE QUADRATIC

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-4 \pm \sqrt{40}}{2 \times 3} = \frac{-4 \pm 2\sqrt{10}}{6} = -\frac{2}{3} \pm \frac{1}{3}\sqrt{10}$$

HENCE WE OBTAIN A FINAL SKETCH OF THE QUARTIC FUNCTION



AND FINALLY THE RANGE LOOKING AT THE ABOVE GRAPH IS

$-27 \leq f(x) \leq 37$

NYGB-SYNOPTIC PAPER F-QUESTION 11

a) EASIER TO REWRITE & USE TRANSFORMATIONS


$$f(x) = \frac{3x+3}{x-2} = \frac{3(x-2)+9}{x-2} = \frac{3(x-2)}{x-2} + \frac{9}{x-2} = 3 + \frac{9}{x-2}$$

$$\frac{1}{x} \mapsto 9\left(\frac{1}{x}\right) \mapsto \frac{9}{x-2} \mapsto \frac{9}{x-2} + 3$$


VERTICAL STRETCH
By s.f 9

TRANSLATION,
2 UNITS, "RIGHT"

TRANSLATION,
3 UNIT "UPWARDS"



HENCE THE SKETCH CAN NOW BE OBTAINED



$x=0, y=-\frac{3}{2} \quad (0, -\frac{3}{2})$
 $y=0, x=-1 \quad (-1, 0)$

b) SOLVING $f(x) = 2$

$$\frac{3x+3}{x-2} = 2$$

$$3x+3 = 2x-4$$

$$x = -7$$

MGB-SYNOPTIC PAPER F-QUESTION 11

OR USING THE ALTERNATIVE FORM FOUND


$$\Rightarrow 3 + \frac{9}{x-2} = 2$$

$$\Rightarrow \frac{9}{x-2} = -1$$

$$\Rightarrow 9 = -x + 2$$

$$\Rightarrow \underline{x = -7}$$

c) LOOKING AT THE SKETCH BELOW



$$\underline{x \leq -7 \text{ OR } x > 2}$$

LYGB - SYNOPTIC PAPER F - QUESTION 12

a) OBTAIN THE FIRST DERIVATIVE

$$y = 2 - 2\ln(x^2 + 4)$$

$$\frac{dy}{dx} = 1 - 2 \times \frac{1}{x^2 + 4} \times 2x$$

$$\frac{dy}{dx} = 1 - \frac{4x}{x^2 + 4}$$

DIFFERENTIATE AGAIN BY THE QUOTIENT RULE

$$\frac{d^2y}{dx^2} = 0 - \frac{(x^2 + 4) \times 4 - 4x(2x)}{(x^2 + 4)^2}$$

$$\frac{d^2y}{dx^2} = - \frac{4x^2 + 16 - 8x^2}{(x^2 + 4)^2}$$

$$\frac{d^2y}{dx^2} = - \frac{16 - 4x^2}{(x^2 + 4)^2}$$

$$\frac{d^2y}{dx^2} = \frac{4x^2 - 16}{(x^2 + 4)^2} = \frac{4(x^2 - 4)}{(x^2 + 4)^2} \quad \text{As required}$$

b) Solving $\frac{dy}{dx} = 0$

$$1 - \frac{4x}{x^2 + 4} = 0$$

$$x^2 + 4 - 4x = 0$$

$$x^2 - 4x + 4 = 0$$

$$(x - 2)^2 = 0$$

$$x = 2$$

$$y = 2 - 2\ln(2^2 + 4)$$

$$y = 2 - 2\ln 8$$

$$y = 2 - 6\ln 2$$

$$\left. \frac{d^2y}{dx^2} \right|_{x=2} = \frac{4(2^2 - 4)}{8^2} = 0$$

$$\therefore (2, 2 - 6\ln 2) \text{ is}$$

$$\text{A POINT OF INFLEXION}$$

YGB - SYNOPTIC PAPER F - QUESTION 13

a) SETTING UP TWO EQUATIONS

● $S_n = \frac{n}{2} [a + L]$

● $u_n = a + (n-1)d$

$S_{25} = 1050$

$u_{25} = 72$

$\frac{25}{2}(a+L) = 1050$

$a + 24d = 72$

$25(a+72) = 2100$

$12 + 24d = 72$

$a + 72 = \frac{2100}{25}$

$24d = 60$

$a + 72 = \frac{4200}{50}$

$d = \frac{250}{24}$

$a + 72 = \frac{8400}{100}$

$d = \frac{5}{2}$

$a + 72 = 84$

$a = 12$

b) LET US NOTE THAT $\sum_{h=1}^{25} u_h = 1050$ (GIVEN IN PART a)

$\Rightarrow 117 [1050 - T_k] = 233 T_k$

$T_k = \sum_{h=1}^k u_h$

$\Rightarrow 117 \times 1050 - 117 T_k = 233 T_k$

$\Rightarrow 117 \times 1050 = 350 T_k$

$\Rightarrow T_k = \frac{117 \times 1050}{350}$

$\Rightarrow T_k = 351$

$\Rightarrow \sum_{h=1}^k u_h = 351$

1YGB - SYNOPTIC PAPER F - QUESTION 13

USING $a=12$ & $d=\frac{5}{2}$

$$\Rightarrow \frac{k}{2} \left[2 \times 12 + (k-1) \times \frac{5}{2} \right] = 351$$

$$\Rightarrow \frac{k}{2} \left[24 + \frac{5}{2}(k-1) \right] = 351$$

$$\Rightarrow k \left[12 + \frac{5}{4}(k-1) \right] = 351$$

$$\Rightarrow 4k \left[12 + \frac{5}{4}(k-1) \right] = 1404 \quad \left. \begin{array}{l} \text{red arrow} \\ \times 4 \end{array} \right\}$$

$$\Rightarrow k \left[48 + 5(k-1) \right] = 1404$$

$$\Rightarrow k \left[5k + 43 \right] = 1404$$

NOW BY TRIAL & IMPROVEMENT, NOTING THAT $k < 24$

$$\text{IF } k=10 \Rightarrow 10 \times 93 < 1404$$

$$\text{IF } k=15 \Rightarrow 15 \times 118 > 1404$$

$$\text{IF } k=13 \Rightarrow 13 \times 108 = 1404$$

so $k=13$

ALTERNATIVE

$$\frac{k}{2} \left[2 \times 12 + (k-1) \times \frac{5}{2} \right] = 351$$

$$\frac{k}{2} \left[24 + \frac{5}{2}k - \frac{5}{2} \right] = 351$$

$$12k + \frac{5}{4}k^2 - \frac{5}{4}k = 351$$

$$48k + 5k^2 - 5k = 1404$$

$$5k^2 + 43k - 1404 = 0$$

$$k = \frac{-43 \pm \sqrt{43^2 - 4 \times 5 \times (-1404)}}{2 \times 5} = \begin{cases} \frac{-43 + 173}{10} = 13 \\ \frac{-43 - 173}{10} = -216 \end{cases}$$

1 YGB - SYNOPTIC PAPER F - QUESTION 14

$$\sqrt{2}x + \sqrt{3}y = 5$$

$$(5\sqrt{3} - \sqrt{2})x + (5\sqrt{2} - \sqrt{3})y = 10\sqrt{6}$$

START WITH THE SECOND EQUATION

$$\Rightarrow (5\sqrt{3} - \sqrt{2})x + (5\sqrt{2} - \sqrt{3})y = 10\sqrt{6}$$

$$\Rightarrow 5\sqrt{3}x - \sqrt{2}x + 5\sqrt{2}y - \sqrt{3}y = 10\sqrt{6}$$

$$\Rightarrow 5\sqrt{3}x + 5\sqrt{2}y = 10\sqrt{6} + \sqrt{2}x + \sqrt{3}y$$

$$\Rightarrow 5\sqrt{3}x + 5\sqrt{2}y = 10\sqrt{6} + 5$$

$$\Rightarrow \sqrt{3}x + \sqrt{2}y = 2\sqrt{6} + 1$$

NEXT ELIMINATE AS FOLLOWS

$$\begin{array}{l} \sqrt{2}x + \sqrt{3}y = 5 \\ \sqrt{3}x + \sqrt{2}y = 2\sqrt{6} + 1 \end{array} \begin{array}{l} \times \sqrt{3} \\ \times \sqrt{2} \end{array} \Rightarrow \begin{array}{l} \sqrt{6}x + 3y = 5\sqrt{3} \\ \sqrt{6}x + 2y = 2\sqrt{12} + \sqrt{2} \end{array}$$

SUBTRACTING

$$\Rightarrow y = 5\sqrt{3} - (2\sqrt{12} + \sqrt{2})$$

$$\Rightarrow y = 5\sqrt{3} - 4\sqrt{3} - \sqrt{2}$$

$$\Rightarrow y = \sqrt{3} - \sqrt{2}$$

AND TO FIND x

$$\sqrt{2}x + \sqrt{3}y = 5$$

$$\sqrt{2}x + \sqrt{3}(\sqrt{3} - \sqrt{2}) = 5$$

$$\sqrt{2}x + 3 - \sqrt{6} = 5$$

$$\sqrt{2}x = 2 + \sqrt{6} \quad \left. \begin{array}{l} \\ \\ \end{array} \right) \times \sqrt{2}$$


$$2x = 2\sqrt{2} + \sqrt{12}$$

$$2x = 2\sqrt{2} + 2\sqrt{3}$$

$$\therefore x = \sqrt{3} + \sqrt{2}$$

1YGB - SYNOPTIC PAPER F - QUESTION 15

LOOKING AT THE DIAGRAM BELOW



DETERMINE THE DISTANCE

$$|AB| = \sqrt{(y_2 - y_1)^2 + (x_2 - x_1)^2} = \sqrt{(2 - 3)^2 + (8 - 1)^2} = \sqrt{10}$$

LET C(x, y)

$$|AC| = \sqrt{(8 - x)^2 + (2 - y)^2}$$

NOW |AC| = 2|AB|

$$\Rightarrow \sqrt{(8 - x)^2 + (2 - y)^2} = 2\sqrt{10}$$

$$\Rightarrow (8 - x)^2 + (2 - y)^2 = 40$$

$$\Rightarrow (x - 8)^2 + (y - 2)^2 = 40$$

$$\left. \begin{aligned} (8 - x)^2 &= 64 - 16x + x^2 \\ (x - 8)^2 &= x^2 - 16x + 64 \end{aligned} \right\}$$

BUT THE POINT C LIES ON $x + y = 14$

$$\Rightarrow [(14 - y) - 8]^2 + (y - 2)^2 = 40$$

$$\Rightarrow (6 - y)^2 + (y - 2)^2 = 40$$

$$\Rightarrow \begin{pmatrix} 36 - 12y + y^2 \\ 4 - 4y + y^2 \end{pmatrix} = 40$$

$$\Rightarrow 2y^2 - 16y + 40 = 40$$

1YGB - SYNOPTIC PAPER F - QUESTION 15

$$\Rightarrow 2y^2 - 16y = 0$$

$$\Rightarrow 2y(y - 8)$$

$$\Rightarrow y = \begin{cases} 0 \\ 8 \end{cases}$$

or using $x = 14 - y$

$$\Rightarrow x = \begin{cases} 14 \\ 6 \end{cases}$$

\therefore $C(14,0)$ or $C(6,8)$ //

- 1 -

YGB - SYNOPTIC PAPER F - QUESTION 16

a) STARTING FROM THE R.H.S

$$\begin{aligned} \text{R.H.S.} &= \frac{2 \tan x}{1 + \tan^2 x} = \frac{2 \tan x}{\sec^2 x} = 2 \tan x \times \cos^2 x \\ &= 2 \times \frac{\sin x}{\cancel{\cos x}} \times \cos^2 x = 2 \sin x \cos x = 2 \sin 2x \\ &= \text{L.H.S.} \end{aligned}$$

As required

b) THE PARTIAL FRACTIONS NEXT

$$\frac{8}{(3t+1)(t+3)} \equiv \frac{A}{3t+1} + \frac{B}{t+3}$$

$$\boxed{8 \equiv A(t+3) + B(3t+1)}$$

• If $t = -3$

$$8 = -8B$$

$$\underline{\underline{B = -1}}$$

• If $t = -\frac{1}{3}$

$$8 = \frac{8}{3}A$$

$$\underline{\underline{A = 3}}$$

$$\therefore \frac{8}{(3t+1)(t+3)} = \frac{3}{3t+1} - \frac{1}{t+3}$$

c) USING THE SUBSTITUTION GIVEN & THE PREVIOUS PARTS

$$\Rightarrow t = \tan x$$

$$\Rightarrow \frac{dt}{dx} = \sec^2 x$$

1YGB - SYNOPTIC PAPER F - QUESTION 16

$$\Rightarrow dx = \frac{dt}{\sec^2 x}$$

$$\Rightarrow dx = \frac{dt}{\tan^2 x + 1}$$

$$\Rightarrow \boxed{dx = \frac{dt}{1+t^2}}$$

- when $x=0$, $t=0$
- when $x=\frac{\pi}{4}$, $t=1$

TRANSFORMING THE INTEGRAL

$$\int_0^{\frac{\pi}{4}} \frac{8}{3+5\sin 2x} dx = \int_0^{\frac{\pi}{4}} \frac{8}{3+5\left(\frac{2\tan x}{1+\tan^2 x}\right)} dx \quad \leftarrow \text{BY PART (a)}$$

$$= \int_0^1 \frac{8}{3+5\left(\frac{2t}{1+t^2}\right)} \left(\frac{dt}{1+t^2}\right) \quad \leftarrow \text{BY THE SUBSTITUTION}$$

$$= \int_0^1 \frac{8}{3(1+t^2) + 10t} dt = \int_0^1 \frac{8}{3t^2 + 10t + 3} dt$$

$$= \int_0^1 \frac{8}{(3t+1)(t+3)} dt = \int_0^1 \left(\frac{3}{3t+1} - \frac{1}{t+3}\right) dt \quad \leftarrow \text{BY (b)}$$

$$= \left[\ln|3t+1| - \ln|t+3| \right]_0^1 = \cancel{(\ln 4 - \ln 4)} - \cancel{(\ln 1 - \ln 3)}$$

$$= \ln 3$$

AS REQUIRED

- -

YGB - SYNOPTIC PAPER F - QUESTION 17

a) FIND THE GRADIENT FUNCTION IN TERMS OF t

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{3at^2}{3a} = t^2$$

SO NORMAL GRADIENT AT A GENERAL POINT WILL BE $\frac{-1}{t^2}$

EQUATION OF NORMAL AT (3at, at^3)

$$\begin{aligned} \Rightarrow y - y_0 &= m(x - x_0) \\ \Rightarrow y - at^3 &= -\frac{1}{t^2}(x - 3at) \\ \Rightarrow ty - at^5 &= -x + 3at \\ \Rightarrow \underline{yt^2 + x} &= \underline{at^5 + 3at} \end{aligned}$$

// AS REQUIRED

b) USING THE TWO POINTS GIVEN INTO THE GENERAL NORMAL

$$\begin{aligned} (7, 3) &\Rightarrow 3t^2 + 7 = 3at + at^5 \\ (-1, 5) &\Rightarrow 5t^2 - 1 = 3at + at^5 \end{aligned} \left. \vphantom{\begin{aligned} (7, 3) \\ (-1, 5) \end{aligned}} \right\} \begin{aligned} \Rightarrow 5t^2 - 1 &= 3t^2 + 7 \\ \Rightarrow 2t^2 &= 8 \\ \Rightarrow t^2 &= 4 \\ \Rightarrow t &= \begin{matrix} -2 \\ 2 \end{matrix} \end{aligned}$$

Now if t = 2

$$\begin{aligned} 3 \times 2^2 + 7 &= 6a + 32a \\ 19 &= 38a \\ a &= \frac{1}{2} \end{aligned}$$

AND IF t = -2

$$\begin{aligned} 3(-2)^2 + 7 &= -6a - 32a \\ 19 &= -38a \\ a &= \cancel{-\frac{1}{2}} \quad a > 0 \end{aligned}$$

∴ a = 1/2, t = 2 YIELDS

P(3at, at^3) = P(3, 4)

//

1YCB - SYNOPTIC PAPER F - QUESTION 17

ALTERNATIVE BY FINDING THE EQUATION OF THE NORMAL

$$(7,3) \text{ \& } (-1,5) \implies m = \frac{5-3}{-1-7} = \frac{2}{-8} = -\frac{1}{4}$$

NORMAL GRADIENT = $-\frac{1}{t^2} = -\frac{1}{4}$

$$\therefore t^2 = 4$$
$$t = \begin{cases} 2 \\ -2 \end{cases}$$

• IF t=2

$$4y + x = 6a + 32a$$

$$4y + x = 38a$$

$$(7,3): 12 + 7 = 38a$$

$$19 = 38a$$

$$a = \frac{1}{2}$$

OR $(-1,5): 20 - 1 = 38a$

• IF t=-2

$$4y + x = -6a - 32a$$

$$4y + x = -38a$$

$$(7,3): 12 + 7 = -38a$$

$$a = \frac{1}{2} \quad a > 0$$

As before $a = \frac{1}{2}, t = 2$ yields $P(3,4)$

- 1 -

1YGB - SYNOPTIC PAPER F - QUESTION 18

USING THE IDENTITY FOR $\tan(A-B)$

$$\tan(A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

$$\Rightarrow \tan(\overset{A}{\arctan 3x} - \overset{B}{\arctan 2}) + \tan(\overset{A}{\arctan 3} - \overset{B}{\arctan 2x}) = \frac{3}{8}$$

$$\Rightarrow \frac{3x - 2}{1 + (3x)(2)} + \frac{3 - 2x}{1 + 3(2x)} = \frac{3}{8}$$

$$\Rightarrow \frac{3x - 2}{1 + 6x} + \frac{3 - 2x}{1 + 6x} = \frac{3}{8}$$

$$\Rightarrow \frac{x+1}{1+6x} = \frac{3}{8}$$


$$\Rightarrow 8x + 8 = 3 + 18x$$

$$\Rightarrow 5 = 10x$$


$$\Rightarrow x = \frac{1}{2}$$

1YGB - SYNOPSIS PAPER F - QUESTION 19

START WITH A DIAGRAM ; A(3,1), B(7,4), C(10,-4)



$\tan \theta = \frac{4}{3}$ $\tan \phi = \frac{3}{8}$



HENCE BY USING THE COMPOUND ANGLE IDENTITY FOR $\tan(\theta + \phi)$

$\Rightarrow \tan(\theta + \phi) = \frac{\tan \theta + \tan \phi}{1 - \tan \theta \tan \phi}$

$= \frac{\frac{4}{3} + \frac{3}{8}}{1 - \frac{4}{3} \times \frac{3}{8}}$


$= \frac{\frac{4}{3} + \frac{3}{8}}{\frac{1}{2}}$

$= 2 \left(\frac{4}{3} + \frac{3}{8} \right)$

$= 2 \left(\frac{32 + 9}{24} \right)$

$= 2 \times \frac{41}{24}$

$= \frac{41}{12}$



-1-

NGB - SYNOPSIS PAPER F - QUESTION 20

a) LOOKING AT THE DIAGRAM & BY RELATING VARIABLES

$$\Rightarrow \frac{dv}{dt} = +k(2r-y) \quad k > 0$$


$$\Rightarrow \frac{dv}{dy} \times \frac{dy}{dt} = k(2r-y)$$

$$\Rightarrow \pi(2ry - y^2) \frac{dy}{dt} = k(2r-y)$$

$$\Rightarrow \pi y(2r-y) \frac{dy}{dt} = k(2r-y)$$

$$\Rightarrow \pi y \frac{dy}{dt} = k \quad (2r-y \neq 0)$$

$$\Rightarrow \frac{dy}{dt} = \frac{k}{\pi y} \quad \text{AS REQUIRED}$$



ii) SOLVING THE O.D.E BY SEPARATION OF VARIABLES

$$\Rightarrow y \, dy = \frac{k}{\pi} \, dt$$

$$\Rightarrow \int_{y=0}^r y \, dy = \int_{t=0}^t \frac{k}{\pi} \, dt$$

$$\Rightarrow \left[\frac{1}{2}y^2 \right]_{y=0}^{y=r} = \left[\frac{k}{\pi} t \right]_0^t$$

$$\Rightarrow \frac{1}{2}r^2 - 0 = \frac{k}{\pi} t - 0$$

$$\Rightarrow \frac{r^2}{2} = \frac{kt}{\pi}$$

$$\Rightarrow t = \frac{\pi r^2}{2k} \quad \text{AS REQUIRED}$$

MYGB - SYNOPSIS PAPER F - QUESTION 20

b) REMODEL THE O.D.E

WATER GOING IN AT CONSTANT RATE

$$\frac{dv}{dt} = k(2r - y) = \underbrace{2kr}_{\text{WATER GOING IN AT CONSTANT RATE}} - \underbrace{ky}_{\text{WATER LEAKING OUT PROPORTIONAL TO } y}$$

HENCE THE REQUIRED O.D.E FOR LEAVING ONLY IS

$$\frac{dv}{dt} = -ky, \quad k > 0 \quad \text{SUBJECT TO } t=0 \quad y=r$$

BY RELATING VARIABLES AS BEFORE

$$\Rightarrow \frac{dv}{dy} \frac{dy}{dt} = -ky$$

$$\Rightarrow \pi y(2r - y) \frac{dy}{dt} = -ky$$

$$\Rightarrow (2r - y) dy = -\frac{k}{\pi} dt$$

INTEGRATING SUBJECT TO $t=0, y=r$ & REQUIRING t AT $y=0$

$$\Rightarrow \int_{y=r}^{y=0} (2r - y) dy = \int_{t=0}^t -\frac{k}{\pi} dt$$

$$\Rightarrow \left[2ry - \frac{1}{2}y^2 \right]_{y=r}^{y=0} = \left[-\frac{k}{\pi} t \right]_0^t$$

$$\Rightarrow 0 - \left(2r^2 - \frac{1}{2}r^2 \right) = -\frac{k}{\pi} t - 0$$

$$\Rightarrow 2r^2 - \frac{1}{2}r^2 = \frac{k}{\pi} t$$

$$\Rightarrow \frac{3}{2}r^2 = \frac{k}{\pi} t$$

$$\Rightarrow t = \frac{3\pi r^2}{2k}$$

$$\Rightarrow t = 3 \left(\frac{\pi r^2}{2k} \right)$$

IT TAKES $\frac{\pi r^2}{2k}$ TO FILL UP

IT TAKES $3 \left(\frac{\pi r^2}{2k} \right)$ TO EMPTY

IF 3 TIMES AS LONG