

YGB - SYNOPSIS PAPER L - QUESTION 1

a) EXPANDING BINOMIALLY

$$(1+x)^{-1} = 1 + \frac{-1}{1}(x)^1 + \frac{-1(-2)}{1 \times 2}(x)^2 + \frac{-1(-2)(-3)}{1 \times 2 \times 3}(x)^3 + o(x^4)$$

$$= \underline{1 - x + x^2 - x^3 + o(x^3)}$$

b) LET $g(x) = (1+x)^{-1}$

THEN $(1-2x)^{-1} = g(-2x)$

Hence $g(-2x) = (1-2x)^{-1} = 1 - (-2x) + (-2x)^2 - (-2x)^3 + o(x^4)$

$$= \underline{1 + 2x + 4x^2 + 8x^3 + o(x^4)}$$

c) BY PARTIAL FRACTIONS OR DIRECT MULTIPLICATION

$$f(x) = (4x+1)(1+x)^{-1}(1-2x)^{-1}$$

$$f(x) = (1+4x)(1-x+x^2-x^3+\dots)(1+2x+4x^2+8x^3+\dots)$$

$$f(x) = (1+4x) \begin{bmatrix} 1 + 2x + 4x^2 + 8x^3 + o(x^4) \\ -x - 2x^2 - 4x^3 + o(x^4) \\ x^2 + 2x^3 + o(x^4) \\ -x^3 + o(x^4) \end{bmatrix}$$

$$f(x) = (1+4x)(1 + 2 + 3x^2 + 5x^3 + o(x^4))$$

$$f(x) = \underline{1 + 2 + 3x^2 + 5x^3 + o(x^4) + 4x + 4x^2 + 12x^3 + o(x^4)}$$

\therefore $f(x) = 1 + 5x + 7x^2 + 17x^3 + o(x^4)$

- $(1+x)^{-1}$ is valid $|x| < 1$, i.e. $-1 < x < 1$
 - $(1-2x)^{-1}$ is valid $|2x| < 1$, i.e. $-\frac{1}{2} < x < \frac{1}{2}$
- \therefore Hence $-\frac{1}{2} < x < \frac{1}{2}$

YGB - SYNOPTIC PAPER L - QUESTION 2

a) WHAT t BECOMES VERY LARGE, IF $t \rightarrow \infty$

$$e^{-0.1t} \rightarrow 0$$

$$200e^{-0.1t} \rightarrow 0$$

$\therefore \theta \rightarrow 225$

\therefore MAX TEMPERATURE IS 225°C

b) WHAT $\theta = 125$

$$\Rightarrow 125 = 225 - 200e^{-0.1t}$$

$$\Rightarrow 200e^{-0.1t} = 100$$

$$\Rightarrow e^{-0.1t} = \frac{1}{2}$$

$$\Rightarrow e^{0.1t} = 2$$

$$\Rightarrow 0.1t = \ln 2$$

$$\Rightarrow t = 10 \ln 2 \approx 6.93147 \dots \approx \underline{7 \text{ min}}$$

c) WORK AS FOLLOWS

$$\theta = 225 - 200e^{-0.1t}$$

$$\frac{d\theta}{dt} = 0 + 20e^{-0.1t}$$

$$\frac{d\theta}{dt} = 20e^{-0.1t}$$

$$10 \frac{d\theta}{dt} = 200e^{-0.1t}$$

$$10 \frac{d\theta}{dt} = 225 - \theta$$

$$\frac{d\theta}{dt} = \frac{1}{10}(225 - \theta)$$

~~AS REQUIRED~~

$$\left. \frac{d\theta}{dt} \right|_{\theta=125} = \underline{10^\circ/\text{min}}$$

FROM THE ORIGINAL EQUATION

$$\theta = 225 - 200e^{-0.1t}$$

$$200e^{-0.1t} = 225 - \theta$$

1YGB - SYNOPTIC PAPER L - QUESTION 3

a) USING THE STANDARD EQUATION OF A CIRCLE

$$\Rightarrow (x-a)^2 + (y-b)^2 = r^2$$

$$\Rightarrow (x+3)^2 + (y-8)^2 = \left(\frac{1}{2}\sqrt{20}\right)^2$$

$$\Rightarrow \underline{(x+3)^2 + (y-8)^2 = 20}$$

b) SOLVING SIMULTANEOUSLY WITH $y = 3x + 7$

$$\Rightarrow (x+3)^2 + ((3x+7)-8)^2 = 20$$

$$\Rightarrow (x+3)^2 + (3x-1)^2 = 20$$

$$\Rightarrow x^2 + \cancel{6x} + 9 + 9x^2 - \cancel{6x} + 1 = 20$$

$$\Rightarrow 10x^2 = 10$$

$$\Rightarrow x^2 = 1$$

$$\Rightarrow x = \begin{cases} 1 \\ -1 \end{cases} \quad y = \begin{cases} 3(1)+7 = 10 \\ 3(-1)+7 = 4 \end{cases}$$


$\therefore \underline{A(1,10) \text{ \& } B(-1,4)}$

c) USING GRADIENTS

$$m_{AC} = \frac{8-10}{-3-1} = \frac{-2}{-4} = \frac{1}{2}$$

$$m_{BC} = \frac{8-4}{-3-(-1)} = \frac{4}{-2} = -2$$

NEGATIVE RECIPROALS $\Rightarrow \hat{ACB} = 90^\circ$



$$\text{Area} = \frac{1}{2} |AC| |BC|$$

$$= \frac{1}{2} \sqrt{20} \sqrt{20}$$

$$= \underline{10}$$

- 1 -


1YGB - SYNOPTIC PAPER 2 - QUESTION 4

LOOKING AT THE DIAGRAM

$$\sin \theta = \frac{6}{12} \quad (\text{TRIANGLE ADE})$$

$$\sin \theta = \frac{1}{2}$$

$$\theta = 30^\circ$$



AREA OF RECTANGLE ABCD

$$6 \times 12 = 72$$

AREA OF TRIANGLE ADE

$$= \frac{1}{2} |AD| |AE| \sin \phi$$

$$= \frac{1}{2} \times 6 \times 12 \times \sin(90 - 30)$$

$$= 36 \sin 60^\circ$$

$$= 18\sqrt{3}$$

AREA OF CIRCULAR SECTOR ABE

$$= \pi r^2 \times \frac{\theta}{360}$$

$$= \pi \times 12^2 \times \frac{30}{360}$$

$$= 12\pi$$

FINALLY WE OBTAIN THE AREA

$$\text{REQUIRED AREA} = 72 - 18\sqrt{3} - 12\pi$$

$$\approx \underline{3.12}$$

1YGB - SYNOPTIC PAPER L - QUESTION 5

a) SOLVING SIMULTANEOUSLY TO FIND "A"

$$\left. \begin{aligned} x^3 + xy + y^3 &= 10 \\ y &= x+2 \end{aligned} \right\} \Rightarrow x^3 + x(x+2) + (x+2)^3 = 10$$

$$\Rightarrow x^3 + x^2 + 2x - 10 + (x+2)^3 = 0$$

↑
NO NEED TO EXPAND

LET $f(x) = x^3 + x^2 + 2x - 10 + (x+2)^3$

$f(0.1) = -0.528 < 0$
 $f(0.2) = 1.096 > 0$

AS $f(x)$ IS CONTINUOUS AND CHANGES SIGN IN $(0.1, 0.2)$ THERE IS AT LEAST ONE ROOT IN THE INTERVAL

b) PREPARE THE "N-R ITEMS"

- $f(0.1) = -0.528$
- $f'(x) = 3x^2 + 2x + 2 + 3(x+2)^2$
- $f'(0.1) = 15.46$

HENCE WE HAVE

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

$$x = 0.1 - \frac{-0.528}{15.46}$$

$$x = 0.1341526 \dots$$

∴ APPROXIMATELY 0.134

YGB - SYNOPTIC PAPER L - QUESTION 6


a) WITHOUT THE ORDER BEING IMPORTANT

$$\ln(x) \xrightarrow{\quad} \ln(x+e) \xrightarrow{\quad} 2\ln(x+3)$$

TRANSLATION BY
 $\begin{pmatrix} -e \\ 0 \end{pmatrix}$

VERTICAL STRETCH
BY SCALE FACTOR 2

b) SKETCHING $g(x)$ AND $|g(x)|$



c) SOLVING $|g(x)| = 2$


$$|2\ln(x+e)| = 2 \implies 2\ln(x+e) = 2 \quad \text{OR} \quad 2\ln(x+e) = -2$$

$$\ln(x+e) = 1 \qquad \qquad \qquad \ln(x+e) = -1$$

$$x+e = e \qquad \qquad \qquad x+e = e^{-1}$$

$$x = 0 \qquad \qquad \qquad x = \frac{1}{e} - e$$

d) LOOKING AT THE GRAPH



1YGB - SYNOPTIC PAPER L - QUESTION 7

START BY FINDING THE EQUATION OF THE TANGENT.

• When $x=4$, $y = 4^2 - 6 \times 4 + 10$
 $y = 2$

$\therefore A(4, 2)$

• $y = x^2 - 6x + 10$


$\frac{dy}{dx} = 2x - 6$

$\left. \frac{dy}{dx} \right|_{x=4} = 2 \quad \leftarrow \text{TANGENT GRADIENT}$

• $y - y_0 = m(x - x_0)$

$y - 2 = 2(x - 4)$

$y = 2x - 6$




FIND THE x & y INTERCEPT OF THE TANGENT.

$x=0 \quad y = -6 \quad (0, -6)$

$y=0 \quad x = 3 \quad (3, 0)$

LOOKING AT THE DIAGRAM BELOW



$\int_0^4 x^2 - 6x + 10 \, dx = \left[\frac{1}{3}x^3 - 3x^2 + 10x \right]_0^4$

1YGB - SYNOPTIC PAPER 1 - QUESTION 7

$$= \left(\frac{64}{3} - 48 + 40 \right) - (0)$$

$$= \frac{64}{3}$$

∴ THE REQUIRED AREA IS $\frac{64}{3}$



1708 - SYNOPTIC PAPER L - QUESTION 8

a) DIFFERENTIATE WITH RESPECT TO x

$$\Rightarrow \frac{d}{dx}(4xy) - \frac{d}{dx}((x+2)^2) = \frac{d}{dx}(y^2) - \frac{d}{dx}(5)$$

↑
PRODUCT RULE
↑
CHAIN RULE

$$\Rightarrow 4y + 4x \frac{dy}{dx} - 2(x+2) = 2y \frac{dy}{dx} - 0$$

$$\Rightarrow (4x - 2y) \frac{dy}{dx} = 2(x+2) - 4y$$

$$\Rightarrow \frac{dy}{dx} = \frac{2x - 4y + 4}{4x - 2y}$$

$$\Rightarrow \frac{dy}{dx} = \frac{x - 2y + 2}{2x - y}$$

b) "STATIONARY" $\Rightarrow \frac{dy}{dx} = 0$

$$\therefore x - 2y + 2 = 0$$

$$x = 2y - 2$$

SUBSTITUTE INTO THE EQUATION OF THE CURVE

$$\Rightarrow 4y(2y-2) - (2y-2+2)^2 = y^2 - 5$$

$$\Rightarrow 8y^2 - 8y - 4y^2 = y^2 - 5$$

$$\Rightarrow 3y^2 - 8y + 5 = 0$$

$$\Rightarrow (3y - 5)(y - 1) = 0$$

$$y = \begin{cases} \frac{5}{3} \\ 1 \end{cases} \quad x = \begin{cases} 2 \times \frac{5}{3} - 2 = \frac{4}{3} \\ 2(1) - 2 = 0 \end{cases}$$

$$\therefore \left(\frac{4}{3}, \frac{5}{3} \right) \text{ \& } (0, 1)$$

1YGB - SYNOPSIS PAPER L - QUESTION 9

SWITCHING INTO INDICES FOR "COMFORT"

$$\begin{aligned} \frac{\sqrt[3]{16} - \sqrt[3]{2}}{\sqrt[3]{4}} &= \frac{16^{\frac{1}{3}} - 2^{\frac{1}{3}}}{4^{\frac{1}{3}}} = \frac{(2^4)^{\frac{1}{3}} - 2^{\frac{1}{3}}}{(2^2)^{\frac{1}{3}}} = \frac{2^{\frac{4}{3}} - 2^{\frac{1}{3}}}{2^{\frac{2}{3}}} \\ &= \frac{2 \times 2^{\frac{1}{3}} - 2^{\frac{1}{3}}}{2^{\frac{2}{3}}} = \frac{2^{\frac{1}{3}}}{2^{\frac{2}{3}}} \end{aligned}$$

NOW RATIONALIZING

$$\begin{aligned} &= \frac{2^{\frac{1}{3}} \times 2^{\frac{2}{3}}}{2^{\frac{2}{3}} \times 2^{\frac{1}{3}}} = \frac{2^{\frac{2}{3}}}{2} = \frac{1}{2} \times 2^{\frac{2}{3}} \\ &= \frac{1}{2} \times (2^2)^{\frac{1}{3}} = \frac{1}{2} \times 4^{\frac{1}{3}} = \underline{\underline{\frac{1}{2} \sqrt[3]{4}}} \end{aligned}$$

1YGB - SYNOPSIS PAGE 1 - QUESTION 10

a) PROCEED AS FOLLOWS

$$\Rightarrow \frac{ds}{dt} = 24\pi$$

$$\Rightarrow \frac{ds}{dr} \times \frac{dr}{dt} = 24\pi$$

$$\Rightarrow 8\pi r \times \frac{dr}{dt} = 24\pi$$

$$\Rightarrow \frac{dr}{dt} = \frac{24}{8r}$$

$$\Rightarrow \frac{dr}{dt} = \frac{3}{r}$$

SURFACE AREA OF SPHERE

$$s = 4\pi r^2$$

$$\frac{ds}{dr} = 8\pi r$$

b) RELATING THE VOLUME NEXT

$$\frac{dv}{dt} = \frac{dv}{dr} \times \frac{dr}{dt}$$

$$\frac{dv}{dt} = 4\pi r^2 \times \frac{3}{r}$$

$$\frac{dv}{dt} = 12\pi r$$

$$\frac{dv}{dt} = \sqrt[3]{(12\pi r)^3}$$

$$\frac{dv}{dt} = \sqrt[3]{1728\pi^3 r^3}$$

$$\frac{dv}{dt} = \sqrt[3]{1728\pi^3} \times \frac{3V}{4\pi}$$

$$\frac{dv}{dt} = \sqrt[3]{1296\pi^2 V}$$

AS REQUIRED

VOLUME OF A SPHERE

$$V = \frac{4}{3}\pi r^3$$

$$\frac{dv}{dr} = 4\pi r^2$$

$$r^3 = \frac{3V}{4\pi}$$

c) SEPARATING VARIABLES & SOLVING THE O.D.E

$$\Rightarrow \frac{dv}{dt} = (1296\pi^2)^{\frac{1}{3}} V^{\frac{1}{3}}$$

$$\Rightarrow \frac{1}{V^{\frac{1}{3}}} dv = (1296\pi^2)^{\frac{1}{3}} dt$$

YGB - SYNOPSIS PAPER L - QUESTION 10

$$\begin{aligned} \Rightarrow \int v^{-\frac{1}{3}} dv &= \int (1296\pi^2)^{\frac{1}{3}} dt \\ \Rightarrow \frac{3}{2} v^{\frac{2}{3}} &= (1296\pi^2)^{\frac{1}{3}} t + C \\ \Rightarrow v^{\frac{2}{3}} &= \frac{2}{3} (1296\pi^2)^{\frac{1}{3}} t + C \end{aligned}$$

/ / required

d) FINALLY WHEN $t = \sqrt[3]{36} = 36^{\frac{1}{3}}$, GIVEN THAT $t=0$ $v=6\pi$

$$\begin{aligned} (64\pi)^{\frac{2}{3}} &= C \\ C &= 16\pi^{\frac{2}{3}} \end{aligned}$$

$$\therefore v^{\frac{2}{3}} = \frac{2}{3} (1296\pi^2)^{\frac{1}{3}} t + 16\pi^{\frac{2}{3}}$$

$$\Rightarrow v^{\frac{2}{3}} = \frac{2}{3} (1296\pi^2)^{\frac{1}{3}} \times 36^{\frac{1}{3}} + 16\pi^{\frac{2}{3}}$$

$$\Rightarrow v^{\frac{2}{3}} = \frac{2}{3} (46656\pi^2)^{\frac{1}{3}} + 16\pi^{\frac{2}{3}}$$

$$\Rightarrow v^{\frac{2}{3}} = \frac{2}{3} \times 36\pi^{\frac{2}{3}} + 16\pi^{\frac{2}{3}}$$

$$\Rightarrow v^{\frac{2}{3}} = 24\pi^{\frac{2}{3}} + 16\pi^{\frac{2}{3}}$$

$$\Rightarrow v^{\frac{2}{3}} = 40\pi^{\frac{2}{3}}$$

$$\Rightarrow v = (40\pi^{\frac{2}{3}})^{\frac{3}{2}}$$

$$\Rightarrow v = 40^{\frac{3}{2}} \pi$$

$$\Rightarrow v = 40\sqrt{40}\pi$$

/ /

1YGB - SYNOPSIS PAPER L - QUESTION 11

USING THE SUBSTITUTION GIVEN

$$\Rightarrow u = 1 + x^2 \operatorname{cosec} x$$

$$\Rightarrow \frac{du}{dx} = 2x \operatorname{cosec} x - x^2 \operatorname{cosec} x \cot x$$

$$\Rightarrow \frac{du}{dx} = x \operatorname{cosec} x (2 - x \cot x)$$

$$\Rightarrow du = \frac{du}{(2 - x \cot x)(x \operatorname{cosec} x)}$$

SUBSTITUTE INTO THE INTEGRAL

$$\int \frac{2x - x^2 \cot x}{x^2 + \sin x} dx = \int \frac{\cancel{x(2 - x \cot x)}}{x^2 + \sin x} \times \frac{du}{\cancel{(2 - x \cot x)} x \operatorname{cosec} x}$$

$$= \int \frac{1}{(x^2 + \sin x) \operatorname{cosec} x} du$$

$$= \int \frac{1}{x^2 \operatorname{cosec} x + 1} du$$

$$= \int \frac{1}{u} du$$

$$= \ln |u| + C$$

$$= \underline{\underline{\ln |1 + x^2 \operatorname{cosec} x| + C}}$$

LYGB - SYNOPTIC PART L - QUESTION 12

a) AS THE TWO EXPRESSIONS OF $f(x)$ ARE IDENTICAL, WE MAY TRY DIFFERENT SENSIBLE VALUES OF x TO ELIMINATE

$$f(x) \equiv Ax^5 + Bx^4 + 8x^2 \equiv (x-1)(x-2)g(x) + 169x - 82$$

$$f(2) \equiv 32A + 16B + 32 = 0 + 338 - 82$$

$$f\left(\frac{1}{2}\right) \equiv \frac{1}{32}A + \frac{1}{16}B + 2 = 0 + \frac{169}{2} - 82$$

TIDY THE EQUATIONS & SOLVE

$$\left. \begin{aligned} 32A + 16B &= 224 \\ \frac{1}{32}A + \frac{1}{16}B &= \frac{1}{2} \end{aligned} \right\} \Rightarrow \begin{aligned} 2A + B &= 14 \\ A + 2B &= 16 \end{aligned}$$

$$\boxed{B = 14 - 2A}$$

$$\Rightarrow A + 2(14 - 2A) = 16$$

$$\Rightarrow A + 28 - 4A = 16$$

$$\Rightarrow 2 = 3A$$

$$\Rightarrow \underline{A = 4} \quad \text{and} \quad \underline{B = 6}$$

b) Using the answers from part (a)

$$\Rightarrow 4x^5 + 6x^4 + 8x^2 \equiv (x-1)(x-2)g(x) + 169x - 82$$

$$\Rightarrow 4x^5 + 6x^4 + 8x^2 - 169x + 82 \equiv (x-1)(x-2)g(x)$$

$$\Rightarrow 4x^5 + 6x^4 + 8x^2 - 169x + 82 \equiv (2x^2 - 5x + 2)g(x)$$

1YGB - SYNOPTIC PAPER L - QUESTION 12

BY LONG DIVISION

$$\begin{array}{r}
 2x^3 + 8x^2 + 18x + 41 \\
 \hline
 4x^5 + 6x^4 + 0x^3 + 8x^2 - 164x + 82 \\
 -4x^5 + 10x^4 - 4x^3 \\
 \hline
 16x^4 - 4x^3 + 8x^2 - 164x + 82 \\
 -16x^4 + 40x^3 - 16x^2 \\
 \hline
 36x^3 - 8x^2 - 164x + 82 \\
 -36x^3 + 90x^2 - 36x \\
 \hline
 82x^2 - 205x + 82 \\
 -82x^2 + 205x - 82 \\
 \hline
 0
 \end{array}$$

$\therefore g(x) = 2x^3 + 8x^2 + 18x + 41$

c) PROCEED AS FOLLOWS

$$\begin{cases}
 f(x) \equiv 4x^5 + 6x^4 + 8x^2 \equiv (x+2)^2 h(x) + Px + Q \\
 f'(x) \equiv 20x^4 + 24x^3 + 16x \equiv 2(x+2)h(x) + (x+2)^2 h'(x) + P
 \end{cases}$$


$$\begin{cases}
 f(-2) = -128 + 96 + 32 = 0 = -2P + Q \\
 f'(-2) = 320 - 192 - 32 = P
 \end{cases}$$

$\therefore P = 96$

$$\begin{aligned}
 \text{q } Q - 2P &= 0 \\
 Q &= 2P \\
 Q &= 192
 \end{aligned}$$

1YGB - SYNOPTIC PART 2 L - QUESTION 13

a) THIS IS A TRANSLATION OF $y = \frac{1}{x}$ BY 1 UNIT TO THE RIGHT



b) LOOKING FOR "INTERSECTIONS", ALTHOUGH WE ARE TOLD THERE ARE NOT ANY

$$\begin{aligned}
 \left. \begin{aligned} y &= \frac{1}{x-1} \\ y &= a-2x \end{aligned} \right\} & \Rightarrow \frac{1}{x-1} = a-2x \\
 & \Rightarrow 1 = (x-1)(a-2x) \\
 & \Rightarrow 1 = ax - 2x^2 - a + 2x \\
 & \Rightarrow 2x^2 - ax - 2x + 1 + a = 0 \\
 & \Rightarrow 2x^2 - (a+2)x + (a+1) = 0
 \end{aligned}$$

BUT THIS EQUATION HAS NO REAL ROOTS (NO INTERSECTIONS)

$$\begin{aligned}
 & \Rightarrow B^2 - 4AC < 0 \\
 & \Rightarrow [-(a+2)]^2 - 4 \times 2 \times (a+1) < 0 \\
 & \Rightarrow (a+2)^2 - 8(a+1) < 0
 \end{aligned}$$

- 2 -

1YGB - SYNOPTIC PAPER L - QUESTION 13

$$\Rightarrow a^2 + 4a + 4 - 8a - 8 < 0$$

$$\Rightarrow a^2 - 4a - 4 < 0$$

$$\Rightarrow (a - 2)^2 - 8 < 0$$

$$\Rightarrow (a - 2)^2 < 8$$

$$\Rightarrow -\sqrt{8} < a - 2 < \sqrt{8}$$

$$\Rightarrow 2 - \sqrt{8} < a < 2 + \sqrt{8}$$

$$\therefore \underline{2 - 2\sqrt{2} < a < 2 + 2\sqrt{2}}$$

1XGB - SYNOPTIC PAPER L - QUESTION 14

a) REWRITE IN INDEX NOTATION & DIFFERENTIATE

$$y = \ln(x^{\frac{1}{2}}) + (\ln x)^{\frac{1}{2}}$$

$$\frac{dy}{dx} = \frac{1}{x^{\frac{1}{2}}} \times \frac{1}{2} x^{-\frac{1}{2}} + \frac{1}{2} (\ln x)^{-\frac{1}{2}} \times \frac{1}{x}$$

$$\frac{dy}{dx} = \frac{1}{2x} + \frac{1}{2x} (\ln x)^{-\frac{1}{2}}$$

$$\frac{dy}{dx} = \frac{1}{2x} + \frac{1}{2x\sqrt{\ln x}}$$

ALTERNATIVE FOR THE FIRST TERM

$$\frac{d}{dx} [\ln \sqrt{x}] = \frac{d}{dx} [\ln(x^{\frac{1}{2}})] = \frac{d}{dx} [\frac{1}{2} \ln x] = \frac{1}{2x}$$

b) DIFFERENTIATING THE PRODUCT

$$y = (2x+1)^{\frac{1}{2}} (1-4x)^{-\frac{1}{2}}$$

$$\frac{dy}{dx} = (2x+1)^{-\frac{1}{2}} (1-4x)^{-\frac{1}{2}} + (2x+1)^{\frac{1}{2}} \times (-\frac{1}{2}) (1-4x)^{-\frac{3}{2}} (-4)$$

$$\frac{dy}{dx} = (2x+1)^{-\frac{1}{2}} (1-4x)^{-\frac{1}{2}} + 2(2x+1)^{\frac{1}{2}} (1-4x)^{-\frac{3}{2}}$$

$$\frac{dy}{dx} = (2x+1)^{-\frac{1}{2}} (1-4x)^{-\frac{3}{2}} [(1-4x) + 2(2x+1)]$$

$$\frac{dy}{dx} = (2x+1)^{-\frac{1}{2}} (1-4x)^{-\frac{3}{2}} (1-4x+4x+2)$$

$$\frac{dy}{dx} = 3(2x+1)^{-\frac{1}{2}} (1-4x)^{-\frac{3}{2}}$$

→ -

YGB-SYNOPTIC PAPER L - QUESTION 14

Solving for zero

$$\Rightarrow 3(1+x)^{-\frac{1}{2}}(1-4x)^{-\frac{3}{2}} = 0$$

$$\Rightarrow \frac{3}{2(1+x)^{\frac{1}{2}}(1-4x)^{\frac{3}{2}}} = 0$$

No solutions of hence no turning points

d By the quotient rule

$$\begin{aligned} \frac{d}{dx} \left[\frac{2x-1}{(2x+1)^{\frac{1}{2}}} \right] &= \frac{(2x+1)^{\frac{1}{2}} \times 2 - (2x-1)(2x+1)^{-\frac{1}{2}} \times \frac{1}{2} \times 2}{(2x+1)^1} \\ &= \frac{2(2x+1)^{\frac{1}{2}} - (2x-1)(2x+1)^{\frac{1}{2}}}{2x+1} \\ &= \frac{(2x+1)^{-\frac{1}{2}} [2(2x+1) - (2x-1)]}{2x+1} \\ &= \frac{(2x+1)^{-\frac{1}{2}} (4x+2-2x+1)}{2x+1} = \frac{2x+3}{(2x+1)^{\frac{1}{2}}(2x+1)} \\ &= \frac{2x+3}{(2x+1)^{\frac{3}{2}}} \end{aligned}$$

As required

YGB - SYNOPSIS PAGE L - QUESTION IS

START BY FINDING THE GRADIENT AT P

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{4\cos\theta}{-8\sin\theta} = -\frac{1}{2}\cot\theta$$

$$\left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{4}} = -\frac{1}{2}\cot\frac{\pi}{4} = -\frac{1}{2}$$

OBTAIN THE EQUATION OF THE TANGENT

$$\bullet x \Big|_{\frac{\pi}{4}} = 8\cos\frac{\pi}{4} = 8 \times \frac{\sqrt{2}}{2} = 4\sqrt{2}$$

$$\bullet y \Big|_{\frac{\pi}{4}} = 4\sin\frac{\pi}{4} = 4 \times \frac{\sqrt{2}}{2} = 2\sqrt{2}$$

i.e. $P(4\sqrt{2}, 2\sqrt{2})$


$$y - y_0 = m(x - x_0)$$

$$y - 2\sqrt{2} = -\frac{1}{2}(x - 4\sqrt{2})$$

$$2y - 4\sqrt{2} = -x + 4\sqrt{2}$$

$$2y + x = 8\sqrt{2}$$

NEXT WE FIND THE AREA OF THE TRIANGLE
IN THE FOLLOWING DIAGRAM



$$\Delta \text{ AREA} = \frac{1}{2} \times 2\sqrt{2} \times 4\sqrt{2} = 8$$

BY SETTING $y=0$
IN THE EQUATION
OF THE TANGENT

FINALLY THE AREA SHOWN SHADDED IN THE ABOVE
DIAGRAM (AREA BETWEEN CURVE & X AXIS)


FIRSTLY BY INSPECTION, THE CURVE MEETS

THE X AXIS AT $x=8$, i.e. $\theta=0$

-2-

YGB - QUESTION 15 - SHORTCUT PAPER L

$$\text{AREA} = \int_{x_1}^{x_2} y(x) dx = \int_{\theta_1}^{\theta_2} y(\theta) \frac{dx}{d\theta} d\theta$$



$$\text{AREA} = \int_0^{\pi/4} 4\sin\theta (-8\sin\theta) d\theta$$

$$\text{AREA} = \int_0^{\pi/4} -32\sin^2\theta d\theta = \int_0^{\pi/4} 32\left(\frac{1}{2} - \frac{1}{2}\cos 2\theta\right) d\theta$$

$$\text{AREA} = \int_0^{\pi/4} 16 - 16\cos 2\theta d\theta = [16\theta - 8\sin 2\theta]_0^{\pi/4}$$

$$\text{AREA} = (4\pi - 8) - (0)$$

$$\text{AREA} = 4\pi - 8$$

FINALLY THE REQUIRED AREA CAN BE FOUND

$$\Rightarrow \text{AREA OF TRIANGLE} - \text{AREA UNDER CURVE} = 8 - (4\pi - 8)$$

$$= 16 - 4\pi$$

IXGB - SYNOPTIC PAPER L - QUESTION 16

● IF $f(x) = \sqrt{1+x^2}$ THEN $f(x+h) = \sqrt{1+(x+h)^2}$

● BY THE FORMAL DEFINITION OF THE DERIVATIVE

$$f'(x) = \lim_{h \rightarrow 0} \left[\frac{f(x+h) - f(x)}{h} \right] = \lim_{h \rightarrow 0} \left[\frac{\sqrt{1+(x+h)^2} - \sqrt{1+x^2}}{h} \right]$$

$$= \lim_{h \rightarrow 0} \left[\frac{[\sqrt{1+(x+h)^2} - \sqrt{1+x^2}] [\sqrt{1+(x+h)^2} + \sqrt{1+x^2}]}{h [\sqrt{1+(x+h)^2} + \sqrt{1+x^2}]} \right]$$

$$= \lim_{h \rightarrow 0} \left[\frac{[1+(x+h)^2] - [1+x^2]}{h [\sqrt{1+(x+h)^2} + \sqrt{1+x^2}]} \right]$$

$$= \lim_{h \rightarrow 0} \left[\frac{x^2 + 2xh + h^2 - x^2}{h [\sqrt{1+(x+h)^2} + \sqrt{1+x^2}]} \right]$$

$$= \lim_{h \rightarrow 0} \left[\frac{h(2x+h)}{h [\sqrt{1+(x+h)^2} + \sqrt{1+x^2}]} \right]$$

$$= \lim_{h \rightarrow 0} \left[\frac{2x+h}{\sqrt{1+(x+h)^2} + \sqrt{1+x^2}} \right]$$

$$= \frac{2x}{\sqrt{1+x^2} + \sqrt{1+x^2}}$$

$$= \frac{2x}{2\sqrt{1+x^2}}$$

$$= \frac{x}{\sqrt{1+x^2}}$$

IYGB - SYNOPTIC PAPER L - QUESTION 17.

a) LOOKING AT THE DIAGRAM


$$\frac{|XB|}{|XO|} = \tan \frac{\pi}{8}$$

$$\frac{|XB|}{1} = \tan \frac{\pi}{8}$$

$$|XB| = \tan \frac{\pi}{8}$$

$$|AB| = 2 \tan \frac{\pi}{8}$$

$$\underline{\text{PERIMETER OF OCTAGON}} = 8 \times 2 \tan \frac{\pi}{8} = 16 \tan \frac{\pi}{8}$$



$$360 \div 8 = 45 \text{ or } \frac{\pi}{4}$$

b) LOOKING AT THE DIAGRAM AGAIN


$$\frac{|YN|}{|ON|} = \sin \frac{\pi}{8}$$

$$\frac{|YN|}{1} = \sin \frac{\pi}{8}$$

$$|YN| = \sin \frac{\pi}{8}$$

$$|MN| = 2 \sin \frac{\pi}{8}$$

$$\underline{\text{PERIMETER OF OCTAGON}} = 8 \times 2 \sin \frac{\pi}{8} = 16 \sin \frac{\pi}{8}$$



c) USING $\cos 2\theta \equiv 1 - 2\sin^2 \theta$ WITH $\theta = \frac{\pi}{8}$

$$\Rightarrow \cos \frac{\pi}{4} = 1 - 2\sin^2 \frac{\pi}{8}$$

$$\Rightarrow 2\sin^2 \frac{\pi}{8} = 1 - \cos \frac{\pi}{4}$$

$$\Rightarrow 2\sin^2 \frac{\pi}{8} = 1 - \frac{\sqrt{2}}{2}$$

$$\Rightarrow \sin^2 \frac{\pi}{8} = \frac{1}{2} - \frac{\sqrt{2}}{4}$$

$$\Rightarrow \sin^2 \frac{\pi}{8} = \frac{2 - \sqrt{2}}{4}$$

19GB - SYNOPTIC PAPER L - QUESTION 17

$$\Rightarrow \sin \frac{\pi}{8} = + \sqrt{\frac{2 - \sqrt{2}}{4}} \quad \left(\frac{\pi}{8} \text{ is acute} \right)$$

$$\Rightarrow \sin \frac{\pi}{8} = \frac{1}{2} \sqrt{2 - \sqrt{2}} \quad \text{AS REQUIRED}$$

d) USING THE DOUBLE ANGLE IDENTITY FOR $\tan 2\theta$ WITH $\theta = \frac{\pi}{8}$

$$\Rightarrow \tan \frac{\pi}{4} = \frac{2 \tan \frac{\pi}{8}}{1 - \tan^2 \frac{\pi}{8}}$$

$$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$$

$$\Rightarrow 1 = \frac{2T}{1 - T^2}$$

$$T = \tan \frac{\pi}{8}$$

$$\Rightarrow 1 - T^2 = 2T$$

$$\Rightarrow 0 = T^2 + 2T - 1$$

$$\Rightarrow (T+1)^2 - 2 = 0$$

$$\Rightarrow (T+1)^2 = 2$$

$$\Rightarrow T+1 = \pm \sqrt{2}$$

$$\Rightarrow T = -1 \pm \sqrt{2}$$

$$\Rightarrow \tan \frac{\pi}{8} = -1 \pm \sqrt{2}$$

$$\Rightarrow \tan \frac{\pi}{8} = -1 + \sqrt{2} \quad \text{(AS } \frac{\pi}{8} \text{ IS ACUTE \& } -1 - \sqrt{2} \text{ IS NEGATIVE)}$$


FINALLY WE HAVE

PERIMETER P_2	$<$	CIRCUMFERENCE C	$<$	PERIMETER P_1
$16 \sin \frac{\pi}{8}$		$2\pi \times 1$		$16 \tan \frac{\pi}{8}$
$16 \left(\frac{1}{2} \sqrt{2 - \sqrt{2}} \right)$	$<$	2π	$<$	$(-1 + \sqrt{2}) \times 16$
<u>3.06</u>	$<$	<u>π</u>	$<$	<u>3.31</u>

YGB - SYNOPTIC PAPER 1 - QUESTION 19

START BY SKETCHING THE FUNCTION

$$f(x) = \begin{cases} \frac{5}{2} - x, & x \in \mathbb{R}, -10 < x < 2 \\ \frac{2}{x^2}, & x \in \mathbb{R}, 2 \leq x \leq 4 \end{cases}$$



TREAT EACH SECTION SEPARATELY

$$f_1(x) = \frac{5}{2} - x, \quad -10 < x < 2$$

$$y = \frac{5}{2} - x$$

$$2y = 5 - 2x$$

$$2x = 5 - 2y$$

$$x = \frac{5}{2} - y$$

$$f_1^{-1}(x) = \frac{5}{2} - x \quad (\text{SWAP INVERSE})$$

	$f_1(x)$	$f_1^{-1}(x)$
D	$-10 < x < 2$	$\frac{1}{2} < x < \frac{25}{2}$
R	$\frac{1}{2} < f_1(x) < \frac{25}{2}$	$-10 < f_1^{-1}(x) < 2$

$$f_2(x) = \frac{2}{x^2}, \quad 2 \leq x \leq 4$$

$$y = \frac{2}{x^2}$$

$$x^2 = \frac{2}{y}$$

$$x = \pm \sqrt{\frac{2}{y}}$$

$$x = + \sqrt{\frac{2}{y}}$$

$$f_2^{-1}(x) = \sqrt{\frac{2}{x}}$$

	$f_2(x)$	$f_2^{-1}(x)$
D	$2 \leq x \leq 4$	$\frac{1}{8} \leq x \leq \frac{1}{2}$
R	$\frac{1}{8} \leq f_2(x) \leq \frac{1}{2}$	$2 \leq f_2^{-1}(x) \leq 4$

$$f^{-1}(x) = \begin{cases} \sqrt{\frac{2}{x}}, & x \in \mathbb{R}, \frac{1}{8} \leq x \leq \frac{1}{2} \\ \frac{5}{2} - x, & x \in \mathbb{R}, \frac{1}{2} < x < \frac{25}{2} \end{cases}$$

WITH RANGE $-10 \leq f^{-1}(x) \leq 4$

1YGB - SYNOPSIS PAPER L - QUESTION 11

$f(n) = 10, 13, 16, 19, 22, \dots$	$3n + 7$
$g(n) = 6, 12, 24, 48, 96, \dots$	3×2^n

ADDING THE n^{th} TERMS

$$u_n = f(n) + g(n) = 3n + 7 + 3 \times 2^n$$

$$u_{n+1} = 3(n+1) + 7 + 3 \times 2^{n+1} = 3n + 10 + 3 \times 2^{n+1}$$

SUBTRACTING GIVES

$$\begin{aligned} u_{n+1} - u_n &= (3n + 10 + 3 \times 2^{n+1}) - (3n + 7 + 3 \times 2^n) \\ &= \cancel{3n} + 10 + 3 \times 2^{n+1} - \cancel{3n} - 7 - 3 \times 2^n \\ &= 3 + 3 \times 2 \times 2^n - 3 \times 2^n \\ &= 3 + 6 \times 2^n - 3 \times 2^n \\ &= 3 + 3 \times 2^n \\ &= 3(1 + 2^n) \end{aligned}$$

FINALLY WE HAVE

$$\begin{aligned} u_{n+1} - u_n &= 3(1 + 2^n) \\ \underline{u_{n+1}} &= \underline{u_n + 3(2^n + 1)} \end{aligned}$$

As required