

1YGB - MPI PAPER Z - QUESTION 1

USING STANDARD REARRANGING TECHNIQUES

$$\frac{\sqrt{p}}{2p + \sqrt{p}} = \frac{2\sqrt{p} - q}{3p + q}$$

$$\Rightarrow \sqrt{p}(3p + q) = (2p + \sqrt{p})(2\sqrt{p} - q)$$

$$\Rightarrow 3p\sqrt{p} + q\sqrt{p} = 4p\sqrt{p} - 2pq + 2p - q\sqrt{p}$$

$$\Rightarrow 2q\sqrt{p} + 2pq = p\sqrt{p} + 2p$$

$$\Rightarrow q[2\sqrt{p} + 2p] = p\sqrt{p} + 2p$$

$$\Rightarrow q = \frac{p\sqrt{p} + 2p}{2\sqrt{p} + 2p}$$

$$\Rightarrow q = \frac{p\sqrt{p} + 2\sqrt{p}\sqrt{p}}{2\sqrt{p} + 2\sqrt{p}\sqrt{p}} = \frac{p + 2\sqrt{p}}{2 + 2\sqrt{p}} \quad \text{As required}$$

ALTERNATIVE

$$\frac{\sqrt{p}}{2p + \sqrt{p}} = \frac{2\sqrt{p} - q}{3p + q}$$

$$\Rightarrow \frac{1}{2\sqrt{p} + 1} = \frac{2\sqrt{p} - q}{3p + q}$$

DIVIDE "TOP & BOTTOM" OF THE FRACTION IN THE L.H.S BY \sqrt{p}

$$\Rightarrow 3p + q = (2\sqrt{p} - q)(2\sqrt{p} + 1)$$

$$\Rightarrow 3p + q = 4p + 2\sqrt{p} - 2q\sqrt{p} - q$$

$$\Rightarrow 2q + 2q\sqrt{p} = p + 2\sqrt{p}$$

$$\Rightarrow q(2 + 2\sqrt{p}) = p + 2\sqrt{p}$$

$$\Rightarrow q = \frac{p + 2\sqrt{p}}{2 + 2\sqrt{p}}$$

As before

1YGB - MPI PAPER 2 - QUESTION 2

LOOKING AT THE DIAGRAM ON THE TRIANGLE AMO

BY PYTHAGORAS


$$\Rightarrow |AM|^2 + |MO|^2 = |OA|^2$$

$$\Rightarrow 12^2 + (72-r)^2 = r^2$$

$$\Rightarrow 144 + 5184 - 144r + r^2 = r^2$$

$$\Rightarrow 5328 = 144r$$

$$\Rightarrow r = 37$$



1
19GB - MPAI PAPER 2 - QUESTION 3

EXPAND THE EXPRESSION UP TO x^2 , IN TERMS OF a & b

$$\begin{aligned} (2+ax)^2(1+bx)^6 &= (4 + 4ax + a^2x^2) \left[1 + \frac{6}{1}(bx) + \frac{6 \times 5}{1 \times 2}(bx)^2 + \dots \right] \\ &= (4 + 4ax + a^2x^2)(1 + 6bx + 15b^2x^2 + \dots) \\ &= 4 + 24bx + 60b^2x^2 + \dots \\ &\quad 4ax + 24abx^2 + \dots \\ &\quad \quad \quad a^2x^2 + \dots \\ &= 4 + (4a + 24b)x + (60b^2 + 2(4ab + a^2))x^2 + \dots \end{aligned}$$

COMPARING COEFFICIENTS

● $4a + 24b = 44$
 $a + 6b = 11$
 $a = 11 - 6b$

● $60b^2 + 24ab + a^2 = 85$

$\rightarrow 60b^2 + 24b(11 - 6b) + (11 - 6b)^2 = 85$
 $60b^2 + 264b - 144b^2 + 121 - 132b + 36b^2 = 85$
 $- 48b^2 + 132b + 36 = 0$
 $48b^2 - 132b - 36 = 0$
 $4b^2 - 11b - 3 = 0$
 $(4b + 1)(b - 3) = 0$
 $b = \begin{cases} 3 \\ -\frac{1}{4} \end{cases}$

$a = \begin{cases} 11 - 6 \times 3 = -7 \\ 11 - 6 \left(-\frac{1}{4}\right) = \frac{25}{2} \end{cases}$

Either $a=7$ & $b=3$

or $a = \frac{25}{2}$ & $b = -\frac{1}{4}$

- 1 -

IYGB - MPI PAPER Z - QUESTION 4

REARRANGE THE EXPRESSION AS A QUADRATIC IN x

$$f(x) = x^2 + 2x - m(x^2 - 2x + 2) - 2$$

$$f(x) = x^2 + 2x - mx^2 + 2mx - 2m - 2$$

$$f(x) = (1-m)x^2 + (2+2m)x + (-2m-2)$$

$\begin{matrix} \uparrow & \uparrow & \uparrow \\ a & b & c \end{matrix}$

FOR DISTINCT REAL ROOTS OF $f(x)=0$, $b^2 - 4ac > 0$

$$\Rightarrow (2+2m)^2 - 4(1-m)(-2m-2) > 0$$

$$\Rightarrow 4(1+m)^2 + 4(1-m)(2m+2) > 0$$

$$\Rightarrow (1+m)^2 + (1-m)(2m+2) > 0$$

$$\Rightarrow 1 + 2m + m^2 + 2m + 2 - 2m^2 - 2m > 0$$

$$\Rightarrow -m^2 + 2m + 3 > 0$$

$$\Rightarrow m^2 - 2m - 3 < 0$$

$$\Rightarrow (m-3)(m+1) < 0$$

CRITICAL VALUES $\begin{matrix} & -1 \\ & / \\ & 3 \end{matrix}$



$$\underline{-1 < m < 3}$$


$$\underline{m \neq 1}$$

OR $\{-1 < m < 1\} \cup \{1 < m < 3\}$

1YGB - MPI PAPER 2 - QUESTION 5


$|a| = 4$ $|b| = 6$ $\hat{AOB} = 40^\circ$

START WITH A DIAGRAM



REDRAW SHOWING THE POSITION OF P

$\vec{OP} = 2a - 3b$



BY THE COSINE RULE

$$|P|^2 = 18^2 + 8^2 - 2 \times 18 \times 8 \times \cos 40^\circ$$

$$|P|^2 = 324 + 64 - 288 \cos 40^\circ$$

$$|P|^2 = 167.379\dots$$

$$|P| = 12.93751\dots$$

$\therefore |OP| = 12.94$

BY THE SINE RULE:

$$\frac{|P|}{\sin 40^\circ} = \frac{8}{\sin \theta}$$

$$\sin \theta = \frac{8 \sin 40^\circ}{12.93751\dots}$$

$$\theta \approx 23.42^\circ\dots$$

∴ REQUIRED ANGLE ϕ
 $180 - (40 + 23.42^\circ)$
 $\approx 117^\circ$

- 1 -

1YGB - MPI PAPER 2 - QUESTION 6

a) $f(x) = 2\log_4 x$

$g(x) = 1 + 2\log_4 x = 1 + f(x) = f(x) + 1$

∴ TRANSFORMATION BY $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$

b) $g(x) = 1 + 2\log_4 x$

$g(x) = \log_4 4 + \log_4 x^2$

$g(x) = \log_4 (4x^2)$

$g(x) = \log_4 [(2x)^2]$

$g(x) = 2\log_4 (2x)$

$g(x) = f(2x)$

OR

$g(x) = \log_4 4 + 2\log_4 x$

$g(x) = \log_4 2^2 + 2\log_4 x$

$g(x) = 2\log_4 2 + 2\log_4 x$

$g(x) = 2[\log_4 2 + \log_4 x]$

$g(x) = 2\log_4 (2x)$

∴ HORIZONTAL STRETCH BY SCALE FACTOR $\frac{1}{2}$
(STRETCH PARALLEL TO THE x AXIS, BY S.F $\frac{1}{2}$)

IYGB - MPI PAPER 2 - QUESTION 7

THE EQUATION OF l IS GIVEN BY

$$\begin{aligned}y - y_0 &= m(x - x_0) \\y - 5 &= 3(x - 4) \\y - 5 &= 3x - 12 \\y &= 3x - 7\end{aligned}$$

LET THE REQUIRED POINT HAVE CO-ORDINATES Q(a, 3a-7)

$$\begin{aligned}\Rightarrow |PQ| &= 3\sqrt{10} \\ \Rightarrow \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} &= 3\sqrt{10} \\ \Rightarrow \sqrt{(a - 4)^2 + (3a - 7 - 5)^2} &= 3\sqrt{10} \\ \Rightarrow \sqrt{(a - 4)^2 + (3a - 12)^2} &= 3\sqrt{10} \\ \Rightarrow (a - 4)^2 + (3a - 12)^2 &= 9 \times 10\end{aligned}$$

P(4, 5)
Q(a, 3a-7)

$$\begin{aligned}\Rightarrow \left. \begin{aligned}a^2 - 8a + 16 \\ 9a^2 - 72a + 144\end{aligned} \right\} &= 90 \\ \Rightarrow 10a^2 - 80a + 160 &= 90 \\ \Rightarrow 10a^2 - 80a + 70 &= 0 \\ \Rightarrow a^2 - 8a + 7 &= 0 \\ \Rightarrow (a - 7)(a - 1) &= 0\end{aligned}$$


$$a = \begin{cases} 1 \\ 7 \end{cases}$$

$$3a - 7 = \begin{cases} -4 \\ 14 \end{cases}$$


\therefore Q(1, -4)
OR
Q(7, 14)

1YGB - MPI PAPER 2 - QUESTION 7

ALTERNATIVE BY GEOMETRY



• GRADIENT 3



• DISTANCE $3\sqrt{10}$ IS

3 "GRADIENT" STAIRS FROM P

• SEE DIAGRAM FOR THE ANSWERS

Q(1, -4) OR Q(7, 14)

1YGB - MPI PAPER 2 - QUESTIONS 8

ALTHOUGH IT IS "TEMPTING" TO START FROM THE TWO CONDITIONS
GIVEN ($t=0, h=2$ & $t=2, h=3.81$), IT BEST TO START FROM

"LIFETIME MAX HEIGHT OF 12m" \Rightarrow As $t \rightarrow \infty$ $h \rightarrow 12$
 \Rightarrow As $t \rightarrow \infty$ $e^{-kt} \rightarrow 0$
 $Be^{-kt} \rightarrow 0$
 $h \rightarrow A$

$\therefore \underline{\underline{A=12}}$

$$h = 12 - Be^{-kt}$$

$t=0, h=2$

$$2 = 12 - Be^0$$

$$\underline{\underline{B=10}}$$

$t=2, h=3.81$

$$3.81 = 12 - Be^{-2k}$$

$$3.81 = 12 - 10e^{-2k}$$

$$10e^{-2k} = 8.19$$

$$e^{-2k} = 0.819$$

$$-2k = \ln(0.819)$$

$$\underline{\underline{k = 0.099835 \dots \approx 0.1}}$$

FINALLY WE HAVE

$$\Rightarrow h = 12 - 10e^{-0.1t}$$

$$\Rightarrow 10 = 12 - 10e^{-0.1t}$$

$$\Rightarrow 10e^{-0.1t} = 2$$

$$\Rightarrow e^{-0.1t} = \frac{1}{5}$$

$$\Rightarrow e^{0.1t} = 5$$



$$\Rightarrow 0.1t = \ln 5$$

$\therefore \underline{\underline{t = 10 \ln 5 \approx 16.1}}$


- 1 -

1YGB - MPI PAPER 2 Z - QUESTION 9

a) ● $C_1: y = (2x-3)^3$



● $C_2: y = (2x-1)^2 - 4$
 $y = (2x-1-2)(2x-1+2)$
 $y = (2x-3)(2x+1)$
 $(\frac{3}{2}|0), (-\frac{1}{2}|0), (0|-3)$



b) $\Rightarrow (2x-1)^2 + (3-2x)^3 = 4$
 $\Rightarrow (2x-1)^2 - 4 = -(3-2x)^3$
 $\Rightarrow (2x-1)^2 - 4 = +(2x-3)^3$

USING THE FACTORIZATION FOR THE QUADRATIC FROM ABOVE

$$\Rightarrow (2x-3)(2x+1) = (2x-3)^3$$

PREFERABLY NOT EXPANDING AS THERE IS A COMMON FACTOR

$$\Rightarrow 0 = (2x-3)^3 - (2x-3)(2x+1) = 0$$

$$\Rightarrow 0 = (2x-3) [(2x-3)^2 - (2x+1)]$$

$$\Rightarrow 0 = (2x-3) [4x^2 - 12x + 9 - 2x - 1]$$

1YGB - MPI PAPER 2 - QUESTION 9

$$\Rightarrow 0 = (2x-3)(4x^2-14x+8)$$

EITHER

$$2x-3=0$$

$$2x=3$$

$$x = \frac{3}{2}$$

OR

$$4x^2-14x+8=0$$

$$2x^2-7x+4=0$$

QUADRATIC FORMULA

$$\Delta = (-7)^2 - 4 \times 2 \times 4$$

$$\Delta = 49 - 32$$

$$\Delta = 17$$

$$x = \frac{-(-7) \pm \sqrt{17}}{2 \times 2}$$

$$x = \frac{7 \pm \sqrt{17}}{4}$$

THUS THE THREE SOLUTIONS ARE

$$x = \begin{cases} \frac{3}{2} \\ \frac{1}{4}(7 + \sqrt{17}) \\ \frac{1}{4}(7 - \sqrt{17}) \end{cases}$$

IYGB - MFL PAPER 2 - QUESTION 10

a) SOLVING SIMULTANEOUSLY

$$\left. \begin{aligned} x^2 + y^2 - 6x &= 16 \\ x^2 + y^2 - 18x + 16y &= 80 \end{aligned} \right\} \text{SUBTRACT}$$

$$\begin{aligned} 12x - 16y &= -64 \\ 16y - 12x &= 64 \\ 4y - 3x &= 16 \\ 3x &= 4y - 16 \\ 9x^2 &= 16y^2 - 128y + 256 \end{aligned}$$

NOW MULTIPLYING THE FIRST EQUATION BY 9

$$\begin{aligned} 9x^2 + 9y^2 - 54x &= 144 \\ 9x^2 + 9y^2 - 18(3x) &= 144 \\ 16y^2 - 128y + 256 + 9y^2 - 18(4y - 16) &= 144 \\ 16y^2 - 128y + 256 + 9y^2 - 72y + 288 &= 144 \\ 25y^2 - 200y + 400 &= 0 \\ y^2 - 8y + 16 &= 0 \\ (y - 4)^2 &= 0 \end{aligned}$$

$3x = 4y - 16$

$y = 4$ REPEATS, INDEED THEY TOUCH AT $(0, 4)$

b) FIRSTLY WE NEED THE CIRCLE PARTICULARS

$$\begin{aligned} x^2 + y^2 - 6x &= 16 \\ (x-3)^2 - 9 + y^2 &= 16 \\ (x-3)^2 + y^2 &= 25 \\ (3, 0), \text{ radius } 5 \end{aligned}$$

$$\begin{aligned} x^2 + y^2 - 18x + 16y &= 80 \\ x^2 - 18x + y^2 + 16y &= 80 \\ (x-9)^2 - 81 + (y+8)^2 - 64 &= 80 \\ (x-9)^2 + (y+8)^2 &= 225 \\ (9, -8), \text{ radius } 15 \end{aligned}$$

1 YGB, MPI PAPER 2 - QUESTION 10

DISTANCE BETWEEN THE CENTRES (3,0) & (9,-8)

$$d = \sqrt{(3-9)^2 + (0+8)^2}$$


$$d = \sqrt{36 + 64}$$

$$d = \sqrt{100}$$

$$d = 10$$

TOUCHING EXTERNALLY REQUIRES $d = 15 + 5 = 20$

TOUCHING INTERNALLY REQUIRES $d = 15 - 5 = 10$

∴ INTERNALLY 

1YGB - MFL PAPER 2 - QUESTION 11

a) LINE: $5x + 2y = 9$

CUBIC: $y = x^3 - 4x^2 + px + 4$

$\frac{dy}{dx} = 3x^2 - 8x + p$

WHEN $x=1$ THE y CO-ORDS ARE IDENTICAL (POINT A)

• $5 + 2y = 9$

• $y = 1 - 4 + p + 4$
 $y = p + 1$

THE GRADIENT AT B, WHERE $x=2$, IS ZERO

$\Rightarrow 0 = 12 - 16 + p$

$\Rightarrow \underline{p = 4}$

9 THE y CO-ORD OF A IS 5 ($y = p + 1$)

$\Rightarrow 5 + 2y = 9$

$\Rightarrow 5 + 10 = 9$

$\Rightarrow \underline{9 = 15}$

b) FIND THE x INTERCEPT OF L


$5x + 2y = 15$

$5x + 2 \times 0 = 15$

$x = 3$

19GB - MPA PAPER 2 - QUESTION 11

Now looking at the "PICTORIAL" equation



$$\int_0^1 (x^3 - 4x^2 + 4x + 4) dx = \left[\frac{1}{4}x^4 - \frac{4}{3}x^3 + 2x^2 + 4x \right]_0^1$$

$$= \left(\frac{1}{4} - \frac{4}{3} + 2 + 4 \right) - (0)$$

$$= \frac{59}{12}$$

THE REQUIRED REGION HAS AREA

$$\frac{59}{12} + 5 = \frac{119}{12}$$

-|-

1YGB - MPI PAPER 2 - QUESTION 12

ASSERTION: IF $x \in \mathbb{R}, x > 0$ THEN $x^4 + x^{-4} \geq 2$

PROOF BY CALCULUS

● LET $f(x) = x^4 + x^{-4} = x^4 + \frac{1}{x^4}$, $x \in \mathbb{R}, x > 0$

AS $x \rightarrow +\infty$, $f(x) \rightarrow +\infty$ ($f(x) \sim x^4$)

AS $x \rightarrow 0^+$, $f(x) \rightarrow +\infty$ ($f(x) \sim \frac{1}{x^4}$)

● LOOK FOR STATIONARY VALUES

$$\Rightarrow f'(x) = 4x^3 - 4x^{-5} = 4x^{-5}(x^8 - 1)$$

SOLVING FOR ZERO

$$\Rightarrow \frac{4}{x^5}(x^8 - 1) = 0$$

$$\Rightarrow x^8 - 1 = 0 \quad \left(\frac{4}{x^5} \neq 0\right)$$

$$\Rightarrow x = \pm 1 \quad \text{ONLY REAL SOLUTIONS}$$

$$\Rightarrow x = 1 \quad \text{ONLY POSITIVE REAL SOLUTION}$$

$$\therefore f(1) = 1^4 + 1^{-4} = 2$$

● AS $f(x)$ TENDS TO INFINITY AS $x \rightarrow \infty$ OR $x \rightarrow 0^+$, THEN $(1, 2)$ IS MORE THAN A LOCAL MINIMUM, IT A "PROPER" MINIMUM

$$\Rightarrow f(x) \geq 2 \quad \text{WHICH IMPLIES } \underline{x^4 + x^{-4} \geq 2}$$

$x \in \mathbb{R}, x > 0$

1YGB - MPI PAPER 2 - QUESTION 12

PROOF WITHOUT CALCULUS

START FROM THE FACT THAT ANY SQUARED
EXPRESSION IS NON NEGATIVE

$$\Rightarrow (x^4 - 1)^2 \geq 0$$

$$\Rightarrow x^8 - 2x^4 + 1 \geq 0$$

$$\Rightarrow x^8 + 1 \geq 2x^4$$

AS $x > 0$ WE MAY SAFELY DIVIDE THE INEQUALITY

$$\Rightarrow \frac{x^8 + 1}{x^4} \geq 2$$

$$\Rightarrow x^4 + x^{-4} \geq 2$$

AS REQUIRED