

# 1YGB - MP2 PART 1 - QUESTION 1

## USING APPROXIMATIONS FOR "SMALL ANGLES"

$$\begin{aligned}\sin \theta &\approx \theta \\ \cos \theta &\approx 1 - \frac{\theta^2}{2} \\ \cos\left(\frac{1}{2}\theta\right) &\approx 1 - \frac{\left(\frac{1}{2}\theta\right)^2}{2} \\ &\approx 1 - \frac{1}{8}\theta^2\end{aligned}$$

## HENCE WE HAVE

$$\begin{aligned}\Rightarrow \frac{\cos \frac{1}{2}x}{1 + \sin x} &= 0.925 \\ \Rightarrow \frac{1 - \frac{1}{8}x^2}{1 + x} &= 0.925 \\ \Rightarrow \frac{8 - x^2}{8 + 8x} &= 0.925 \\ \Rightarrow 7.4 + 7.4x &= 8 - x^2 \\ \Rightarrow x^2 + 7.4x - 0.6 &= 0\end{aligned}$$

## QUADRATIC FORMULA OR COMPLETING THE SQUARE

$$\begin{aligned}\Rightarrow (x + 3.7)^2 - 3.7^2 - 0.6 &= 0 \\ \Rightarrow (x + 3.7)^2 &= 14.29 \\ \Rightarrow x + 3.7 &= \begin{cases} 0.0802\dots\dots \\ -3.7802\dots\dots \end{cases} \\ \Rightarrow x &= \begin{cases} \cancel{7.4802\dots\dots} \\ 0.0802\dots\dots \end{cases}\end{aligned}$$

$\therefore x \approx 0.08$

1YGB - MP2 PAPER U - QUESTION 2

PROCEED AS FOLLOWS

$$\Rightarrow 2^2 + 4^2 + 8^2 + 16^2 + \dots = 1$$

$$\Rightarrow 2^2 + (2^2)^2 + (2^2)^3 + (2^2)^4 + \dots = 1$$

$$\Rightarrow 2^2 + (2^2)^2 + (2^2)^3 + (2^2)^4 + \dots = 1$$

THIS IS A CONVERGENT G.P WITH  $a = 2^2$  &  $r = 2^2$

$$\Rightarrow \frac{2^2}{1 - 2^2} = 1$$

$$S_{\infty} = \frac{a}{1 - r}$$

$$\Rightarrow 2^2 = 1 - 2^2$$

$$\Rightarrow 2 \times 2^2 = 1$$

$$\Rightarrow 2^2 = \frac{1}{2}$$

$$\Rightarrow 2^2 = 2^{-1}$$

$$\Rightarrow \underline{2 = -1}$$

# IYGB-MP2 PAPER U - QUESTION 3

a) SOLVING THE STANDARD EQUATION FOR  $0 \leq x \leq 2\pi$

$$2 + \sec\left(x - \frac{\pi}{3}\right) = 0$$

$$\sec\left(x - \frac{\pi}{3}\right) = -2$$


$$\cos\left(x - \frac{\pi}{3}\right) = -\frac{1}{2}$$

$$\begin{cases} x - \frac{\pi}{3} = 2\pi/3 \pm 2n\pi \\ x - \frac{\pi}{3} = 4\pi/3 \pm 2n\pi \end{cases} \quad n=0,1,2,3,\dots$$


$$\begin{cases} x = \pi \pm 2n\pi \\ x = \frac{5\pi}{3} \pm 2n\pi \end{cases}$$

$\therefore x = \pi, \frac{5\pi}{3}$

b) STARTING WITH THE GRAPH OF  $\sec x$




TRANSFORMING BY  $\pi/3$  TO THE "RIGHT" & BY 2 "UPWARDS"



$y = f(x) = 2 + \sec\left(x - \frac{\pi}{3}\right)$

1YGB - MP2 PAPER U - QUESTION 4

DRAWING A DIAGRAM



DETERMINE THE POSITION VECTOR OF D

$$\vec{OD} = \frac{1}{3} \vec{OC} = \frac{1}{3} (4, 0, 12) = \left(\frac{4}{3}, 0, 4\right) \text{ i.e. } D\left(\frac{4}{3}, 0, 4\right)$$

DETERMINE THE VECTORS  $\vec{AD}$  &  $\vec{DB}$

$$\vec{AD} = \underline{d} - \underline{a} = \left(\frac{4}{3}, 0, 4\right) - (1, 1, 4) = \left(\frac{1}{3}, -1, 0\right) = 1 \left(\frac{1}{3}, -1, 0\right)$$


$$\vec{DB} = \underline{b} - \underline{d} = (2, -2, 4) - \left(\frac{4}{3}, 0, 4\right) = \left(\frac{2}{3}, -2, 0\right) = 2 \left(\frac{1}{3}, -1, 0\right)$$

AS  $\vec{AD}$  IS IN THE SAME DIRECTION AS  $\vec{DB}$ , AND SHARE A POINT

A, D & B ARE COLLINEAR, SO D LIES ON THE LINE AB

YGB - MP2 PAPER 1 - QUESTION 5

a) SKETCHING EACH SECTION SEPARATELY NOTING THAT BOTH SECTIONS, AGREE AT  $x=2$




b) TREATING EACH SECTION SEPARATELY

●  $f_1(x) = x + 1$   
 $y = x + 1$   
 $x = y - 1$   
 $f_1^{-1}(x) = x - 1$

∴  $f^{-1}(x) = \begin{cases} x - 1 & x \leq 3 \\ 2 + \sqrt{x - 3} & x > 3 \end{cases}$

●  $f_2(x) = (x - 2)^2 + 3$   
 $y = (x - 2)^2 + 3$   
 $y - 3 = (x - 2)^2$   
 $x - 2 = \pm \sqrt{y - 3}$   
 $(x > 2 \Rightarrow \text{RHS IS POSITIVE})$   
 $x - 2 = \sqrt{y - 3}$   
 $x = 2 + \sqrt{y - 3}$   
 $f_2^{-1}(x) = 2 + \sqrt{x - 3}$



1YGB - MP2 PAPER 1 - QUESTION 6

● START BY REARRANGING THE EQUATION

$$4\sin\frac{\theta}{2} + \sqrt{3} = 0$$

$$4\sin\frac{\theta}{2} = -\sqrt{3}$$

$$\sin\frac{\theta}{2} = -\frac{\sqrt{3}}{4}$$

● WE NEED NO SOLUTION OVER AN ACTUAL RANGE, IF WE SUITABLY MANIPULATE THE FUNCTION

$$\Rightarrow f(\theta) = 4\cos\theta - 3\sin\frac{\theta}{2}$$

$$\Rightarrow f(\theta) = 4\left(1 - 2\sin^2\frac{\theta}{2}\right) - 3\sin\frac{\theta}{2}$$

$$\Rightarrow f(\theta) = 4 - 8\sin^2\frac{\theta}{2} - 3\sin\frac{\theta}{2}$$

$\cos 2A \equiv 1 - 2\sin^2 A$   
 $\cos 2\left(\frac{A}{2}\right) \equiv 1 - 2\sin^2\left(\frac{A}{2}\right)$   
 $\cos A \equiv 1 - 2\sin^2\frac{A}{2}$

● EVALUATING THE ABOVE AT  $\sin\frac{\theta}{2} = -\frac{\sqrt{3}}{4}$

$$\Rightarrow f(\theta) \Big|_{\sin\frac{\theta}{2} = -\frac{\sqrt{3}}{4}} = 4 - 8\left(-\frac{\sqrt{3}}{4}\right)^2 - 3\left(-\frac{\sqrt{3}}{4}\right)$$

$$= 4 - 8\left(\frac{3}{16}\right) + \frac{3\sqrt{3}}{4}$$

$$= 4 - \frac{3}{2} + \frac{3\sqrt{3}}{4}$$

$$= \frac{5}{2} + \frac{3\sqrt{3}}{4}$$

$$= \frac{10}{4} + \frac{3\sqrt{3}}{4}$$

$$= \frac{1}{4}(10 + 3\sqrt{3})$$

- 1 -

# 1 YGB - MP2 PAPER U - QUESTION 7

- REWRITE THE EQUATION IN INDEX FORM & DIFFERENTIATE BY USING THE PRODUCT RULE

$$\Rightarrow y = x(\ln x)^{\frac{1}{2}}$$

$$\Rightarrow \frac{dy}{dx} = 1 \times (\ln x)^{\frac{1}{2}} + \cancel{x} \times \frac{1}{2} (\ln x)^{-\frac{1}{2}} \times \frac{1}{\cancel{x}}$$

$$\Rightarrow \frac{dy}{dx} = (\ln x)^{\frac{1}{2}} + \frac{1}{2} (\ln x)^{-\frac{1}{2}}$$

$$\Rightarrow \frac{dy}{dx} = \sqrt{\ln x} + \frac{1}{2\sqrt{\ln x}}$$

- NOW WE REQUIRE GRADIENT OF  $\frac{3}{2}$

$$\Rightarrow \sqrt{\ln x} + \frac{1}{2\sqrt{\ln x}} = \frac{3}{2}$$

$$\Rightarrow a + \frac{1}{2a} = \frac{3}{2}$$

WHERE  $a = \sqrt{\ln x}$

$$\Rightarrow 2a^2 + 1 = 3a$$

$$\Rightarrow 2a^2 - 3a + 1 = 0$$

$$\Rightarrow (2a - 1)(a - 1) = 0$$

$$\Rightarrow a = \begin{cases} 1 \\ \frac{1}{2} \end{cases}$$

$$\Rightarrow \sqrt{\ln x} = \begin{cases} 1 \\ \frac{1}{2} \end{cases}$$

$$\Rightarrow \ln x = \begin{cases} 1 \\ \frac{1}{4} \end{cases}$$

-2-

IYGB - MP2 PAPER U - QUESTION 7

$$\Rightarrow x = \begin{cases} e^1 = e \\ e^{\frac{1}{4}} \end{cases}$$

$$\Rightarrow y = \begin{cases} e \times 1 = e \\ e^{\frac{1}{4}} \times \frac{1}{2} = \frac{1}{2}e^{\frac{1}{4}} \end{cases}$$

● REQUIRED POINTS ARE  $(e, e)$  &  $(e^{\frac{1}{4}}, \frac{1}{2}e^{\frac{1}{4}})$



- 1 -

1YGB - MP2 PAPER 1 - QUESTION 8

ARGUE AS FOLLOWS - LET  $a$  &  $b$  BE ODD POSITIVE INTEGERS

$a+b$  IS A MULTIPLE OF 4, so  $a+b = 4m, m \in \mathbb{N}$

SUPPOSE NOW  $a-b$  IS A MULTIPLE OF 4

$a-b = 4n, n \in \mathbb{N}$

ADDING THE EQUATIONS

$$\left. \begin{array}{l} a+b = 4m \\ a-b = 4n \end{array} \right\} \Rightarrow \begin{array}{l} 2a = 4(m+n) \\ a = 2(m+n) \end{array}$$

$\therefore a$  MUST BE EVEN

THIS IS A CONTRADICTION THAT  $a$  IS ODD

$\therefore a-b$  CANNOT BE A MULTIPLE OF 4

1YGB - MP2 PAPER 1 - QUESTION 9

a) REWRITE THE EQUATION IN FUNCTION FORM

$$x^2 = \frac{2}{\sqrt{x}} + \frac{3}{x^2}$$

$$x^2 - \frac{2}{\sqrt{x}} + \frac{3}{x^2} = 0$$

$$f(x) = x^2 - \frac{2}{\sqrt{x}} - \frac{3}{x^2}$$

- $f(1) = 1 - 2 - 3 = -4 < 0$
- $f(2) = 4 - \sqrt{2} - \frac{3}{4} = 1.8357... > 0$

AS  $f(x)$  IS CONTINUOUS ON  $(1,2)$ , AND CHANGES SIGN ON  $(1,2)$  THERE EXISTS AT LEAST A VALUE  $\alpha$  IN  $(1,2)$  SO THAT  $f(\alpha) = 0$

b)

$$f(x) = x^2 - 2x^{-\frac{1}{2}} - 3x^{-2}$$

$$f'(x) = 2x + x^{-\frac{3}{2}} + 6x^{-3}$$

NEWTON RAPHSON STATES

$$\Rightarrow x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

$$\Rightarrow x_{n+1} = x_n - \frac{x_n^2 - 2x_n^{-\frac{1}{2}} - 3x_n^{-2}}{2x_n + x_n^{-\frac{3}{2}} + 6x_n^{-3}}$$

$$\Rightarrow x_{n+1} = x_n - \frac{x_n^5 - 2x_n^{\frac{5}{2}} - 3x_n}{2x_n^4 + x_n^{\frac{3}{2}} + 6}$$

$$\Rightarrow x_{n+1} = \frac{2x_n^5 + x_n^{\frac{5}{2}} + 6x_n - x_n^5 + 2x_n^{\frac{5}{2}} + 3x_n}{2x_n^4 + x_n^{\frac{3}{2}} + 6}$$

$$\Rightarrow x_{n+1} = \frac{x_n^5 + 3x_n^{\frac{5}{2}} + 9x_n}{2x_n^4 + x_n^{\frac{3}{2}} + 6}$$

AS REQUIRED

1YGB - MP2 PAPER 1 - QUESTION 9

c)

USING ANY VALUE IN THE INTERVAL AND THE RECURRENT OF PART (b)

$$x_{n+1} = \frac{x_n^5 + 3x_n^{\frac{3}{2}} + 9x_n}{2x_n^4 + x_n^{\frac{3}{2}} + 6}$$

- $x_1 = 1.5$
- $x_2 = 1.634594485 \dots$
- $x_3 = 1.637565406 \dots$
- $x_4 = 1.637566228 \dots$
- $x_5 = 1.637566228 \dots$

∴ REQUIRED ROOT IS 1.63756623


CORRECT TO 8 D.P

YGB - MP2 PAGE 0 - QUESTION 10

EVIDENTLY THE AREA OF THE SQUARE ABCD IS  $4a^2$

THE RADIUS OF THE CIRCUMSCRIBING CIRCLE IS

$$\begin{aligned} \frac{1}{2} \sqrt{(2a)^2 + (2a)^2} &= \frac{1}{2} \sqrt{8a^2} \\ &= \frac{1}{2} \times 2\sqrt{2}a \\ &= \sqrt{2}a \end{aligned}$$



THE AREA BETWEEN THE SIDE OF THE SQUARE AND A WINK IS

$$\begin{aligned} \frac{1}{4} [\pi (\sqrt{2}a)^2 - 4a^2] &= \frac{1}{4} [\pi (2a^2) - 4a^2] = \frac{1}{2} [\pi a^2 - 2a^2] \\ &= \frac{1}{2} (\pi - 2) a^2 \end{aligned}$$

THE AREA OF EACH WINK IS

$$\frac{1}{2} \pi a^2 - \frac{1}{2} (\pi - 2) a^2 = \frac{1}{2} \pi a^2 - \frac{1}{2} \pi a^2 + a^2 = a^2$$

$\uparrow$   
 SQUARE

$\therefore$  AREA OF 4 WINKS IS  $4a^2$  WHICH IS THE SAME AREA AS THAT OF THE SQUARE

1YGB - MP2 PAPER U - QUESTION 11

PROCEED AS FOLLOWS

$$\begin{aligned} \frac{A+Bx}{(2-x)^3} &= (A+Bx)(2-x)^{-3} = (A+Bx) \times 2^{-3} (1-\frac{1}{2}x)^{-3} \\ &= \frac{1}{8}(A+Bx) \left[ 1 + \frac{-3}{1}(-\frac{1}{2}x) + \frac{-3(-4)}{1 \times 2}(-\frac{1}{2}x)^2 + \frac{-3(-4)(-5)}{1 \times 2 \times 3}(-\frac{1}{2}x)^3 + o(x^4) \right] \\ &= \frac{1}{8}(A+Bx) \left[ 1 + \frac{3}{2}x + \frac{3}{2}x^2 + \frac{5}{4}x^3 + o(x^4) \right] \\ &= (A+Bx) \left[ \frac{1}{8} + \frac{3}{16}x + \frac{3}{16}x^2 + \frac{5}{32}x^3 + o(x^4) \right] \\ &= \frac{1}{8}A + \frac{3}{16}Ax + \frac{3}{16}Ax^2 + \frac{5}{32}Ax^3 + o(x^4) \\ &\quad + \frac{1}{8}Bx + \frac{3}{16}Bx^2 + \frac{3}{16}Bx^3 + o(x^4) \\ &= \frac{1}{8}A + \left(\frac{3}{16}A + \frac{1}{8}B\right)x + \left(\frac{3}{16}A + \frac{3}{16}B\right)x^2 + \left(\frac{5}{32}A + \frac{3}{16}B\right)x^3 + o(x^4) \end{aligned}$$

$\begin{matrix} \uparrow & \uparrow & \uparrow & \uparrow \\ \frac{1}{4} & 0 & C & D \end{matrix}$

COMPARING COEFFICIENTS

- $\frac{1}{8}A = \frac{1}{4}$   
 $A = 2$
- $\frac{3}{16}A + \frac{1}{8}B = 0$   
 $3A + 2B = 0$   
 $6 + 2B = 0$   
 $B = -3$
- $C = \frac{3}{16}A + \frac{3}{16}B$   
 $C = \frac{3}{16}(A+B)$   
 $C = \frac{3}{16}(-1)$   
 $C = -\frac{3}{16}$
- $D = \frac{5}{32}A + \frac{3}{16}B$   
 $D = \frac{5}{16} - \frac{9}{16}$   
 $D = -\frac{1}{4}$

- 1 -

1YGB - M2 PAPER U - QUESTION 12

a) LOOKING AT THE FIRST SUMMATION WE REQUIRE THE FIRST 3 TERMS

$$t_{n+1} = at_n + 3n + 2$$

- $t_1 = -2$
- $t_2 = at_1 + 3 \times 1 + 2 = a(-2) + 3 + 2 = 5 - 2a$
- $t_3 = at_2 + 3 \times 2 + 2 = a(5 - 2a) + 6 + 2 = 8 + 5a - 2a^2$

NOW WE HAVE

$$\begin{aligned} \sum_{r=1}^3 (t_r + r^3) &= (t_1 + 1^3) + (t_2 + 2^3) + (t_3 + 3^3) \\ &= t_1 + 1 + t_2 + 8 + t_3 + 27 \\ &= t_1 + t_2 + t_3 + 36 \\ &= -2 + (5 - 2a) + (8 + 5a - 2a^2) + 36 \\ &= -2a^2 + 3a + 47 \end{aligned}$$

FINALLY WE HAVE

$$\begin{aligned} \Rightarrow \sum_{r=1}^3 (t_r + r^3) &= 12 \\ \Rightarrow -2a^2 + 3a + 47 &= 12 \\ \Rightarrow 0 &= 2a^2 - 3a - 35 \\ \Rightarrow (a - 5)(2a + 7) &= 0 \end{aligned}$$

$$a = \begin{cases} 5 \\ -\frac{7}{2} \end{cases}$$

1YGB - MP2 PAPER U - QUESTION 12

b) WE PROCEED AS FOLLOWS (THE VALUE OF k IS IRRELEVANT)

$$\sum_{r=8}^{31} [t_{r+1} - a.t_r] = \sum_{r=8}^{31} [(at_r + 3r + 2) - at_r]$$

$$= \sum_{r=8}^{31} (3r + 2)$$

$$= 26 + 29 + 32 + 35 + \dots + 95$$

THIS IS AN ARITHMETIC PROGRESSION WITH  $(31-7) = 24$  TERMS

$$\Rightarrow S_n = \frac{n}{2} [a + L]$$

$$\Rightarrow S_{24} = \frac{24}{2} [26 + 95]$$

$$\Rightarrow S_{24} = 12 \times 121$$

$$\Rightarrow S_{24} = \begin{array}{r} 1210 \\ 242 \end{array}$$

$$\Rightarrow S_{24} = 1452$$

# YGB - MP2 PAPER 1 - QUESTION 13

## FORMING A DIFFERENTIAL EQUATION

$x$  = COFFEE TEMPERATURE ( $^{\circ}\text{C}$ )  
 $t$  = TIME (MINUTES)  
-----  
 $t=0, x=80$   
 $t=10, x=40$

$$\frac{dx}{dt} = -k(x-10)^2$$

↑ RATE  
↑ PROPORTIONAL DECREASING  
↑ "DIFFERENCE ... SQUARED"

## SOLVING BY SEPARATION OF VARIABLES

$$\Rightarrow \frac{dx}{dt} = -k(x-10)^2$$
$$\Rightarrow dx = -k(x-10)^2 dt$$
$$\Rightarrow \frac{1}{(x-10)^2} dx = -k dt$$
$$\Rightarrow \int (x-10)^{-2} dx = \int -k dt$$
$$\Rightarrow -(x-10)^{-1} = -kt + C$$
$$\Rightarrow \frac{1}{x-10} = At + B$$

## APPLY THE CONDITIONS GIVEN

$$t=0, x=80 \Rightarrow \frac{1}{70} = B$$
$$\Rightarrow \frac{1}{x-10} = At + \frac{1}{70}$$
$$t=10, x=40 \Rightarrow \frac{1}{30} = 10A + \frac{1}{70}$$



1YGB - MP2 PAPER U - QUESTION 13

$$\Rightarrow A = \frac{1}{525}$$

$$\Rightarrow \frac{1}{x-10} = \frac{1}{525}t + \frac{1}{70}$$

$$\Rightarrow \frac{1050}{x-10} = 2t + 15$$

$$\Rightarrow \frac{1050}{2t+15} = x-10$$

$$\Rightarrow x = \frac{1050}{2t+15} + 10$$

$$\Rightarrow x = \frac{1050 + 10(2t+15)}{2t+15}$$

$$\Rightarrow x = \frac{20t + 200}{2t+15}$$

AS REQUIRED

FINALLY WITH  $x = 20$

$$\frac{1050}{2-10} = 2t + 15 \quad (\text{FROM EARLIER})$$

$$\frac{1050}{20-10} = 2t + 15$$

$$105 = 2t + 15$$

$$90 = 2t$$

$$t = 45$$

- 1 -

1YGB - MP2 PAPER 1 - QUESTION 14

a) 
$$\frac{1}{u^2 + 5u + 6} \equiv \frac{1}{(u+2)(u+3)} \equiv \frac{A}{u+2} + \frac{B}{u+3}$$

$$1 \equiv A(u+3) + B(u+2)$$

If  $u = -3 \Rightarrow 1 = -B$

If  $u = -2 \Rightarrow 1 = A$

$$\therefore \frac{1}{u^2 + 5u + 6} = \frac{1}{u+2} - \frac{1}{u+3}$$

b) 
$$\int_{\arcsin \frac{3}{5}}^{\arccos \frac{3}{5}} \frac{1}{(\sin x + 2\cos x)(\sin x + 3\cos x)} dx$$

DIVIDE "TOP & BOTTOM" OF THE INTEGRAND BY  $\cos^2 x$

$$= \int_{\arcsin \frac{3}{5}}^{\arccos \frac{3}{5}} \frac{\frac{1}{\cos^2 x}}{\frac{(\sin x + 2\cos x)(\sin x + 3\cos x)}{\cos x \cdot \cos x}} dx$$

$$= \int_{\arcsin \frac{3}{5}}^{\arccos \frac{3}{5}} \frac{\sec^2 x}{(\tan x + 2)(\tan x + 3)} dx$$


c) USING THE SUBSTITUTION GIVEN.

$u = \tan x$

THE UNITS BECOME

$\frac{du}{dx} = \sec^2 x$

$dx = \frac{du}{\sec^2 x}$



$\arccos \frac{3}{5} = \phi = \arctan \frac{4}{3}$

$\arcsin \frac{3}{5} = \theta = \arctan \frac{3}{4}$

•  $x = \arccos \frac{3}{5} \rightarrow u = \frac{4}{3}$

•  $x = \arcsin \frac{3}{5} \rightarrow u = \frac{3}{4}$

1YGB - MP2 PAPER 0 - QUESTION 14

HENCE WE NOW HAVE

$$\int_{\arcsin \frac{3}{5}}^{\arccos \frac{3}{5}} \frac{-\sec^2 x}{(\sec x + 2)(\tan x + 3)} dx = \int_{\frac{3}{4}}^{\frac{4}{3}} \frac{\cancel{\sec^2 x}}{(u+2)(u+3)} \frac{du}{\cancel{\sec^2 x}}$$
$$= \int_{\frac{3}{4}}^{\frac{4}{3}} \frac{1}{(u+2)(u+3)} du = \dots \text{part (a)} \dots = \int_{\frac{3}{4}}^{\frac{4}{3}} \frac{1}{u+2} - \frac{1}{u+3} du$$

$$= \left[ \ln|u+2| - \ln|u+3| \right]_{\frac{3}{4}}^{\frac{4}{3}} = \left( \ln \frac{10}{3} - \ln \frac{13}{3} \right) - \left( \ln \frac{11}{4} - \ln \frac{15}{4} \right)$$

$$= \ln \frac{10/3}{13/3} - \ln \frac{11/4}{15/4} = \ln \frac{10}{13} - \ln \frac{11}{15}$$

$$= \ln \frac{10/3}{11/15} = \ln \frac{150}{43}$$

# 1YGB - SPECIAL PAPER U - QUESTION 15

- START BY FINDING THE EQUATION OF THE TANGENT AT THE POINT WHERE  $t=p$  I.E  $P(p^2, p^2-p)$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2t-1}{2t}$$

$$\frac{dy}{dx}\bigg|_p = \frac{2p-1}{2p}$$

- EQUATION OF TANGENT IS GIVEN BY

$$y - (p^2 - p) = \frac{2p-1}{2p} (x - p^2)$$

- THE TANGENT PASSES THROUGH  $(4, \frac{3}{2})$

$$\frac{3}{2} - p^2 + p = \frac{2p-1}{2p} (4 - p^2)$$

$$3p - 2p^3 + 2p^2 = (2p-1)(4-p^2)$$

$$3p - \cancel{2p^3} + 2p^2 = 8p - \cancel{2p^3} - 4 + p^2$$

$$p^2 - 5p + 4 = 0$$


$$(p-1)(p-4) = 0$$

$$p = \begin{matrix} 1 \\ 4 \end{matrix}$$

- HENCE WE OBTAIN

$$p=1 \quad P(1, 0)$$

$$p=4 \quad P(16, 12)$$



## YGB - MP2 PAPER 1 - QUESTION 16

- DIFFERENTIATE THE EXPRESSION W.R.T  $x$ , USING THE FACT  $\frac{d}{dx}[a^{f(x)}] = a^{f(x)} \ln a \times f'(x)$

$$y = 2^{3e^{2x}} \Rightarrow \frac{dy}{dx} = 2^{3e^{2x}} \times \ln 2 \times 6e^{2x}$$

$$\Rightarrow \frac{dy}{dx} = y \ln 2 \times 2 \times (3e^{2x})$$

NOW WE NOTE THAT

$$\ln y = \ln 2^{3e^{2x}}$$

$$\ln y = (3e^{2x})(\ln 2)$$

$$\Rightarrow \frac{dy}{dx} = 2y \ln y$$

- ALTERNATIVE BY "TAKING LOGS" FIRST FOLLOWS BY IMPLICIT DIFFERENTIATION

$$\Rightarrow y = 2^{3e^{2x}}$$

$$\Rightarrow \ln y = \ln 2^{3e^{2x}}$$

$$\Rightarrow \ln y = (\ln 2)(3e^{2x})$$

DIFFERENTIATE WITH RESPECT TO  $x$

$$\Rightarrow \frac{1}{y} \frac{dy}{dx} = (\ln 2) \times 6e^{2x}$$

$$\Rightarrow \frac{dy}{dx} = y \times (\ln 2) \times 6e^{2x}$$

$$\Rightarrow \frac{dy}{dx} = 2y \times (\ln 2)(3e^{2x})$$

$$\Rightarrow \frac{dy}{dx} = 2y \ln y$$