Mun. My Mains Cloud. com

### FOR EDEXCEL

# GCE Examinations Advanced Subsidiary

## **Core Mathematics C2**

Paper L

Time: 1 hour 30 minutes

## Instructions and Information

Candidates may use any calculator EXCEPT those with the facility for symbolic algebra, differentiation and/or integration.

Full marks may be obtained for answers to ALL questions.

Mathematical formulae and statistical tables are available.

This paper has nine questions.

#### Advice to Candidates

You must show sufficient working to make your methods clear to an examiner. Answers without working may gain no credit.



Written by Shaun Armstrong
© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

- 1. A geometric series has first term 75 and second term -15.
  - Find the common ratio of the series.

**(2)** 

*(b)* Find the sum to infinity of the series. **(2)** 

2. A circle has the equation

$$x^2 + y^2 + 8x - 4y + k = 0,$$

where k is a constant.

Find the coordinates of the centre of the circle.

**(2)** 

Given that the x-axis is a tangent to the circle,

find the value of k. *(b)* 

**(3)** 

3.




Figure 1

Figure 1 shows a circle of radius r and centre O in which AD is a diameter.

The points B and C lie on the circle such that OB and OC are arcs of circles of radius r with centres A and D respectively.

Show that the area of the shaded region *OBC* is  $\frac{1}{6}r^2(3\sqrt{3} - \pi)$ . **(6)** 

- Sketch on the same diagram the graphs of  $y = \sin 2x$  and  $y = \tan \frac{x}{2}$  for x 4. *(a)* in the interval  $0 \le x \le 360^{\circ}$ .

Hence state how many solutions exist to the equation *(b)* 

$$\sin 2x = \tan \frac{x}{2},$$

for x in the interval  $0 \le x \le 360^{\circ}$  and give a reason for your answer.

**(2)** 

5. Find the value of a such that (a)

$$\log_a 27 = 3 + \log_a 8. {3}$$

*(b)* Solve the equation

$$2^{x+3} = 6^{x-1}$$
.

giving your answer to 3 significant figures.

**(4)** 

- Expand  $(2+x)^4$  in ascending powers of x, simplifying each coefficient. **6.** (a) **(4)** 
  - (b) Find the integers A, B and C such that

$$(2+x)^4 + (2-x)^4 \equiv A + Bx^2 + Cx^4.$$
 (2)

Find the real values of x for which (c)

$$(2+x)^4 + (2-x)^4 = 136.$$
 (3)

 $f(x) = 2x^3 - 5x^2 + x + 2$ 7.

- Show that (x 2) is a factor of f(x). (a) **(2)**
- *(b)* Fully factorise f(x). **(4)**
- Solve the equation f(x) = 0. (c) **(1)**
- Find the values of  $\theta$  in the interval  $0 \le \theta \le 2\pi$  for which (d)

$$2\sin^3\theta - 5\sin^2\theta + \sin\theta + 2 = 0,$$

**(4)** giving your answers in terms of  $\pi$ .

Turn over

**8.** The curve *C* has the equation

$$y = 3 - x^{\frac{1}{2}} - 2x^{-\frac{1}{2}}, \quad x > 0.$$

- (a) Find the coordinates of the points where C crosses the x-axis. (4)
- (b) Find the exact coordinates of the stationary point of C. (5)
- (c) Determine the nature of the stationary point. (2)
- (d) Sketch the curve C. (2)

9.




Figure 2

Figure 2 shows the curve C with equation  $y = 3x - 4\sqrt{x} + 2$  and the tangent to C at the point A.

Given that A has x-coordinate 4,

(a) show that the tangent to C at A has the equation 
$$y = 2x - 2$$
. (6)

The shaded region is bounded by C, the tangent to C at A and the positive coordinate axes.

(b) Find the area of the shaded region.

**(8)** 

**END**