

CORE MATHEMATICS (C) UNIT 1 TEST PAPER 2

1. Write down the exact values of

(i) $16^{\frac{3}{4}}$, (ii) $\left(\frac{2}{3}\right)^{-2}$, (iii) $4^{-\frac{1}{2}}$. [3]

2. (i) Find the set of real values of k for which the equation $x^2 - kx + k = 0$ has no real roots. [4]

(ii) State the number of real roots of the equation $x^2 - 4x + 4 = 0$. [2]

3. Solve the simultaneous equations $xy = 1$, $3x + 2y = 5$. [6]

4. The line l_1 passes through $(-2, 5)$ and is parallel to the line $4x + 3y = 0$.

The line l_2 also passes through $(-2, 5)$ and is perpendicular to the line $4x + 3y = 0$.

Find equations of l_1 and l_2 , expressing each answer in the form $ax + by + c = 0$ where a , b , c are integers. [7]

5. Two quantities P and t are related by the equation $P = \frac{k}{\sqrt{t}}$, where k is a positive constant.

(i) Find, in terms of k , the rate of change of P with t when $t = k^2$. [4]

Q also varies with t , such that $Q = \frac{\sqrt{t} + \sqrt{k}}{6}$.

(ii) Show that when $P = Q$, $t + \sqrt{k}\sqrt{t} - 6k = 0$. [3]

(iii) By substituting $t = x^2$, or otherwise, find, in terms of k , the value of t for which $P = Q$. [4]

6. In this question, $f(x) \equiv x^2 - 6x + 11$.

(i) Express $f(x)$ in the form $(x - p)^2 + q$ and hence find the minimum value of $f(x)$. [3]

(ii) If the line of symmetry of the graph of $y = f(x)$ has equation $x = a$, state the value of a . [2]

(iii) With this value of a , the graph of $y = f(x)$ is translated a units in the negative x -direction.

Find the equation of the resulting graph, giving your answer in a form without brackets. [3]

(iv) Sketch these two graphs on the same diagram and find the coordinates of the point where they intersect. [5]

CORE MATHEMATICS 1 (C) TEST PAPER 2 Page 2

7. The equation of a curve is $x^2y = x - 6$.

The normal to the curve at the point where $x = -2$ meets the x -axis at P .

(i) Find the coordinates of P . [9]

(ii) Find the coordinates of the point on the curve at which the gradient is 0. [4]

8. The points A and B have coordinates $(-2, 4)$ and $(6, -2)$ respectively.

(i) Find the coordinates of the mid-point of AB . [2]

(ii) Find the length of AB . [2]

(iii) Find the equation of the circle which has AB as a diameter, giving your answer in the form $x^2 + y^2 + ax + by + c = 0$ where a , b and c are integers. [3]

(iv) Find an equation of the tangent at A to the circle. [4]

The point P lies on the circle, and the straight line AP has gradient 2.

(v) State the gradient of BP . [2]

CORE MATHS 1 (C) TEST PAPER 2 : ANSWERS AND MARK SCHEME

1. (i) $2^3 = 8$ (ii) $(3/2)^2 = 9/4$ (iii) $1/\sqrt{4} = 1/2$ B1 B1 B1 3
2. (i) No real roots if $k^2 - 4k < 0$ $k(k-4) < 0$ $0 < k < 4$ B1 M1 A1 A1
(ii) Here $k = 4$, so there is one (repeated) real root M1 A1 6
3. $y = (5 - 3x)/2$ $x(5 - 3x) = 2$ $3x^2 - 5x + 2 = 0$ B1 M1 A1
 $(3x - 2)(x - 1) = 0$ $x = 2/3, y = 3/2$ or $x = 1, y = 1$ M1 A1 A1 6
4. Gradient of $l_1 = -4/3$, so l_1 is $y - 5 = -4/3(x + 2)$ $4x + 3y - 7 = 0$ M1 A1 A1
Gradient of $l_2 = 3/4$, so l_2 is $y - 5 = 3/4(x + 2)$ $3x - 4y + 26 = 0$ B1 M1 A1 A1 7
5. (i) $P = kt^{-1/2}$ $dP/dt = -1/2 kt^{-3/2}$ When $t = k^2$, $dP/dt = -1/(2k^2)$ B1 M1 A1 A1
(ii) When $P = Q$, $6k = t + \sqrt{kt}$ $t + \sqrt{k}\sqrt{t} - 6k = 0$ M1 A1 A1
(iii) $(\sqrt{t} + 3\sqrt{k})(\sqrt{t} - 2\sqrt{k}) = 0$ $\sqrt{t} > 0$ so $\sqrt{t} = 2\sqrt{k}$ $t = 4k$ M1 A1 M1 A1 11
6. (i) $f(x) = (x - 3)^2 + 2$ Minimum is $f(x) = 2$ M1 A1 A1
(ii) Line of symmetry is $x = 3$, so $a = 3$ M1 A1
(iii) New graph is $f(x + 3)$, so it is $y = x^2 + 2$ M1 A1 A1

(iv) Sketches				B1 B1	
Intersect where $x^2 - 6x + 11 = x^2 + 2$	$x = 3/2$	$(3/2, 17/4)$		M1 A1 A1	13
7. (i) $y = \frac{1}{x} - \frac{6}{x^2}$	$\frac{dy}{dx} = -\frac{1}{x^2} + \frac{12}{x^3} = -\frac{7}{4}$	when $x = -2$		B1 M1 A1 A1	
$y = -2$, so normal is $y + 2 = 4/7 (x + 2)$	When $y = 0$, $x = 3/2$			B1 M1 A1 A1 A1	
(ii) When $-\frac{1}{x^2} + \frac{12}{x^3} = 0$, $x = 12$	Point is $(12, 1/24)$			M1 A1 M1 A1	13
8. (i) Mid-point is $(2, 1)$				B1 B1	
(ii) $AB^2 = 64 + 36$	$AB = 10$			M1 A1	
(iii) $(x - 2)^2 + (y - 1)^2 = 25$	$x^2 + y^2 - 4x - 2y - 20 = 0$			M1 A1 A1	
(iv) Gradient of radius to $A = -3/4$, so gradient of tangent = $4/3$				B1 B1	
Tangent is $y - 4 = 4/3 (x + 2)$				M1 A1	
(v) Angle $ABP = 90^\circ$ (angle in semicircle), so gradient of $BP = -1/2$				M1 A1	13