

CORE MATHEMATICS (C) UNIT 2 TEST PAPER 3

1. Given that $4 \sin x = 3 \cos x$,
 - (i) find the value of $\tan x$. [1]
 - (ii) Find all the values of x between 0 and 360 for which $4 \sin x^\circ = 3 \cos x^\circ$, giving your answers to 1 decimal place. [3]

2. In the binomial expansion of $(k - 3x)^8$ in ascending powers of x , the first two terms are $a - 3072x$.
Find the values of the constants k and a . [6]

3. For the geometric series $a + ar + ar^2 + \dots$,
 - (i) give a formula for the n th term. [1]

The terms of this series are all different and positive. The sum of the first eight terms is 17 times the sum of the first four terms.

 - (ii) Find the value of r . [5]

4. Find the values of x for which $\log_3 (x^2 - 2x - 1) - \log_3 (x + 3) = -1$. [6]

5. At the point (x, y) on a curve C , the gradient is equal to $x^{\frac{1}{3}} + x^{-\frac{2}{3}}$.
The curve passes through the points $(8, 8)$ and $(27, a)$
 - (i) Find the equation of the curve. [5]
 - (ii) Find the value of a . [2]


6. (i) Find an expression, in terms of p , for the n th term of the sequence $\frac{1}{2}, p + \frac{1}{2}, 2p + \frac{1}{2}, 3p + \frac{1}{2}, \dots$ [3]
(ii) If the sum of the first ten terms of this sequence is 35, find the value of p . [4]

7. Given that $f(x) \equiv x^3 - 6x^2 + kx + 10$ and that $(x + 1)$ is a factor of $f(x)$,
 - (i) find the value of k . [2]
 - (ii) Write $f(x)$ as the product of three linear factors. [3]
 - (iii) Solve the equation $2^{3y} - 6(2^{2y}) + k(2^y) + 10 = 0$, giving solutions correct to 1 decimal place where necessary. [3]

CORE MATHEMATICS 2 (C) TEST PAPER 3 Page 2


8. (i) Use the trapezium rule, with four intervals of equal length, to estimate the area bounded by the curve $y = \frac{3}{2\sqrt{x}}$, the lines $x = 1$, $x = 5$ and the x -axis. Give your answer to 2 decimal places. [4]
- (ii) Use integration to find the true value of this area, also to 2 decimal places. [4]
- (iii) Show that the estimate lies within 1.5 % of the true value. [1]

9. In the diagram, a circle is divided into two sectors by the radii OA and OB . The areas of the major and minor sectors are in the ratio 3 : 2. The obtuse angle AOB is θ radians.



- (i) Show that $\theta = \frac{4\pi}{5}$. [4]
- (ii) Express, in its simplest form in terms of π , the ratio of the perimeters of the two sectors. [5]

10. The diagram shows part of the curves with equations $y = x^4 - 16$ and $y = x^2 + 12x + 20$.



- (i) Verify that the curves intersect at the points $(-2, 0)$ and $(3, 65)$. [2]
- (ii) Show that the area of the finite region between the curves is given by $\int_{-2}^3 (-x^4 + x^2 + 12x + 36) dx$. [2]
- (iii) Hence evaluate this area. [6]

CORE MATHS 2 (C) TEST PAPER 3 : ANSWERS AND MARK SCHEME

1. (i) $\tan x = \frac{3}{4}$ (ii) $x = 36.9, x = 216.9$ B1 M1 A1 A1 4
2. $k^8 + 8k^7(-3x) = a - 3072x$ $k^8 = a, 24k^7 = 3072$ M1 A1 A1
 $k = 128^{1/7} = 2$ $a = 2^8 = 256$ M1 A1 A1 6
3. (i) n th term $= ar^{n-1}$ B1
(ii) $a(r^8 - 1)/(r - 1) = 17a(r^4 - 1)/(r - 1)$ $r^8 - 1 = 17(r^4 - 1)$ M1 A1
 $r^8 - 17r^4 + 16 = 0$ $(r^4 - 1)(r^4 - 16) = 0$ $r > 0$ and $r \neq 1$ so $r = 2$ M1 A1 A1 6
4. $\log_3 [(x^2 - 2x - 1)/(x + 3)] = -1$ $(x^2 - 2x - 1)/(x + 3) = 1/3$ B1 M1
 $3x^2 - 7x - 6 = 0$ $(3x + 2)(x - 3) = 0$ $x = -2/3$ or $x = 3$ A1 M1 A1 A1 6
5. (i) Integrating, $y = \frac{3}{4}x^{4/3} + 3x^{1/3} + c$ $8 = 18 + c$, so $c = -10$ M1 A1 A1 M1 A1
(ii) $y = \frac{3}{4}x^{4/3} + 3x^{1/3} - 10$ $a = 60.75 - 1 = 59.75$ M1 A1 7
6. (i) Common difference $= p$, so $T_n = (n - 1)p + \frac{1}{2}$ B1 M1 A1
(ii) $S_n = \frac{n}{2}(1 + (n - 1)p) = 5(9p + 1) = 35$, so $p = \frac{2}{3}$ M1 A1 M1 A1 7
7. (i) $f(-1) = 0$ $-1 - 6 - k + 10 = 0$ $k = 3$ M1 A1
(ii) $f(x) = (x + 1)(x^2 - 7x + 10) = (x + 1)(x - 2)(x - 5)$ M1 A1 A1
(iii) $2^y = x = -1, 2, 5$ $y = 1$ or $y = \log_2 5 = 2.3$ M1 A1 A1 8
8. (i) $(1, 1.5), (2, 1.061), (3, 0.866), (4, 0.75), (5, 0.671)$ B2
Area $\approx \frac{1}{2}(2.171 + 2(2.677)) = 3.76$ M1 A1
(ii) $\int_1^5 \frac{3}{2}x^{-1/2} dx = [3x^{1/2}]_1^5 \approx 3.71$ M1 A1 M1 A1
(iii) $0.054/3.708 \times 100\% \approx 1.45\%$ B1 9
9. (i) $\frac{1}{2}r^2(2\pi - \theta) : \frac{1}{2}r^2\theta = 3 : 2$ $2(2\pi - \theta) = 3\theta$ $5\theta = 4\pi$ M1 A1 M1 A1
(ii) $r(2\pi - \theta) + 2r : \theta + 2r = (6\pi/5 + 2) : (4\pi/5 + 2) = (3\pi + 5) : (2\pi + 5)$ M1 A1 M1 A1 A1 9
10. (i) $16 - 16 = 4 - 24 + 20 = 0$, $81 - 16 = 9 + 36 + 20 = 65$ B1 B1
(ii) Area $= \int_{-2}^3 (x^2 + 12x + 20) - (x^4 - 16) dx = \int_{-2}^3 (-x^4 + x^2 + 12x + 36) dx$ M1 A1
(iii) $\left[-\frac{x^5}{5} + \frac{x^3}{3} + 6x^2 + 36x\right]_{-2}^3 = \left[-\frac{243}{5} + 9 + 54 + 108\right] - \left[\frac{32}{5} - \frac{8}{3} + 24 - 72\right]$ M1 A1 A1 M1 A1
 $= 500/3$ A1 10