

CORE MATHEMATICS (C) UNIT 2 TEST PAPER 5

1. Find, in radians, all the values of θ between 0 and 2π for which $\sin(2\pi - \theta) = \frac{1}{2}$. [4]

2. Given that $2^x = y$,
 (i) find, to 2 decimal places, the value of x when $y = 7$. [3]
 (ii) Express $\log_4 y$ in terms of x . [2]

3. Use the trapezium rule, with six equal intervals, to estimate the value of

$$\int_2^8 \log_{10} \left(\sin \frac{\pi}{x} \right) dx.$$

Give your answer to two decimal places. [6]

4. $f(x) \equiv x^3 + ax^2 + bx + 8$. When $f(x)$ is divided by $(x - 1)$, the remainder is 5.
 When $f(x)$ is divided by $(x + 2)$, the remainder is 20.
 (i) Find the values of the constants a and b . [4]
 (ii) Show that $(x + 4)$ is a factor of $f(x)$. [2]
 (iii) Find the number of real roots of the equation $f(x) = 0$. [2]

5. In the binomial expansion of $(1 + kx)^n$ in ascending powers of x , where n is a positive integer, the first three terms are $1 + 28x + 98kx^2$.
 Find the values of the constants k and n . [8]


6. The first three terms of an arithmetic series are $a(1 + b)$, $a(1 + 3b)$, $a(1 + 5b)$.
 (i) State the common difference of the series. [2]
 (ii) Find an expression for the n th term of the series. [2]
 (iii) If the fifth term of the series is 25 and the tenth term is 55, find the values of a and b . [5]

7. The function f is such that $f'(x) = (3x - 1)^2 - \frac{2}{x^2}$.
 (i) Given that $f(1) = 0$, find $f(x)$. [5]
 (ii) Find an equation of the tangent to the curve $y = f(x)$ at the point $(1, 0)$. [3]
 (iii) Find the second derivative of $f(x)$ with respect to x . [2]

CORE MATHEMATICS 2 (C) TEST PAPER 5 Page 2

8. The curve C has equation $y = \tan\left(x - \frac{\pi}{3}\right)$, where $-\pi \leq x \leq \pi$.
- Sketch C , showing the exact coordinates of all points of intersection with the coordinate axes and the equations of any asymptotes. [5]
 - Explain how the graph shows that for any real constant k there are exactly two values of x in the interval $-\pi < x \leq \pi$ for which $\tan\left(x - \frac{\pi}{3}\right) = k$. [2]
 - Find the solutions of $\tan\left(x - \frac{\pi}{3}\right) = 1$ in this interval, giving your answers in terms of π . [4]

9. The shaded segment is bounded by the arc AB and the chord AB of a circle with centre O and radius 4 cm. The angle AOB is 2θ radians. The arc AB is x cm longer than the chord AB .



- Show that $x = 8(\theta - \sin \theta)$. [4]
- Find an expression for the area of the shaded segment in terms of θ . [3]

Given further that the shaded area is $2x + 8(\sqrt{2} - 1) \text{ cm}^2$,

- show that $\sin 2\theta = 2 \sin \theta + 1 - \sqrt{2}$ [2]
- Verify that this equation is satisfied when $\theta = \frac{\pi}{4}$. [2]

CORE MATHS 2 (C) TEST PAPER 5 : ANSWERS AND MARK SCHEME

- $2\pi - \theta = \frac{\pi}{6}, \frac{5\pi}{6}$
 $\theta = \frac{7\pi}{6}, \frac{11\pi}{6}$

M1 M1 A1 A1 4
- (i) $x = \log 7 / \log 2 = 2.81$ (ii) $y = 2^x = 4^{x/2}$ so $\log_4 y = x/2$

M1 A1 A1 M1 A1 5
- (2, 0), (3, -0.062), (4, -0.151), (5, -0.231), (6, -0.301), (7, -0.363), (8, -0.417)

$\frac{1}{2}(1)(-0.417 + 2(-1.108)) = -1.32$

B1 B1 B1
M1 A1 A1 6
- (i) $f(1) = a + b + 9 = 5$ $f(-2) = 4a - 2b = 20$

$a + b = -4, 2a - b = 10$ $a = 2, b = -6$

(ii) $f(-4) = -64 + 32 + 24 + 8 = 0$ so $(x + 4)$ is a factor

(iii) $f(x) = (x + 4)(x^2 - 2x + 2)$ so one real root

B1
M1 A1 A1
M1 A1
M1 A1 8
- $(1 + kx)^n = 1 + nkx + \frac{n(n-1)}{2}k^2x^2 + \dots$ $nk = 28, \frac{1}{2}n(n-1)k^2 = 98k$

$n(n-1)k = 196$ $28(n-1) = 196$ $n = 8, k = 7/2$

M1 M1 A1 A1
M1 M1 A1 A1 8
- (i) Common difference $d = 2ab$

(ii) $T_n = a + ab + (n-1)(2ab)$ or $a(1 + (2n-1)b)$

(iii) $5d = 30$ so $ab = 3$ $a(1 + 9b) = 25$ so $a + 27 = 25$

$a = -2, b = -3/2$

M1 A1
M1 A1
B1 M1 A1
A1 A1 9
- (i) $f'(x) = 9x^2 - 6x + 1 - 2x^{-2}$ $f(x) = 3x^3 - 3x^2 + x + 2x^{-1} + c$

$3 - 3 + 1 + 2 + c = 0$, so $c = -3$ $f(x) = 3x^3 - 3x^2 + x - 3 + 2x^{-1}$

(ii) At (1, 0), $f'(x) = 2$ Tangent is $y = 2x - 2$

(iii) $f''(x) = 18x - 6 + 4x^{-3}$

B1 M1 A1
M1 A1
B1 M1 A1
B1 B1 10
- (i) Cuts axes at $(-2\pi/3, 0), (0, -\sqrt{3}), (\pi/3, 0)$ Asymp. $x = -\pi/6, x = 5\pi/6$

(ii) Any horizontal line cuts the graph twice in this interval

(iii) $x - \pi/3 = \pi/4, -3\pi/4$ $x = -5\pi/12, x = 7\pi/12$

B1 B2 B2
B2
M1 A1 A1 A1 11
- (i) Arc $AB = 8\theta$ Chord $AB = 2(4 \sin \theta)$ $8\theta - 8 \sin \theta = x$

(ii) Area = $\frac{1}{2}r^2(2\theta) - \frac{1}{2}r^2 \sin 2\theta = 16\theta - 8 \sin 2\theta$

(iii) $16\theta - 8 \sin 2\theta = 16\theta - 16 \sin \theta + 8(\sqrt{2} - 1)$ $\sin 2\theta = 2 \sin \theta + 1 - \sqrt{2}$

(iv) When $\theta = \pi/4$, $\sin 2\theta = 1$ $2 \sin \theta + 1 - \sqrt{2} = 2/\sqrt{2} + 1 - \sqrt{2} = 1$

B1 M1 A1 A1
M1 A1 A1
M1 A1
M1 A1 11