


MECHANICS (C) UNIT 2 TEST PAPER 7

Take $g = 9.8 \text{ ms}^{-2}$ and give all answers correct to 3 significant figures where necessary.

1. A particle of mass 0.6 kg moves in a horizontal circle with constant angular speed $1.5 \text{ radians per second}$. If the force directed towards the centre of the circle has magnitude 0.27 N , find the radius of the circular path. [3]



A key is modelled as a lamina which consists of a circle of radius 3 cm , with a circle of radius 1 cm removed from its centre, attached to a rectangle of length 8 cm and width 1 cm , with a rectangle measuring 3 cm by 1 cm fixed to its end as shown.

Calculate the distance of the centre of mass of the key from the line marked AB . [5]

3. A van of mass 1600 kg is moving with constant speed **down** a straight road inclined at 7° to the horizontal. The non-gravitational resistance to the van's motion has a constant magnitude of 2000 N and the engine of the van is working at a rate of 1.5 kW . Find

- (i) the constant speed of the van, [4]
 (ii) the acceleration of the van if the resistance is suddenly reduced to 1900 N . [2]

4. A body of mass 1 kg moves in a plane under the action of a constant force of magnitude $\sqrt{41} \text{ N}$. The body moves from the point P with coordinates $(-3, -15)$ to the point Q with position vector $(9, 0)$, the unit of distance on each axis being 1 metre .

- (i) Find the work done by the force in moving the body from P to Q . [4]
 (ii) Given that the body started from rest at P , find its speed when it is at Q . [3]

5. Two railway trucks A and B , whose masses are $6m$ and $5m$ respectively, are moving in the same direction along a straight track with speeds $5u$ and $3u$ respectively, and collide directly. Immediately after this impact, the speeds of A and B are v and kv respectively, in the same direction as before. The coefficient of restitution between A and B is e .

Modelling the trucks as particles,

- (i) show that (a) $v = \frac{45u}{5k+6}$, (b) $v = \frac{2eu}{k-1}$. [7]
 (ii) Use the fact that $0 < e < 1$ to deduce the range of possible values of k . [4]

6. A piece of lead and a table tennis ball are dropped together from a point P near the top of the Leaning Tower of Pisa. The lead hits the ground after 3.3 seconds .

- (i) Calculate the height above ground from which the lead was dropped. [2]

According to a simple model, the ball hits the ground at the same time as the lead.


- (ii) State why this may not be true in practice and describe a refinement to the model which could lead to a more realistic solution. [2]

The piece of lead is now thrown again from P , with speed 7 ms^{-1} at an angle of 30° to the horizontal.

- (iii) Find expressions in terms of t for x and y , the horizontal and vertical displacements respectively of the piece of lead from P at time $t \text{ seconds}$ after it is thrown. [4]

- (iv) Deduce that $y = \frac{\sqrt{3}x}{3} - \frac{2x^2}{15}$. [2]

- (v) Find the speed of the piece of lead when it has travelled 10 m horizontally from P . [4]



A uniform ladder AB , of mass m kg and length $2a$ m, rests with its upper end A in contact with a smooth vertical wall and its lower end B in contact with a fixed peg on horizontal ground. The ladder makes an angle α with the ground, where $\tan \alpha = \frac{3}{4}$.

- (i) Show that the magnitude of the resultant force acting on the ladder at B is $\frac{\sqrt{13}}{3}mg$. [6]
- (ii) Find, to the nearest degree, the direction of this resultant force at B . [2]
- The peg will break when the horizontal force acting on it exceeds $2mg$ N.
A painter of mass $6m$ kg starts to climb the ladder from B .
- (iii) Find, in terms of a , the greatest distance up the ladder that the painter can safely climb. [6]

MECHANICS 2 (C) TEST PAPER 7 : ANSWERS AND MARK SCHEME

1. $0.27 = mr\omega^2 = 0.6r(1.52)$ $r = 0.2$ m M1 A1 A1 3
2. $8\pi(3) + 8(10) + 3(13.5) = (11 + 8\pi)\bar{x}$ M1 A1
 $\bar{x} = (24\pi + 120.5) \div (8\pi + 11) = 5.42$ cm M1 A1 A1 5
3. (i) Net resisting force = $2000 - 1600g \sin 7^\circ = 89.1$ N M1 A1
 $1500 = 89.1v$ $v = 16.8$ ms⁻¹ M1 A1
- (ii) Now accelerating force = 100 N = $1600a$ $a = 0.0625$ ms⁻² M1 A1 6
4. (i) Displacement = $\sqrt{(12^2 + 15^2)} = 3\sqrt{41}$ M1 A1
 $F = 41$, so work done = $3\sqrt{41} \times \sqrt{41} = 123$ J M1 A1
- (ii) Work = change in K.E., so $\frac{1}{2}v^2 = 123$ $v = \sqrt{246} = 15.7$ ms⁻¹ M1 A1 A1 7
5. (i) (a) Momentum : $30mu + 15mu = 6mv + 5mkv$ M1 A1
 $45u = (6 + 5k)v$ $v = \frac{45u}{5k + 6}$ A1
- (b) Elasticity : $(kv - v) / (3u - 5u) = -e$ M1 A1
 $(k - 1)v = (-2u)(-e)$ $v = \frac{2eu}{k - 1}$ M1 A1
- (ii) $\frac{45u}{5k + 6} = \frac{2eu}{k - 1}$ $e = \frac{45(k - 1)}{2(5k + 6)}$ M1 A1
 $0 < e < 1$, so $0 < 45k - 45 < 10k + 12$ $k - 1 \neq 0$, so $1 < k < \frac{57}{35}$ M1 A1 11

6. (i) $s = \frac{1}{2}gt^2 = \frac{1}{2} \times 9.8 \times 3.32^2 = 53.4 \text{ m}$ M1 A1
(ii) Ball, being lighter, may be affected by air resistance : include this B1 B1
(iii) $x = (7 \cos 30^\circ)t = \frac{7\sqrt{3}}{2} t$ $y = (7 \sin 30^\circ)t - \frac{1}{2}gt^2 = \frac{7}{2}t - 4.9t^2$ M1 A1 M1 A1
(iv) $t = \frac{2x}{7\sqrt{3}}$ $y = \frac{x}{\sqrt{3}} - \frac{4.9(2x)^2}{(7\sqrt{3})^2} = \frac{\sqrt{3}x}{3} - \frac{2x^2}{15}$ M1 A1
(v) When $x = 10$, $y = -7.56$ B1
 $\frac{1}{2}m(49) + 7.56mg = \frac{1}{2}mv^2$ $v = 14.0 \text{ ms}^{-1}$ M1 A1 A1 14
7. (i) $R = mg, F = S$ M(B) : $mga \cos \alpha = 2a S \sin \alpha$ B1 M1 A1
 $S = mg / 2 \tan \alpha = \frac{2mg}{3} = F$ A1
Resultant force at B = $\sqrt{(mg)^2 + \left(\frac{2mg}{3}\right)^2} = \frac{\sqrt{13}mg}{3}$ M1 A1
(ii) Angle = $\tan^{-1} (2/3) = 33.7^\circ$ to horizontal M1 A1
(iii) M(B) : $mga \cos \alpha + 6mgx \cos \alpha = 2a S \sin \alpha$ M1 A1
 $S = \frac{2mg(a + 6x)}{3a}$ When $S = 2mg$, $a + 6x = 3a$ A1 M1 A1
 $6x = 2a$ $x = \frac{1}{3}a$ A1 14