

STATISTICS (C) UNIT 2 TEST PAPER 7

1. (i) Briefly explain the difference between a one-tailed test and a two-tailed test. [2]
(ii) State, with a reason, which type of test would be more appropriate to test the claim that this decade's average temperature is greater than the last decade's. [2]

2. A company that makes string wants to assess the breaking strain of its product.
 - (i) Explain why a sample, and not the whole population, should be used. [2]
A child cuts a 30 cm piece of string into two parts, cutting at a random point.
 - (ii) Find the probability that one part of the string is more than twice as long as the other. [2]
 - (iii) Sketch the probability density function of L , the length of the longer part of string. [2]

3. When a park is redeveloped, it is claimed that 70% of the local population approve of the new design. A conservation group, however, carries out a survey of 20 people, and finds that only 9 approve.
 - (i) Use this information to carry out a hypothesis test on the original claim, working at the 5% significance level. State your conclusion clearly. [5]
If the conservationists are right, and only 45% approve of the new park,
 - (ii) use a suitable approximation to the binomial distribution to estimate the probability that in a larger survey, of 500 people, less than half will approve. [6]

4. A certain type of steel is produced in a foundry. It has flaws (small bubbles) randomly distributed, and these can be detected by X-ray analysis. On average, there are 0.1 bubbles per cm^3 , and the number of bubbles per cm^3 has a Poisson distribution.
In an ingot of 40 cm^3 , find
 - (i) the probability that there are less than two bubbles, [3]
 - (ii) the probability that there are between 3 and 10 bubbles (inclusive). [3]A new machine is being considered. Its manufacturer claims that it produces fewer bubbles per cm^3 . In a sample ingot of 60 cm^3 , there are just two bubbles.
 - (iii) Carry out a hypothesis test at the 5% significance level to decide whether the new machine is better. State your hypotheses and conclusion carefully. [5]
 - (iv) Explain what a Type I error is in this context. [2]

5. The fraction of sky covered by cloud is modelled by the random variable X with probability density function

$$f(x) = kx(1 - x) \quad 0 \leq x \leq 1,$$

$$f(x) = 0 \quad \text{otherwise.}$$

(i) Find k and sketch the graph of $f(x)$. [4]

(ii) Find the mean and the standard deviation of X . [6]

(iii) Given that flying is prohibited when 81% of the sky is covered by cloud, show that cloud conditions allow flying nearly 90% of the time. [3]

6. In a particular parliamentary constituency, the percentage of Conservative voters at the last election was 35%, and the percentage who voted for the Monster Raving Loony party was 2%. Use suitable approximations to find

(i) the probability that a random sample of 500 electors will include at least 200 who voted either Conservative or Monster Raving Loony, [6]

(ii) the probability that a random sample of 200 electors will have at least 5 Monster Raving Loony voters in it. [5]

One of (i) or (ii) requires an adjustment to be made before a calculation is done. Explain what this adjustment is, and why it is necessary. [2]

STATISTICS 2 (C) TEST PAPER 7 : ANSWERS AND MARK SCHEME

1. (i) One-tailed : is a parameter greater (or less) than a given value? B1
Two-tailed : is a parameter different from a given value? B1
(ii) One-tailed, as testing for 'warmer' rather than 'different' B1 B1 4
2. (i) If every rope were tested to breaking point, none would be left B2
(ii) Needs to be cut in either of the 10 cm ends, so prob. = $\frac{2}{3}$ M1 A1
(iii) Graph drawn : $\frac{1}{15}$ for $15 \leq L \leq 30$, 0 elsewhere B2 6
3. (i) Taking $H_0 : p = 0.7$, no. approving is $X \sim B(20, p)$ B1 B1
Under H_0 , $P(X < 10) = P(X \leq 9) = 0.0171 < 5\%$ M1 A1
so at 5% level, reject H_0 and conclude that less than 70% approve A1
(ii) No. of approvals is $B(500, 0.45) \approx N(225, 123.75)$, so M1 A1
 $P(X < 250) = P(X < 249.5) = P(Z < 24.5/11.12)$ M1 A1
 $= P(Z < 2.20) = 0.986$ M1 A1 11

4. (i) $X \sim \text{Po}(4)$, so $P(X < 2) = 0.0916$ B1 M1 A1
(ii) $P(3 \leq X \leq 10) = 0.9972 - 0.2381 = 0.759$ M1 M1 A1
(iii) H_0 : mean number of bubbles is still 0.1 per cm^3 ;
 H_1 : mean < 0.1 B1
Under H_0 , no. of bubbles in 60 cm^3 is $\text{Po}(6)$ B1
Then $P(X \leq 2) = 0.062$, so do not reject H_0 at 5% level M1 A1 A1
(iv) Type I error is to reject the old machine in favour of the new,
when in fact it is no better B2 13
5. (i) Need $k \int x - x^2 \text{ dx} = 1 \quad k \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = 1 \quad k = 6$ M1 A1 A1
Graph sketched : parabola, vertex upwards, through $(0, 0)$, $(1, 0)$ B1
(ii) Mean = 0.5 , by symmetry M1 A1 A1
 $\text{Var}(X) = 6 \int x^3 - x^4 \text{ dx} - 0.5^2 = 6(0.25 - 0.2) - 0.25 = 0.05$ M1 A1
so standard deviation = $\sqrt{0.05} = 0.224$ A1
(iii) $P(x \leq 81\%) = 6 \int_0^{0.81} x^3 - x^4 \text{ dx} = 0.9054$, so cloud M1 A1
cover is $\leq 81\%$ for about 90% of the time A1 13
6. (i) No. of Cons or MRL $\sim \text{B}(500, 0.37) \approx \text{N}(185, 116.55)$, so M1 A1
 $P(X \geq 200) = P(X > 199.5) = P(Z > 14.5/10.79) = P(Z > 1.34)$ M1 A1 M1
 $= 1 - 0.9099 = 0.0901$ A1
(ii) No. of MRL $\sim \text{B}(200, 0.02) \approx \text{Po}(4)$ M1 A1
so $P(X \geq 5) = 1 - 0.6288 = 0.371$ M1 A1 A1
Binomial to Normal needs continuity correction, going from a discrete B1
to a continuous distribution B1 13