


## C2 Paper L – Marking Guide

1. (i)



B2

B2

(ii) 4 solutions

the graphs intersect at 4 points

B1

B1

(6)

2.

$$\text{area of segment} = \left(\frac{1}{2} \times r^2 \times \frac{\pi}{3}\right) - \left(\frac{1}{2} \times r^2 \times \sin \frac{\pi}{3}\right)$$

$$= \frac{1}{6} r^2 \pi - \frac{1}{4} r^2 \sqrt{3}$$

B1 M2

A1

$$\text{shaded area} = \frac{1}{6} r^2 \pi - 2\left(\frac{1}{6} r^2 \pi - \frac{1}{4} r^2 \sqrt{3}\right)$$

$$= \frac{1}{6} r^2 \pi - \frac{1}{3} r^2 \pi + \frac{1}{2} r^2 \sqrt{3}$$

$$= \frac{1}{2} r^2 \sqrt{3} - \frac{1}{6} r^2 \pi = \frac{1}{6} r^2 (3\sqrt{3} - \pi)$$

M1

A1

(6)

3.

$$(i) \quad u_2 = k^2 - 1$$

$$u_3 = (k^2 - 1)^2 - 1 = k^4 - 2k^2$$

B1

M1 A1

$$(ii) \quad k^4 - 2k^2 + k^2 - 1 = 11$$

M1

$$k^4 - k^2 - 12 = 0$$

M1

$$(k^2 + 3)(k^2 - 4) = 0$$

A1

$$k^2 = -3 \text{ (no solutions)} \text{ or } 4$$

A1

$$k = \pm 2$$

(7)

4.

(i)

$x$	0	0.5	1	1.5	2
$\frac{1}{x^2+1}$	1	0.8	0.5	0.3077	0.2

M1 A1

$$\text{area} \approx \frac{1}{2} \times 0.5 \times [1 + 0.2 + 2(0.8 + 0.5 + 0.3077)]$$

$$= 1.10 \text{ (3sf)}$$

B1 M1

A1

$$(ii) \quad \text{area} = 8^2 \times 1.10385 = 70.6464$$

M1

$$\text{volume} = 2 \times 70.6464 = 141 \text{ cm}^3 \text{ (3sf)}$$

A1

(7)

5.

$$(i) \quad \log_a 27 - \log_a 8 = 3$$

$$\log_a \frac{27}{8} = 3$$

M1

$$a^3 = \frac{27}{8}, \quad a = \sqrt[3]{\frac{27}{8}} = \frac{3}{2}$$

M1 A1

$$(ii) \quad (x+3) \lg 2 = (x-1) \lg 6$$

M1

$$x(\lg 6 - \lg 2) = 3 \lg 2 + \lg 6$$

M1

$$x = \frac{3 \lg 2 + \lg 6}{\lg 6 - \lg 2} = 3.52$$

M1 A1

(7)

6. (i)  $= [2x + x^{-1}]_2^4$  M1 A1  
 $= (8 + \frac{1}{4}) - (4 + \frac{1}{2}) = 3\frac{3}{4}$  M1 A1

(ii)  $y = \int (2x^3 + 1) dx$   
 $y = \frac{1}{2}x^4 + x + c$  M1 A1  
 $x = 0, y = 3 \therefore c = 3$  B1  
 $y = \frac{1}{2}x^4 + x + 3$   
when  $x = 2, y = 8 + 2 + 3 = 13$  M1 A1 (9)

---

7. (i)  $\frac{1-8x^3}{x^2} = 0 \Rightarrow 1-8x^3 = 0$  M1  
 $x^3 = \frac{1}{8}$  M1  
 $x = \frac{1}{2}$  A1

(ii)  $f(x) = x^{-2} - 8x$   
 $\int f(x) dx = \int (x^{-2} - 8x) dx$   
 $= -x^{-1} - 4x^2 + c$  M1 A2

(iii)  $= -[-x^{-1} - 4x^2]_{\frac{1}{2}}^2$  M1  
 $= -\{(-\frac{1}{2} - 16) - (-2 - 1)\} = 13\frac{1}{2}$  M1 A1 (9)

---

8. (i)  $S_6 = \frac{6}{2}[3000 + (5 \times -x)] = 8100$  M1 A1  
 $3000 - 5x = 2700, x = 60$  M1 A1

(ii)  $= 1500 - (7 \times 60) = 1500 - 420 = £1080$  M1 A1

(iii)  $S_n = \frac{n}{2}[3000 - 60(n - 1)]$  M1  
 $= n[1500 - 30(n - 1)]$   
 $= 30n[50 - (n - 1)] = 30n(51 - n) \quad [k = 30]$  M1 A1

(iv) the value of sales in a month would become negative  
which is not possible B1 (10)

---

9. (i)  $f(2) = 16 - 20 + 2 + 2 = 0 \therefore (x - 2)$  is a factor M1 A1

(ii) 
$$\begin{array}{r} 2x^2 - x - 1 \\ x - 2 \overline{) 2x^3 - 5x^2 + x + 2} \\ 2x^3 - 4x^2 \\ \hline -x^2 + x \\ -x^2 + 2x \\ \hline -x + 2 \\ -x + 2 \\ \hline \end{array}$$
 M1 A1

$f(x) = (x - 2)(2x^2 - x - 1)$   
 $f(x) = (x - 2)(2x + 1)(x - 1)$  M1 A1

(iii)  $x = -\frac{1}{2}, 1, 2$  B1

(iv)  $\sin \theta = 2$  (no solutions),  $-\frac{1}{2}$  or 1  
 $\theta = \pi + \frac{\pi}{6}, 2\pi - \frac{\pi}{6}$  or  $\frac{\pi}{2}$  M1 B1  
 $\theta = \frac{\pi}{2}, \frac{7\pi}{6}, \frac{11\pi}{6}$  A2 (11)

---

Total (72)